

7.23 P

Evaluation of Thermotherapy against Huanglongbing (Citrus Greening) under Laboratory Conditions

Fan, G.¹, Xia, Y.², Lin, X.¹, Cai, Z.¹, Hu, H.¹, Wang, X.¹, Ruan, C.³, Lu, L.⁴, Sequeira, R.⁵, and Liu, B.³

¹Fruit Research Institute, Fujian Academy of Agricultural Sciences, Fuzhou, Fujian, China, 350013

²NSF Center for Integrated Pest Management, North Carolina State University, Raleigh, NC 27606, USA

³Agricultural Bio-Resources Research Institute, Fujian Academy of Agricultural Sciences, Fuzhou, Fujian, China, 350003

⁴Zhejiang Citrus Research Institute, Taizhou, Zhejiang, China, 318020

⁵The United States Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine, Center for Plant Health Science and Technology, Raleigh, NC 27606, USA

Huanglongbing (HLB, citrus greening) is the most destructive disease of citrus. The disease is associated with “*Candidatus Liberibacter asiaticus*”. Few management options are available, besides preventive measures such as the removal of affected plants, planting disease-free stock and maintaining vector-free production in quarantine areas. In this study, we assessed the efficacy of thermotherapy against the disease under controlled laboratory conditions. A total of sixty, 2-year old graft-infected *Citrus reticulata* Blanco seedlings were used for the study. The plants were randomly divided into 3 treatment groups (45, 48°C, and untreated), with 5 plants/rep, 4 reps/trt. The treated plants were placed in temperature chambers for a 4-h treatment sessions, repeated weekly 3 times. Disease remission was observed beginning 8 weeks post-treatment. Real-time PCR assays revealed that pathogen “*Ca. L. asiaticus*” concentration of all HLB-affected seedlings were significantly reduced except for 8 plants under 45 and 48°C treatments at 4 weeks after treatment. In contrast, pathogen concentration in the untreated control plants exhibited a significant increase, with the highest increase of about 30-fold compared to the initial pathogen concentration (pre-treatment). Except for 7 plants (7 out of 40 total plants), pathogen concentration in the new flushes of the treated plants decreased 90% at ca. 8 weeks after treatment, compared to the initial pathogen concentration. Nested and real-time PCR were used for confirmation of HLB infection in the seedlings, and for pathogen titer assessment. Although the result is considered preliminary, it provides a foundation for further work in developing the technique for HLB management. Complementary work will include exploration of additional exposure time and temperature combinations as well as treatments using commercial field settings.