

CLINICAL VIGNETTE

A Case of Severe Pernicious Anemia and Subacute Combined Degeneration Associated with Recreational Nitrous Oxide Use

Rintu Sajū, BS¹, Natalie Bransky, MD², Stanley Yuan, MD³ and Joseph Zaky, MD⁴

¹David Geffen School of Medicine at UCLA, Los Angeles, CA

²Department of Medicine, UCLA Medical Center, Los Angeles, CA

³Department of Medicine, Cedars-Sinai Medical Center, Los Angeles, CA

⁴Department of Medicine, Olive View Medical Center, Sylmar, CA

Key Words: vitamin B12 deficiency, nitrous oxide use, subacute combined degeneration of the spinal cord, autoimmune gastritis

Case

A 44-year-old male with history of polysubstance use with nitrous oxide, tobacco, marijuana, and cocaine presented to the emergency department with diffuse paresthesias, abdominal bloating, and dyspnea.

Four weeks prior, the patient reported onset of progressive numbness and paresthesias of all four extremities and trunk along with diffuse weakness that limited ambulation and activities of daily living. His family reported confusion with poor concentration and problem-solving skills. The patient also reported constipation with abdominal bloating and dyspnea also over the past month. He denied headaches, fevers, nausea, vomiting, diarrhea, or urinary or bowel incontinence. He did not follow a restrictive diet and had no recent travel or sick contacts. Family history was notable for pernicious anemia on the paternal side. Social history included polysubstance use with recreational nitrous oxide for the last two years, using up to 24 canisters a day for the past two months prior to admission. He also used tobacco daily, and intermittent cocaine, marijuana, and alcohol.

On admission, the patient was afebrile and hemodynamically stable. Physical exam was notable for diminished motor strength (3/5) globally, decreased fine touch and vibratory sensation, decreased proprioception, ataxic gait, and moderate abdominal distension with tympany. Labs included normal Hgb, WBC and platelets on admission but developed mild normocytic anemia on hospital day 3 with hemoglobin of 12.3 g/dL and hematocrit of 37.1%. Vitamin B12 level was decreased at 146 pg/mL (reference of 213 to 816) with elevated homocysteine of 171.9 umol/L (reference <11.4), elevated methylmalonic acid of 44620 nmol/L (reference of 87-318), and positive intrinsic factor antibody. Parietal cell antibody was negative. Metabolic panel, iron studies, folate, vitamin B1, vitamin B6, and TSH. HIV and RPR were also normal. Peripheral blood smear was notable for occasional hypersegmented neutrophils.

MRI brain was unremarkable, however, MRI spine demonstrated degenerative disc disease of the cervical spine from C5-7 and thickening of the ligamentum flavum, moderate spinal canal narrowing with effacement of the CSF surrounding the cord and minimal ventral cord flattening. CT of the abdomen and pelvis was negative.

The patient had severe vitamin B12 deficiency complicated by subacute combined degeneration of the spinal cord. His genetic predisposition with the positive intrinsic factor antibody suggested autoimmune gastritis, which along with extensive nitrous oxide use contributed to his Vitamin B12 deficiency, which led to his acute clinical and functional decline.

The patient was started on daily intramuscular (IM) vitamin B12 1000mcg supplementation while in the hospital with planned taper to weekly followed by monthly IM supplementation after discharge. He received inpatient physical and occupational therapy to improve mobility and strength. On day 7 of hospitalization, patient's vitamin B12 level normalized, and he was discharged to an acute rehabilitation center. Neuropathic pain persisted at the rehabilitation center despite different medical therapies. Transcutaneous electric nerve stimulation provided significant benefits. He will schedule outpatient endoscopies given increased risk/ incidence of GI cancers among patients with pernicious anemia.

Discussion

Vitamin B12, also referred to as cobalamin, is a critical cofactor necessary for cell maturation and neuronal development. Deficiencies can result in a myriad of hematologic and neurodegenerative complications. Fortunately, the clinical manifestations of vitamin B12 deficiency can be reversible, making early diagnosis and treatment important.¹

Cobalamin is derived primarily from animal food sources such as meats, eggs, fish, shellfish, and dairy products.^{2,3} Once con-

sumed, Vitamin B12 is freed as a micronutrient by stomach acid and binds with salivary R-binder in the gastric lumen.³ In the duodenum, pancreatic protease dissociates vitamin B12, which then binds with intrinsic factor (produced by gastric parietal cells) and is absorbed in the terminal ileum, with the assistance of specialized “cubam” receptors.^{3,4} Disruption anywhere along this pathway can result in low serum B12 levels and associated clinical manifestations. Those following a strict vegan or strict vegetarian diets are at increased risk of developing vitamin B12 deficiency due to decreased dietary intake of vitamin B12.⁵ Reduced absorption due to *H. Pylori* infection, use of medications such as metformin or proton-pump inhibitors, pancreatic insufficiency, small bowel pathology, and surgical changes to the gastrointestinal tract have all been implicated in B12 deficiency.^{1,5–7} However, autoimmune atrophic gastritis, also known as pernicious anemia, remains the most common cause of B12 deficiency with reported prevalence ranging from 50 to 4000 cases per 100,000 persons.^{1,8} Pernicious anemia results from antibodies that target H⁺/K⁺ ATPase in the gastric parietal cells and/or intrinsic factor, leading to atrophic gastritis and poor B12 absorption.⁹ Theories have also explored the possible role of *H. pylori* in the development of autoimmune gastritis through molecular mimicry, although evidence remains equivocal.^{10–12}

At the cellular level, nitrous oxide, a colorless gas colloquially known as “whippets” or “laughing gas”, is a potent inhibitor of vitamin B12 and can precipitate deficiencies and associated neurologic disease.^{13–16} In clinical medicine, nitrous oxide is commonly used as an anesthetic, especially in obstetrics and dentistry, and has received increased recognition in treating refractory depression.^{17,18} However, with its ability to produce instantaneous physiologic and psychedelic effects (e.g., euphoria, analgesia, depersonalization/ derealization), recreational nitrous oxide is gaining popularity, with reported lifetime prevalence as high as 29% in the United States in 2014.^{19,20} Nitrous oxide exhibits deleterious effects by oxidizing vitamin B12, changing its valency, and rendering it inactive.¹⁹ Deficiencies and associated clinical manifestations can be seen after prolonged or even single use, especially among those with underlying susceptibility for vitamin B12 deficiencies.^{14,15,21}

Vitamin B12 is used as a cofactor for two major biochemical reactions involving methionine synthase and L-methyl malonyl-coenzyme A mutase.¹ In the former reaction, B12 acts as the integral cofactor in the recycling of methyl-tetrahydrofolate to tetrahydrofolate (THF), while simultaneously coupling it to generation of methionine from homocysteine.¹ Generation of THF is necessary for purine and pyrimidine synthesis, and disruption to this process leads to improper DNA and RNA synthesis. The subsequent nuclear-cytoplasmic dyssynchrony most notably affects rapidly dividing cells such as the hematopoietic cell lines. This classically results in isolated anemia with macrocytosis and/or megaloblastic on peripheral blood smears.⁶

High levels of homocysteine are seen in B12 deficiency and are often used as a marker to diagnose deficiency, although ele-

vated levels are not specific to Vitamin B12 deficiency. Elevated levels of methylmalonic acid (used as a proxy for L-methylmalonyl CoA) have a higher specificity for diagnosing cobalamin deficiency. However, isolated elevations of methylmalonic acid have been described in the context of renal insufficiency.^{1,3,6}

Subacute combined degeneration (SACD) of the spinal cord is a potentially devastating complication of vitamin B12 deficiency. While the exact mechanism is unclear, evidence suggests that elevated levels of methylmalonyl CoA and associated byproducts like propionyl CoA leads to improper myelin synthesis and aberrant fatty acid accumulation.²² Lack of methionine, a critical precursor for neuronal sheath stability, is also thought to play a role in the demyelinating manifestations of cobalamin deficiency. Demyelination classically affects the dorsal column, lateral corticospinal tract, and spinocerebellar tract leading to decrease proprioception and vibratory sensation, symmetric paresthesias, muscle weakness, hyperreflexia, spasticity, and sensory ataxia, which were seen in our patient.^{1,22} Diagnosis can include MRI of the spine, which classically appears as the “inverted V” sign, alluding to the symmetric linear T2-hyperintensity of the dorsal column seen in the cervical and thoracic spine on imaging.²³ However, the absence of MRI findings, as was the case in our patient, does not completely rule out SACD of the spine.²⁴ Some studies report lower sensitivity of MRI findings in diagnosing SACD, underscoring the importance of considering SACD in the right clinical context even in the absence of classic MRI findings.^{24,25} Notably, MRI in our patient did show moderate disc disease and the effacement of CSF adjacent to the cord and ventral cord flattening, which could also indicate underlying spondylotic myelopathy, in part contributing to some of his neurologic findings.

Although described as pernicious “anemia”, frank anemia is often a later manifestation of disease.⁶ Many patients, like ours, present first with neuropsychiatric manifestations of B12 deficiency, which can make diagnosis challenging.^{26,27} Such presentations may develop with normal serum B12 level,^{28–30} as serum levels may not be reflective of cellular level deficiencies. Diagnosis should therefore be made in the context of metabolite levels (e.g., homocysteine and methylmalonic acid) and correlated to the patient’s clinical presentation.⁶ Recreational nitrous oxide is a risk factor for B12 deficiencies and should be screened for in the right clinical context.

Treatment for pernicious anemia involves lifelong supplementation, usually given intramuscularly,⁹ although recent data suggest that high dose oral supplementation might be equally effective.^{31–33} Gastroenterology should also be consulted for thorough evaluation. Patients may need to undergo an upper endoscopy, which is the gold standard in diagnosing autoimmune gastritis but can also evaluate for GI malignancies given increased risk among this population.³⁴

Conclusion

Pernicious anemia is a common cause of vitamin B12 deficiency.¹ However, frank anemia is usually a later manifestation of this disease process, and patients are often asymptomatic or present first with neuropsychiatric manifestations, as was the case with our patient.⁶ The patient's family history of pernicious anemia along with the presence of intrinsic factor antibodies suggested a genetic susceptibility for an underlying autoimmune process. His daily nitrous oxide use likely contributed to his B12 deficiency and rapid functional decline.

Paresthesias and muscle weakness can have significant impact on a patient's functional status. A multidisciplinary approach involving vitamin supplementation along with intensive physical and occupational therapy should be offered to improve a patient's overall functional status. Screening for inhalant drug use is often missed and with it, potential opportunities for adequate counseling for patients. Meetings with a substance use counselor may be beneficial to address any underlying substance use (e.g., nitrous oxide use) contributing to B12 deficiency.

REFERENCES

1. **Stabler SP.** Clinical practice. Vitamin B12 deficiency. *N Engl J Med.* 2013 Jan 10;368(2):149-60. doi: 10.1056/NEJMc1113996. PMID: 23301732.
2. **Allen LH.** Vitamin B-12. *Adv Nutr.* 2012 Jan;3(1):54-5. doi: 10.3945/an.111.001370. Epub 2012 Jan 5. PMID: 22332101; PMCID: PMC3262614.
3. **Zemleni J, Suttie JW, Gregory III JF, Stover PJ (eds.).** *Handbook of Vitamins.* Boca Raton: CRC Press; 2013. 605 p.
4. **Fyfe JC, Madsen M, Højrup P, Christensen EI, Tanner SM, de la Chapelle A, He Q, Moestrup SK.** The functional cobalamin (vitamin B12)-intrinsic factor receptor is a novel complex of cubilin and amnionless. *Blood.* 2004 Mar 1;103(5):1573-9. doi: 10.1182/blood-2003-08-2852. Epub 2003 Oct 23. PMID: 14576052.
5. **Allen LH.** Causes of vitamin B12 and folate deficiency. *Food Nutr Bull.* 2008 Jun;29(2 Suppl):S20-34; discussion S35-7. doi: 10.1177/15648265080292S105. PMID: 18709879.
6. **Green R.** Vitamin B12 deficiency from the perspective of a practicing hematologist. *Blood.* 2017 May 11;129(19):2603-2611. doi: 10.1182/blood-2016-10-569186. Epub 2017 Mar 30. PMID: 28360040.
7. **de Jager J, Kooy A, Leher P, Wulffelé MG, van der Kolk J, Bets D, Verburg J, Donker AJ, Stehouwer CD.** Long term treatment with metformin in patients with type 2 diabetes and risk of vitamin B-12 deficiency: randomised placebo controlled trial. *BMJ.* 2010 May 20;340:c2181. doi: 10.1136/bmj.c2181. PMID: 20488910; PMCID: PMC2874129.
8. **Stabler SP, Allen RH.** Vitamin B12 deficiency as a worldwide problem. *Annu Rev Nutr.* 2004;24:299-326. doi: 10.1146/annurev.nutr.24.012003.132440. PMID: 15189123.
9. **Bizzaro N, Antico A.** Diagnosis and classification of pernicious anemia. *Autoimmun Rev.* 2014 Apr-May;13(4-5):565-8. doi: 10.1016/j.autrev.2014.01.042. Epub 2014 Jan 11. PMID: 24424200.
10. **Kaptan K, Beyan C, Ural AU, Cetin T, Avcu F, Gülşen M, Finci R, Yalçın A.** Helicobacter pylori--is it a novel causative agent in Vitamin B12 deficiency? *Arch Intern Med.* 2000 May 8;160(9):1349-53. doi: 10.1001/archinte.160.9.1349. PMID: 10809040.
11. **Appelmelk BJ, Simoons-Smit I, Negrini R, Moran AP, Aspinall GO, Forte JG, De Vries T, Quan H, Verboom T, Maaskant JJ, Ghiara P, Kuipers EJ, Bloemena E, Tadema TM, Townsend RR, Tyagarajan K, Crothers JM Jr, Monteiro MA, Savio A, De Graaff J.** Potential role of molecular mimicry between Helicobacter pylori lipopolysaccharide and host Lewis blood group antigens in autoimmunity. *Infect Immun.* 1996 Jun;64(6):2031-40. doi: 10.1128/iai.64.6.2031-2040.1996. PMID: 8675304; PMCID: PMC174033.
12. **Lewerin C, Jacobsson S, Lindstedt G, Nilsson-Ehle H.** Serum biomarkers for atrophic gastritis and antibodies against Helicobacter pylori in the elderly: Implications for vitamin B12, folic acid and iron status and response to oral vitamin therapy. *Scand J Gastroenterol.* 2008;43(9):1050-6. doi: 10.1080/00365520802078341. PMID: 18609169.
13. **Miller MA, Martinez V, McCarthy R, Patel MM.** Nitrous oxide "whippit" abuse presenting as clinical B12 deficiency and ataxia. *Am J Emerg Med.* 2004 Mar;22(2):124. doi: 10.1016/j.ajem.2003.12.015. PMID: 15011232.
14. **Campdesuner V, Teklie Y, Alkayali T, Pierce D, George J.** Nitrous Oxide-Induced Vitamin B12 Deficiency Resulting in Myelopathy. *Cureus.* 2020 Jul 9;12(7):e9088. doi: 10.7759/cureus.9088. PMID: 32685323; PMCID: PMC7366039.
15. **McArdle DJT, Gaillard F.** Pernicious azotaemia? A case series of subacute combined degeneration of the cord secondary to nitrous oxide abuse. *J Clin Neurosci.* 2020 Feb;72:277-280. doi: 10.1016/j.jocn.2019.11.003. Epub 2020 Jan 11. PMID: 31937498.
16. **Irwin MG, Trinh T, Yao CL.** Occupational exposure to anaesthetic gases: a role for TIVA. *Expert Opin Drug Saf.* 2009 Jul;8(4):473-83. doi: 10.1517/14740330903003778. PMID: 19480607.
17. **Xiang Y, Li L, Ma X, Li S, Xue Y, Yan P, Chen M, Wu J.** Recreational Nitrous Oxide Abuse: Prevalence, Neurotoxicity, and Treatment. *Neurotox Res.* 2021 Jun;39(3):975-985. doi: 10.1007/s12640-021-00352-y. Epub 2021 Mar 26. PMID: 33770366.
18. **Nagele P, Duma A, Kopec M, Gebara MA, Parsoei A, Walker M, Janski A, Panagopoulos VN, Cristancho P, Miller JP, Zorumski CF, Conway CR.** Nitrous Oxide for Treatment-Resistant Major Depression: A Proof-of-Concept Trial. *Biol Psychiatry.* 2015 Jul 1;78(1):10-18.

doi: 10.1016/j.biopsych.2014.11.016. Epub 2014 Dec 9. PMID: 25577164.

19. **Garakani A, Jaffe RJ, Savla D, Welch AK, Protin CA, Bryson EO, McDowell DM.** Neurologic, psychiatric, and other medical manifestations of nitrous oxide abuse: A systematic review of the case literature. *Am J Addict.* 2016 Aug;25(5):358-69. doi: 10.1111/ajad.12372. Epub 2016 Apr 1. PMID: 27037733.
20. **Kaar SJ, Ferris J, Waldron J, Devaney M, Ramsey J, Winstock AR.** Up: The rise of nitrous oxide abuse. An international survey of contemporary nitrous oxide use. *J Psychopharmacol.* 2016 Apr;30(4):395-401. doi: 10.1177/0269881116632375. Epub 2016 Feb 24. PMID: 26912510.
21. **Hadzic A, Glab K, Sanborn KV, Thys DM.** Severe neurologic deficit after nitrous oxide anesthesia. *Anesthesiology.* 1995 Oct;83(4):863-6. doi: 10.1097/00000542-199510000-00028. PMID: 7574068.
22. **Qudsiya Z, De Jesus O.** Subacute Combined Degeneration of the Spinal Cord. 2022 Jun 21. In: *StatPearls* [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan–. PMID: 32644742.
23. **Sun HY, Lee JW, Park KS, Wi JY, Kang HS.** Spine MR imaging features of subacute combined degeneration patients. *Eur Spine J.* 2014 May;23(5):1052-8. doi: 10.1007/s00586-014-3191-4. Epub 2014 Jan 28. PMID: 24469885.
24. **Jain KK, Malhotra HS, Garg RK, Gupta PK, Roy B, Gupta RK.** Prevalence of MR imaging abnormalities in vitamin B12 deficiency patients presenting with clinical features of subacute combined degeneration of the spinal cord. *J Neurol Sci.* 2014 Jul 15;342(1-2):162-6. doi: 10.1016/j.jns.2014.05.020. Epub 2014 May 15. PMID: 24857760.
25. **Xiao CP, Ren CP, Cheng JL, Zhang Y, Li Y, Li BB, Fan YZ.** Conventional MRI for diagnosis of subacute combined degeneration (SCD) of the spinal cord due to vitamin B-12 deficiency. *Asia Pac J Clin Nutr.* 2016;25(1):34-8. doi: 10.6133/apjcn.2016.25.1.04. PMID: 26965759.
26. **Conti N.** Vitamin B12 deficiency presenting with neurologic dysfunction in a 20-year-old male without anemi. *Proc of UCLA Health.* 2021, v. 25. Available at: <https://proceedings.med.ucla.edu/index.php/2022/03/21/vitamin-b12-deficiency-presenting-with-neurologic-dysfunction-in-a-20-year-old-male-without-anemia/>.
27. **Ralapanawa DM, Jayawickreme KP, Ekanayake EM, Jayalath WA.** B12 deficiency with neurological manifestations in the absence of anaemia. *BMC Res Notes.* 2015 Sep 18;8:458. doi: 10.1186/s13104-015-1437-9. PMID: 26385097; PMCID: PMC4575440.
28. **Lindenbaum J, Savage DG, Stabler SP, Allen RH.** Diagnosis of cobalamin deficiency: II. Relative sensitivities of serum cobalamin, methylmalonic acid, and total homocysteine concentrations. *Am J Hematol.* 1990 Jun;34(2):99-107. doi: 10.1002/ajh.2830340205. PMID: 2339684.
29. **Briani C, Dalla Torre C, Citton V, Manara R, Pompanin S, Binotto G, Adami F.** Cobalamin deficiency: clinical picture and radiological findings. *Nutrients.* 2013 Nov 15;5(11):4521-39. doi: 10.3390/nu5114521. PMID: 24248213; PMCID: PMC3847746.
30. **Cao J, Xu S, Liu C.** Is serum vitamin B12 decrease a necessity for the diagnosis of subacute combined degeneration?: A meta-analysis. *Medicine (Baltimore).* 2020 Apr;99(14):e19700. doi: 10.1097/MD.00000000000019700. PMID: 32243408; PMCID: PMC7440176.
31. **Bolaman Z, Kadikoylu G, Yukselen V, Yavasoglu I, Barutca S, Senturk T.** Oral versus intramuscular cobalamin treatment in megaloblastic anemia: a single-center, prospective, randomized, open-label study. *Clin Ther.* 2003 Dec;25(12):3124-34. doi: 10.1016/s0149-2918(03)90096-8. PMID: 14749150.
32. **Castelli MC, Friedman K, Sherry J, Brazzillo K, Genoble L, Bhargava P, Riley MG.** Comparing the efficacy and tolerability of a new daily oral vitamin B12 formulation and intermittent intramuscular vitamin B12 in normalizing low cobalamin levels: a randomized, open-label, parallel-group study. *Clin Ther.* 2011 Mar;33(3):358-371.e2. doi: 10.1016/j.clinthera.2011.03.003. PMID: 21600388.
33. **Kuzminski AM, Del Giacco EJ, Allen RH, Stabler SP, Lindenbaum J.** Effective treatment of cobalamin deficiency with oral cobalamin. *Blood.* 1998 Aug 15;92(4):1191-8. PMID: 9694707.
34. **Kulnigg-Dabsch S.** Autoimmune gastritis. *Wien Med Wochenschr.* 2016 Oct;166(13-14):424-430. doi: 10.1007/s10354-016-0515-5. Epub 2016 Sep 26. PMID: 27671008; PMCID: PMC5065578.