

and short answer items. Descriptive statistics were reported. Qualitative analysis of short answers used a thematic approach.

Results: 205 educators participated. The most common publication was peer-reviewed research manuscript. 31% (61/197) had training in research methodology. Time constraint was the greatest barrier to scholarship (8.61/10). There was a mismatch between actual and ideal hours spent on job related tasks. 69.8% (111/159) of researchers perform research in education. Barriers to research were lack of time, support, expertise, mentorship, funding, reward, challenges of adhering to scientifically rigorous methods, and achieving publication. The most motivating factors to performing research were personal intellectual stimulation and to be a better teacher, 7.57/10 and 6.91/10 respectively. Research study design and scientific writing were the most desired skills to acquire, 61.2%(112/183) and 49.7%(91/183) respectively. Preferred formats for developing research skills were online and a home institution faculty development course; 65/181 and 61/181 favor these respectively. 49.7%(91/183) have a mentor.

Conclusions: Multiple barriers to performing scholarship were identified and impact educators to varying degrees. Potential strategies for improvement were suggested. This information may inform interventions to help support educators in their scholarly pursuits.

7 Barriers to the Remediation of Struggling Learners: A Qualitative Study

Krzyzaniak S, Wolf S, Byyny R, Kaplan B, Barker L, Guerrasio J / OSF Saint Francis Medical Center, Peoria, IL; University of Virginia School of Medicine, Charlottesville, VA; Denver Health Medical Center, Denver, CO; University of Colorado Hospital, Denver, CO

Background: Physician trainees are expected to progress through educational milestones to achieve competence appropriate to their level of training. Meeting this expectation requires identification of the struggling resident through outcomes-based and learner-centered assessments with subsequent action taken to remediate the observed deficiencies.

Objectives: This study was designed to better understand faculty perspectives on remediation, including the barriers to the implementation of a remediation plan for struggling residents.

Methods: The authors conducted structured focus groups of regional stakeholders in medical education to explore barriers to the process of remediating struggling residents. Due to limited existing research and theory around remediation, the authors utilized a constructivist qualitative design with conventional content analysis to evaluate the data. Concepts related to barriers to remediation were identified and overarching themes were developed.

Results: Major themes identified as barriers to remediation were: (1) faculty concern about the premature labeling of

residents as “struggling” and the stigma that comes with remediation; (2) limited availability of resources to devote to remediation; (3) inadequate faculty development and training around the development and implementation of remediation plans; (4) a lack of, or an unwillingness, of resident participation; and (5) a lack of consistent and honest documentation.

Conclusions: The process of remediation is hindered by the emotional response of faculty, a dearth of resources and expertise, learner factors, and a lack of honest and consistent documentation. When implementing a remediation program for a struggling resident, educators should address these elements prior to initiation of the plan.

8 Can Active Learning via the Socratic Method Improve Knowledge Retention Amongst Emergency Medicine Residents?

Wood S, Teran F, Strayer R, Shah K / Mount Sinai, New York, NY

Background: Socratic instruction utilizes targeted questions to expand learners’ understanding of a subject. In medical education, a form of the Socratic method known as “pimping” is used, which at its best may help learners attain greater knowledge in a slightly stressful environment. Although “pimping” is a common teaching strategy for medical students, it has not been studied as an educational tool for Emergency Medicine residents.

Objectives: The focus of this pilot study was to assess Emergency Medicine residents’ perceptions of “pimping” and to evaluate whether teaching via Socratic instruction can enhance knowledge retention amongst residents compared with a standard lecture format.

Methods: This was a prospective, randomized-controlled study performed during emergency department shifts at an urban, academic hospital. Groups of 3-4 residents received a bedside teaching session on head trauma either in a Socratic method i.e. “pimping” style or a lecture style. Groups were taught by one of two emergency physicians trained via an online module and a live session. Residents were asked the same pre-defined set of questions during the session. Afterwards, both groups completed a questionnaire assessing their perceptions of “pimping”. Four weeks after the teaching session, residents completed a follow up quiz.

Results: 72 residents participated in our study. Results of our questionnaire showed that 54% of residents found “pimping” to be an effective teaching method most or all of the time. 82% would use Socratic instruction as a teaching method at least some of the time.

57 out of 72 residents (79%) completed the follow up quiz. Average scores were identical for residents in the “pimping” group (66.2%) compared with the non-pimping group (65.9%). Interns in the “pimping” group had higher