

## Omental Prolapse Through Vaginal Cuff Dehiscence

Jon Solberg, MD\*  
Karan Saravana, BS†

\*University of North Dakota, Department of Emergency Medicine, Bismarck, North Dakota  
†University of North Dakota, School of Medicine and Health Sciences, Bismarck, North Dakota

Section Editor: Manish Amin, DO

Submission History: Submitted February 3, 2022; Revision received February 28, 2022; Accepted May 19, 2022

Electronically published July 18, 2022

Full text available through open access at [http://escholarship.org/uc/uciem\\_cpem](http://escholarship.org/uc/uciem_cpem)

DOI: 10.5811/cpem.2022.2.56353

**Case Presentation:** A 31-year-old female presented to the emergency department with abdominal pain and a 15-centimeter bloody vaginal protrusion, which resulted during an attempted bowel movement. Reduction of the mass was unsuccessful, and the patient was taken to the operating room for examination.

**Discussion:** In patients with a history of vaginal hysterectomy, the vaginal cuff can dehisce and abdominal contents may protrude through the vaginal canal. In this case presentation, the vaginal mass was found to be omental tissue, which could be mistaken for a prolapse of vaginal mucosa. Therefore, a proper pelvic exam is imperative, as prolapse through a cuff dehiscence can lead to severe complications. [Clin Pract Cases Emerg Med. XXXX;X(X)t:X-X.]

**Keywords:** *vaginal cuff dehiscence; omental prolapse; vaginal mass.*

### CASE PRESENTATION

A 31-year-old female with a history of laparoscopic-assisted vaginal hysterectomy two years prior presented by ambulance to the emergency department with an acute onset of abdominal pain and a vaginal protrusion that occurred while straining to pass a bowel movement. Physical examination was notable for a flat but slightly tender abdomen, normal bowel sounds, scant vaginal bleeding, and a 15-centimeter long, blood-tinged mass protruding from the vagina.

A brief and unsuccessful attempt at reduction was made by the emergency physician. Obstetrics and gynecology was consulted, and the patient was taken to the operating room for exam under anesthesia.

### DISCUSSION

The diagnosis was *omental prolapse through vaginal cuff dehiscence*. Following a vaginal hysterectomy, the vaginal cuff is closed surgically.<sup>1</sup> Occasionally, this site can dehisce, allowing abdominal contents such as the small bowel or omentum to enter the vagina or protrude through the vaginal canal.<sup>2,3</sup> In existing literature, reports of prolapses of omentum are uncommon, and photographed cases may illustrate an anterior or apical vaginal bulge, as opposed to a completely



**Image.** External pelvic examination demonstrates a blood-tinged mass (arrow) protruding from the vagina.

visible omental mass, as seen here.<sup>4</sup> Vaginal cuff dehiscence is estimated to have a rate of 0.39%. It is more commonly seen after total laparoscopic hysterectomy (1.35%) compared with

laparoscopic-assisted vaginal hysterectomy (0.28%).<sup>4</sup>

Risk factors include trauma from sexual intercourse, repetitive Valsalva maneuvers, smoking, malnutrition, anemia, diabetes, immunosuppression, and corticosteroid use.<sup>2</sup> Cases typically present as vaginal spotting or post-coital bleeding, and occasionally as pelvic pressure or protrusion.<sup>2</sup> Most cases occur within weeks to months after the procedure, but some can present years later. Patients are at risk for infection due to exposure of peritoneal contents to vaginal and skin flora. Management includes administration of broad-spectrum antibiotics. Partial dehiscence can be managed with rest, but large dehiscence is usually managed surgically.

This case highlights the importance of the pelvic exam in patients with vaginal bleeding and abdominal pain, and care should be taken to not mistake protruding omental tissue for prolapsed vaginal mucosa.

Documented patient informed consent and/or Institutional Review Board approval has been obtained and filed for publication of this case report.

---

*Address for Correspondence:* Karan Saravana, BS, University of North Dakota, School of Medicine and Health Sciences, 1301 N Columbia Rd Stop 9037, Grand Folks, ND 58202. Email: k.saravana@umd.edu.

*Conflicts of Interest:* By the CPC-EM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.

*Copyright:* © 2022 Solberg. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

---

### *CPC-EM Capsule*

What do we already know about this clinical entity?

*Vaginal cuff dehiscence is seen in patients with a history of vaginal hysterectomy. Abdominal contents can prolapse through the vaginal cuff and protrude through the vagina.*

What makes this presentation of disease reportable?

*The image illustrates a rare photograph of omental prolapse via cuff dehiscence.*

What is the major learning point?

*This condition will appear as a bloody mass, which should not be mistaken for prolapsed vaginal mucosa, due to risk of infection.*

How might this improve emergency medicine practice?

*Immediate recognition allows for decreased misidentification and decreased chance of risk, such as infection of peritoneal contents.*

### REFERENCES

1. Binz NM. Complications of gynecologic procedures. In: *Emergency Medicine: A Comprehensive Study Guide*, 9e. McGraw Hill; 2020.
2. McGrane O and Owens S. Elderly female with abdominal pain. *Images Emerg Med*. 2012; 60 (1):132.
3. Sokol ER and Munro EG. Incarcerated vaginal herniation of the omentum mimicking vaginal prolapse. *Am J Obstet Gynecol*. 2007 Mar;196(3):e7-8.
4. Clarke-Pearson DL, Geller EJ. Complications of hysterectomy. *Obstet Gynecol*. 2013 Mar;121(3):654-73.