



RV LV
TV MV
A

WestJEM

Volume 26, Issue 1.2, January 2025

Open Access at www.westjem.com

ISSN 1936-900X

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health Indexed in MEDLINE



Special Issue on Educational Research and Practice



CALIFORNIA ACEP
AMERICAN COLLEGE OF EMERGENCY PHYSICIANS



UC Irvine Health



CALAAEM
CALIFORNIA CHAPTER DIVISION OF THE
AMERICAN ACADEMY OF EMERGENCY MEDICINE

A Peer-Reviewed, International Professional Journal

SAVE THE DATE!



SPRING SEMINAR

**Renaissance
Glendale
Hotel & Spa**
Phoenix, AZ

Sunday
April 27, 2025
Pre-Conference
Workshops

Sunday-Monday
April 27-28, 2025
Resident Student
Organization Session

Monday-Thursday
April 28-May 1, 2025
Spring
Seminar

**Registration is
Coming Soon –
Watch your inbox for the
official announcement!**





PennState Health

Penn State Health Emergency Medicine



About Us: Penn State Health is a multi-hospital health system serving patients and communities across central Pennsylvania. We are the only medical facility in Pennsylvania to be accredited as a Level I pediatric trauma center and Level I adult trauma center. The system includes Penn State Health Milton S. Hershey Medical Center, Penn State Health Children’s Hospital and Penn State Cancer Institute based in Hershey, Pa.; Penn State Health Hampden Medical Center in Enola, Pa.; Penn State Health Holy Spirit Medical Center in Camp Hill, Pa.; Penn State Health Lancaster Medical Center in Lancaster, Pa.; Penn State Health St. Joseph Medical Center in Reading, Pa.; Pennsylvania Psychiatric Institute, a specialty provider of inpatient and outpatient behavioral health services, in Harrisburg, Pa.; and 2,450+ physicians and direct care providers at 225 outpatient practices. Additionally, the system jointly operates various healthcare providers, including Penn State Health Rehabilitation Hospital, Hershey Outpatient Surgery Center and Hershey Endoscopy Center.



We foster a collaborative environment rich with diversity, share a passion for patient care, and have a space for those who share our spark of innovative research interests. Our health system is expanding and we have opportunities in both academic hospital as well community hospital settings.

Benefit highlights include:

- Competitive salary with sign-on bonus
- Comprehensive benefits and retirement package
- Relocation assistance & CME allowance
- Attractive neighborhoods in scenic central Pennsylvania

FOR MORE INFORMATION PLEASE CONTACT:

Heather Peffley, PHR CPRP

Penn State Health Lead Physician Recruiter

hpeffley@pennstatehealth.psu.edu



Penn State Health is fundamentally committed to the diversity of our faculty and staff. We believe diversity is unapologetically expressing itself through every person’s perspectives and lived experiences. We are an equal opportunity and affirmative action employer. All qualified applicants will receive consideration for employment without regard to age, color, disability, gender identity or expression, marital status, national or ethnic origin, political affiliation, race, religion, sex (including pregnancy), sexual orientation, veteran status, and family medical or genetic information.



SAVE THE DATE

SAEM25

PHILADELPHIA, PA • MAY 13-16, 2025

SAEM25 CDEM ACADEMY MEETING



Clerkship Directors in Emergency Medicine

Clerkship Directors in Emergency Medicine (CDEM) Meeting

Philadelphia Marriott Downtown | Wednesday, May 14, 2025 | 3:30 PM - 5:30 PM

www.saem.org/cdem

ADDITIONAL RESOURCES OFFERED BY SAEM/CDEM

• **SAEMTests**

• **CDEM Curriculum**

Seattle

Register Here:



CORD

EM

ACADEMIC ASSEMBLY

March 2-5, 2025

ARE YOU REGISTERED? EARLY BIRD PRICING ENDS **01/18/25**

Western Journal of Emergency Medicine: CDEM/CORD 2025 Special Education Issue

A Note from the Editors:

We are excited to publish the 10th issue of the Western Journal of Emergency Medicine (WestJEM) Education Issue and first year of a rolling decision process. Over 10 years ago a unique relationship was formed between WestJEM, the Council of Residency Director for Emergency Medicine and the Clerkship Directors of Emergency Medicine. The idea was to promote and disseminate educational scholarship which has been accomplished over the past decade. Senior and junior researchers have an opportunity to publish in the education issue because of the diverse nature of our submission categories, ranging from original research to brief educational advances. A successful issue requires the courage of the authors to submit their work for peer review and we do our best to provide detailed feedback regardless of the final decision. Publication of the issue requires the commitment and hard work of the publication staff, leadership of the organizations, editors, and peer reviewers. We want to thank them all for their efforts and professionalism. The topics of this year's education issue likely reflect the focus of educators as we entered a post-covid reality. Many of the topics were related to innovative curriculums and focused on the benefits derived. There were also several articles that were dedicated to the administrative aspects of residency and fellowships and how that has changed after COVID. We have already started to receive and review submissions for next year's education issue. The editorial staff review every submission on a rolling basis and, once accepted, the articles are available on PubMed. There are also no processing fees when accepted to the Education Issue. This is a great opportunity to submit your educational scholarship, thereby enhancing your professional development and disseminating your work to others. We are excited that this experiment has flourished, and we look forward to seeing your work in our 11th issue.

*Jeffrey Love, MD
Georgetown University School of Medicine
Co-Editor of Annual Special Issue on Education Research and Practice*

*Douglas Ander, MD
Emory University
Co-Editor of Annual Special Issue on Education Research and Practice*

*The Western Journal of Emergency Medicine: Integrating
Emergency Care with Population Health would like to thank The
Clerkship Directors in Emergency Medicine (CDEM) and the
Council of Residency Directors in Emergency Medicine (CORD)
for helping to make this collaborative special issue possible.*

Western Journal of Emergency Medicine:

Integrating Emergency Care with Population Health

Indexed in MEDLINE, PubMed, and Clarivate Web of Science, Science Citation Index Expanded

JOURNAL FOCUS

Emergency medicine is a specialty which closely reflects societal challenges and consequences of public policy decisions. The emergency department specifically deals with social injustice, health and economic disparities, violence, substance abuse, and disaster preparedness and response. This journal focuses on how emergency care affects the health of the community and population, and conversely, how these societal challenges affect the composition of the patient population who seek care in the emergency department. The development of better systems to provide emergency care, including technology solutions, is critical to enhancing population health.

Table of Contents

- 1** **Program Signaling in Emergency Medicine: The 2022–2023 Program Director Experience**
AE Pelletier-Bui, T Fallon, L Smith, T Strout, M Fischer, M Olaf, E McDonough, B Barbas, M Cirone, E Barrall Werley
- 11** **Emergency Medicine Milestones Final Ratings Are Often Subpar**
DL Gorgas, KB Joldersma, FK Ankel, WA Carter, MA Barton, EJ Reisdorff
- 15** **Emergency Department Slit Lamp Interdisciplinary Training Via Longitudinal Assessment in Medical Practice**
S Hamou, S Ghiaee, C Chung, M Lloyd, K Khem, XC Zhang
- 25** **Teaching the New Ways: Improving Resident Documentation for the New 2023 Coding Requirements**
N Zapolsky, A Cardell, R Desai, S Frisch, N Jobeun, D Novak, M Silver, AS Chung
- 29** **Telesimulation Use in Emergency Medicine Residency Programs: National Survey of Residency Simulation Leaders**
M Berger, J Buckanavage, J Jordan, S Lai, L Regan
- 35** **Substantial Variation Exists in Clinical Exposure to Chief Complaints Among Residents Within an Emergency Medicine Training Program**
CM Jewell, AT Hummel, DJ Hekman, BH Schnapp
- 41** **The Effect of Hospital Boarding on Emergency Medicine Residency Productivity**
P Moffett, A Best, N Lewis, S Miller, G Hickam, H Kissel-Smith, L Barrera, S Huang, J Moll
- 50** **Making A Difference: Launching a Multimodal, Resident-Run Social Emergency Medicine Program**
NP Newton, C Freeman, P Panakos
- 59** **Palliative Care Boot Camp Offers Skill Building for Emergency Medicine Residents**
J Cooper, J Fredette
- 63** **Effectiveness of a Collaborative, Virtual Outreach Curriculum for 4th-Year EM-bound Students at a Medical School Affiliated with a Historically Black College and University**
C Brown, R Carter, N Hartman, A Hammond, E MacNeill, L Holden, A Pierce, L Campbell, M Norman

Policies for peer review, author instructions, conflicts of interest and human and animal subjects protections can be found online at www.westjem.com.

2024 Gold Standard Reviewers

The *WestJEM* Special Issue in Educational Research & Practice couldn't exist without our many reviewers. To all, we wish to express our sincerest appreciation for their contributions to this year's success. Each year a number of reviewers stand out for their (1) detailed reviews, (2) grasp of the tenets of education scholarship and (3) efforts to provide feedback that mentors authors on how to improve.

This year's "Gold Standard" includes:

- Dave Carlberg
- Michael Cassara
- Max Griffith/Bjorn Watsjold*
- Alana Harp/Deena Bengiamin*
- Mojibade Hassan/Kathleen White/Brad Barth*
- Heather Hechter
- Adam Heilmann/Erin Kane/John Schneider/Albert Kim*
- Corlin Jewel
- Julia Isaacson/Kerry McCabe*
- Kaitlin Lipner/Chris Merritt*
- Evie Marcolini
- Dan Mayer
- Colin McMahon/Anne Messman*
- Elspeth Pearce
- Adam Rieves
- Eric Pellegrini /Emily Rose/Aarti Jain/Taku Taira/Jeff Riddell*
- John Priester/Richard Bounds*
- Thaddeus Schmitt
- Jeffrey Siegelman
- Jacqueline Sippel/Ashley Foreman/Erica Shaver/Chris Kiefer*
- Olivia Urbanoor/Jessica Baez/Sally Santen*

*Mentored Peer Reviews from Emergency Medicine Education Fellowship Programs

CDEM/CORD Guest Consulting Editors

We would also like to recognize our guest consulting editors who assisted with pre-screening submissions during our initial peer-review stages.

Thank you for all of your efforts and contributions.

CDEM

- Christine Stehman
- Eric Shappell
- Sharon Bord
- Andrew Golden

CORD

- Jenna Fredette
- Danielle Hart
- William Soares III
- Jamie Jordan
- Anne Messman
- Logan Weygandt

Consulting Statistician/ Psychometrician

- David Way
-



Clerkship Directors in Emergency Medicine



COUNCIL OF RESIDENCY DIRECTORS IN EMERGENCY MEDICINE

Western Journal of Emergency Medicine:

Integrating Emergency Care with Population Health

Indexed in MEDLINE, PubMed, and Clarivate Web of Science, Science Citation Index Expanded

Jeffrey N. Love, MD, Guest Editor

Georgetown School of Medicine- Washington, District of Columbia

Chris Merritt, MD, MPH, Associate Guest Editor

Alpert Medical School of Brown University-Providence, Rhode Island

Benjamin Schnapp, MD, MEd, Associate Guest Editor

University of Wisconsin-Madison, Wisconsin

Wendy Macias-Konstantopoulos, MD, MPH, Associate Editor

Massachusetts General Hospital- Boston, Massachusetts

Danya Khoujah, MBBS, Associate Editor

University of Maryland School of Medicine- Baltimore, Maryland

Patrick Joseph Maher, MD, MS, Associate Editor

Ichan School of Medicine at Mount Sinai- New York, New York

Yanina Purim-Shem-Tov, MD, MS, Associate Editor

Rush University Medical Center-Chicago, Illinois

Gayle Galletta, MD, Associate Editor

University of Massachusetts Medical School- Worcester, Massachusetts

Dan Mayer, MD, Associate Editor

Retired from Albany Medical College- Niskayuna, New York

Julianna Jung, MD, Associate Guest Editor

Johns Hopkins Hospital, Baltimore, Maryland

Douglas Franzen, MD, Associate Guest Editor

Harborview Medical Center, Seattle, Washington

Henry Wilkerson, MD, Associate Editor

University of Maryland

Michael Gottlieb, MD, Associate Editor

Rush Medical Center-Chicago, Illinois

Sara Krzyzaniak, MD Associate Guest Editor

Stanford University-Palo Alto, California

Susan R. Wilcox, MD, Associate Editor

Massachusetts General Hospital- Boston, Massachusetts

Donna Mendez, MD, EdD, Associate Editor

University of Texas-Houston/McGovern Medical School- Houston, Texas

Taku Taira, MD, EDD, Associate Guest Editor

LAC + USC Medical Center-Los Angeles, California

Douglas S. Ander, MD, Guest Editor

Emory University School of Medicine-Atlanta, Georgia

Edward Ullman, MD, Associate Guest Editor

Harvard University-Cambridge, Massachusetts

Abra Fant MD, MS, Associate Guest Editor

Northwestern Medicine-Chicago, Illinois

Kendra Parekh, MD, MS, Associate Guest Editor

Vanderbilt University-Nashville, Tennessee

Matthew Tews, DO, MS, Associate Guest Editor

Indiana University School of Medicine, Augusta, Georgia

Rick A. McPheeters, DO, Associate Editor

Kern Medical- Bakersfield, California

Niels K. Rathlev MD, MS, Associate Editor

Tufts University School of Medicine-Boston, Massachusetts

Shahram Lotfipour, MD, MPH, Managing Associate Editor

University of California, Irvine School of Medicine- Irvine, California

Mark I. Langdorf, MD, MHPE, Editor-in-Chief

University of California, Irvine School of Medicine- Irvine, California

Resident Editors

AAEM/RSA

Tehreem Rehman, MD
Advocate Christ Medical Center

ACOEP

Justina Truong, DO
Kingman Regional Medical Center

Section Editors

Behavioral Emergencies

Erin Dehon, PhD
University of Mississippi Medical Center

Leslie Zun, MD, MBA

Chicago Medical School

Marc L. Martel, MD

Hennepin County Medical Center

Cardiac Care

Fred A. Severyn, MD
University of Colorado School of Medicine

Michael C. Kurz, MD
University of Alabama at Birmingham

Sam S. Torbati, MD
Cedars-Sinai Medical Center

Clinical Practice

Cortlyn W. Brown, MD
Carolinas Medical Center

Casey Clements, MD, PhD
Mayo Clinic

Patrick Meloy, MD
Emory University

Nicholas Pettit, DO, PhD
Indiana University

Eric Snoey, MD
Alameda County Medical Center

David Thompson, MD
University of California, San Francisco

Kenneth S. Whitlow, DO
Kaweah Delta Medical Center

Critical Care

Christopher "Kit" Tainter, MD
University of California, San Diego

Gabriel Wardi, MD
University of California, San Diego

Joseph Shiber, MD
University of Florida-College of Medicine

Matt Prekker MD, MPH
Hennepin County Medical Center

David Page, MD
University of Alabama

Erik Melnychuk, MD
Geisinger Health

Quincy Tran, MD, PhD
University of Maryland

Disaster Medicine

Christopher Kang, MD
Madigan Army Medical Center

Education

Danya Khoujah, MBBS
University of Maryland School of Medicine

Jeffrey Druck, MD
University of Colorado

John Burkhardt, MD, MA
University of Michigan Medical School

Michael Epter, DO
Maricopa Medical Center

ED Administration, Quality, Safety

David C. Lee, MD
Northshore University Hospital

Gary Johnson, MD
Upstate Medical University

Brian J. Yun, MD, MBA, MPH
Harvard Medical School

Laura Walker, MD
Mayo Clinic

León D. Sánchez, MD, MPH
Beth Israel Deaconess Medical Center

William Fernandez, MD, MPH
University of Texas Health-San Antonio

Emergency Medical Services

Daniel Joseph, MD
Yale University

Joshua B. Gaither, MD
University of Arizona, Tucson

Julian Mapp
University of Texas, San Antonio

Shira A. Schlesinger, MD, MPH
Harbor-UCLA Medical Center

Geriatrics

Cameron Gettel, MD
Yale School of Medicine

Stephen Meldon, MD
Cleveland Clinic

Luna Ragsdale, MD, MPH
Duke University

Health Equity

Emily C. Manchanda, MD, MPH
Boston University School of Medicine

Mandy J. Hill, DrPH, MPH
UT Health McGovern Medical School

Infectious Disease

Elissa Schechter-Perkins, MD, MPH
Boston University School of Medicine

Ioannis Koutroulis, MD, MBA, PhD
Drexel University College of Medicine

Kevin Lunney, MD, MHS, PhD
University of Maryland School of Medicine

Robert Derlet, MD
Founding Editor, California Journal of Emergency
Medicine
University of California, Davis

Stephen Liang, MD, MPH
Washington University School of Medicine

Injury Prevention

Mark Faul, PhD, MA
Centers for Disease Control and Prevention

Wirachin Hoonpongsimanont, MD, MSBATS
Eisenhower Medical Center

International Medicine

Heather A.. Brown, MD, MPH
Prisma Health Richland

Taylor Burkholder, MD, MPH
Keck School of Medicine of USC

Christopher Greene, MD, MPH
University of Alabama

Chris Mills, MD, MPH
Santa Clara Valley Medical Center

Shada Rouhani, MD
Brigham and Women's Hospital

Legal Medicine

Melanie S. Heniff, MD, JD
Indiana University School of Medicine

Greg P. Moore, MD, JD
Madigan Army Medical Center

Statistics and Methodology

Shu B. Chan MD, MS
Resurrection Medical Center

Stormy M. Morales Monks, PhD, MPH
Texas Tech Health Science University

Soheil Saadat, MD, MPH, PhD
University of California, Irvine

James A. Meltzer, MD, MS
Albert Einstein College of Medicine

Musculoskeletal

Juan F. Acosta DO, MS
Pacific Northwest University

Neurosciences

Antonio Siniscalchi, MD
Annunziata Hospital, Cosenza, Italy

Rick Lucarelli, MD
Medical City Dallas Hospital

William D. Whetstone, MD
University of California, San Francisco

Pediatric Emergency Medicine

Paul Walsh, MD, MSc
University of California, Davis

Muhammad Waseem, MD
Lincoln Medical & Mental Health Center

Deena Berkowitz, MD, MPH
Children's National Hospital

Cristina M. Zeretzke-Bien, MD
University of Florida

Public Health

Jeremy Hess, MD, MPH
University of Washington Medical Center

Jacob Manteuffel, MD
Henry Ford Hospital

John Ashurst, DO
Lehigh Valley Health Network

Tony Zitek, MD
Kendall Regional Medical Center

Trevor Mills, MD, MPH
Northern California VA Health Care

Erik S. Anderson, MD
Alameda Health System-Highland Hospital

Technology in Emergency Medicine

Nikhil Goyal, MD
Henry Ford Hospital

Phillips Perera, MD
Stanford University Medical Center

Trauma

Pierre Borczuk, MD
Massachusetts General Hospital/Harvard
Medical School

Toxicology

Brandon Wills, DO, MS
Virginia Commonwealth University

Jeffrey R. Suchard, MD
University of California, Irvine

Ultrasound

J. Matthew Fields, MD
Thomas Jefferson University

Shane Summers, MD
Brooke Army Medical Center

Robert R. Ehrman
Wayne State University

Ryan C. Gibbons, MD
Temple Health

Official Journal of the California Chapter of the American College of Emergency Physicians, the American College of Osteopathic Emergency Physicians, and the California Chapter of the American Academy of Emergency Medicine



Available in MEDLINE, PubMed, PubMed Central, CINAHL, SCOPUS, Google Scholar, eScholarship, Melvyl, DOAJ, EBSCO, EMBASE, Medscape, HINARI, and MDLinx Emergency Med. Members of OASPA. Editorial and Publishing Office: WestJEM/Department of Emergency Medicine, UC Irvine Health, 3800 W Chapman Ave Ste 3200, Orange, CA 92868, USA. Office: 1-714-456-6389; Email: Editor@westjem.org.

Western Journal of Emergency Medicine:

Integrating Emergency Care with Population Health

Indexed in MEDLINE, PubMed, and Clarivate Web of Science, Science Citation Index Expanded

Editorial Board

Amin A. Kazzi, MD, MAAEM
*The American University of Beirut,
Beirut, Lebanon*

Brent King, MD, MMM
University of Texas, Houston

Christopher E. San Miguel, MD
*Ohio State University Wexner Medical
Center*

Daniel J. Dire, MD
*University of Texas Health Sciences
Center San Antonio*

Douglas Ander, MD
Emory University

Edward Michelson, MD
Texas Tech University

Edward Panacek, MD, MPH
University of South Alabama

Francesco Della Corte, MD
*Azienda Ospedaliera Universitaria
"Maggiore della Carità," Novara, Italy*

Jaqueline Le, MD
Desert Regional Medical Center

Jeffrey Love, MD
Georgetown School of Medicine

Katsuhiro Kanemaru, MD
*University of Miyazaki Hospital, Miyazaki,
Japan*

Leslie Zun, MD, MBA
Chicago Medical School

Linda S. Murphy, MLIS
*University of California, Irvine School of
Medicine Librarian*

Pablo Aguilera Fuenzalida, MD
*Pontificia Universidad Catolica de Chile,
Región Metropolitana, Chile*

Peter A. Bell, DO, MBA
Baptist Health Sciences University

Peter Sokolove, MD
University of California, San Francisco

Rachel A. Lindor, MD, JD
Mayo Clinic

Robert Suter, DO, MHA
UT Southwestern Medical Center

Robert W. Derlet, MD
University of California, Davis

Scott Rudkin, MD, MBA
University of California, Irvine

Scott Zeller, MD
University of California, Riverside

Steven H. Lim Hoon Chin, MD
*Changi General Hospital, Simei,
Singapore*

Wirachin Hoonpongmanont, MD,
MSBATS
*Siriraj Hospital, Mahidol University,
Bangkok, Thailand*

Advisory Board

Elena Lopez-Gusman, JD
*California ACEP
American College of Emergency
Physicians*

Jennifer Kanapicki Comer, MD
FAAEM
*California Chapter Division of AAEM
Stanford University School of Medicine*

DeAnna McNett
*American College of Osteopathic
Emergency Physicians*

Kimberly Ang, MBA
UC Irvine Health School of Medicine

Randall J. Young, MD, MMM, FACEP
*California ACEP
American College of Emergency
Physicians
Kaiser Permanente*

Mark I. Langdorf, MD, MHPE,
MAAEM, FACEP
UC Irvine Health School of Medicine

Robert Suter, DO, MHA
*American College of Osteopathic
Emergency Physicians
UT Southwestern Medical Center*

Shahram Lotfipour, MD, MPH
FAAEM, FACEP
UC Irvine Health School of Medicine

Jorge Fernandez, MD, FACEP
UC San Diego Health School of Medicine

Editorial Staff

Isabelle Nepomuceno, BS
Executive Editorial Director

Ian Ollife, BS
WestJEM Editorial Director

Emily Kane, BS
WestJEM Editorial Director

Tran Nguyen, BS
CPC-EM Editorial Director

Stephanie Burmeister, MLIS
WestJEM Staff Liaison

Cassandra Saucedo, MS
Executive Publishing Director

Nicole Valenzi, BA
WestJEM Publishing Director

Alyson Tsai, BS
CPC-EM Publishing Director

June Casey, BA
Copy Editor

Official Journal of the California Chapter of the American College of Emergency Physicians, the American College of Osteopathic Emergency Physicians, and the California Chapter of the American Academy of Emergency Medicine



Available in MEDLINE, PubMed, PubMed Central, Europe PubMed Central, PubMed Central Canada, CINAHL, SCOPUS, Google Scholar, eScholarship, Melvyl, DOAJ, EBSCO, EMBASE, Medscape, HINARI, and MDLinx Emergency Med. Members of OASPA.

Editorial and Publishing Office: *WestJEM*/Department of Emergency Medicine, UC Irvine Health, 333 City Blvd, West, Rt 128-01, Orange, CA 92866, USA

Office: 1-714-456-6389; Email: Editor@westjem.org

Western Journal of Emergency Medicine:

Integrating Emergency Care with Population Health

Indexed in MEDLINE, PubMed, and Clarivate Web of Science, Science Citation Index Expanded

This open access publication would not be possible without the generous and continual financial support of our society sponsors, department and chapter subscribers.

Professional Society Sponsors

AMERICAN COLLEGE OF OSTEOPATHIC EMERGENCY PHYSICIANS
CALIFORNIA ACEP

CALIFORNIA CHAPTER DIVISION OF
AMERICAN ACADEMY OF EMERGENCY MEDICINE

Academic Department of Emergency Medicine Subscriber

Alameda Health System-Highland Hospital Oakland, CA	Emory University Atlanta, GA	Loma Linda University Medical Center Loma Linda, CA	Ohio State University Wexner Medical Center Columbus, OH
Ascension Resurrection Chicago, IL	Franciscan Health Carmel, IN	Louisiana State University Shreveport Shreveport, LA	Oregon Health and Science University Portland, OR
Arnot Ogden Medical Center Elmira, NY	Geisinger Medical Center Danville, PA	Massachusetts General Hospital/ Brigham and Women's Hospital/ Harvard Medical Boston, MA	Penn State Milton S. Hershey Medical Center Hershey, PA
Atrium Health Wake Forest Baptist Winston-Salem, NC	Healthpartners Institute/ Regions Hospital Minneapolis, MN	Mayo Clinic in Florida Jacksonville, FL	Poliklinika Drinkovic Zagreb, Croatia
Baylor College of Medicine Houston, TX	Hennepin Healthcare Minneapolis, MN	Mayo Clinic College of Medicine in Rochester Rochester, MN	Prisma Health/ University of South Carolina SOM Greenville Greenville, SC
Baystate Medical Center Springfield, MA	Henry Ford Hospital Detroit, MI	Mayo Clinic in Arizona Phoenix, AZ	Rush University Medical Center Chicago, IL
Beth Israel Deaconess Medical Center Boston, MA	Henry Ford Wyandotte Hospital Wyandotte, MI	Medical College of Wisconsin Affiliated Hospital Milwaukee, WI	Rutgers Robert Wood Johnson Medical School New Brunswick, NJ
Brigham and Women's Hospital Boston, MA	Howard County Department of Fire and Rescue Marriottsville, MD	Mount Sinai Medical Center Miami Beach Miami Beach, FL	St. Luke's University Health Network Bethlehem, PA
Brown University-Rhode Island Hospital Providence, RI	Icahn School of Medicine at Mt Sinai New York, NY	Mount Sinai Morningside New York, NY	Southern Illinois University School of Medicine Springfield, IL
Carolinas Medical Center Charlotte, NC	Indiana University School of Medicine Indianapolis, IN	New York University Langone Health New York, NY	Stony Brook University Hospital Stony Brook, NY
Cedars-Sinai Medical Center Los Angeles, CA	INTEGRIS Health Oklahoma City, OK	North Shore University Hospital Manhasset, NY	SUNY Upstate Medical University Syracuse, NY
Cleveland Clinic Cleveland, OH	Kaweah Delta Health Care District Visalia, CA	NYC Health and Hospitals/ Jacobi New York, NY	Temple University Philadelphia, PA
Desert Regional Medical Center Palm Springs, CA	Kent Hospital Warwick, RI	Ochsner Medical Center New Orleans, LA	Texas Tech University Health Sciences Center El Paso, TX
Eisenhower Health Rancho Mirage, CA	Kern Medical Bakersfield, CA		

State Chapter Subscriber

ARIZONA CHAPTER DIVISION OF THE
AMERICAN ACADEMY OF EMERGENCY MEDICINE
CALIFORNIA CHAPTER DIVISION OF THE
AMERICAN ACADEMY OF EMERGENCY MEDICINE
FLORIDA CHAPTER DIVISION OF THE
AMERICAN ACADEMY OF EMERGENCY MEDICINE

GREAT LAKES CHAPTER DIVISION OF THE
AMERICAN ACADEMY OF EMERGENCY MEDICINE
TENNESSEE CHAPTER DIVISION OF THE
AMERICAN ACADEMY OF EMERGENCY MEDICINE

UNIFORMED SERVICES CHAPTER DIVISION OF THE
AMERICAN ACADEMY OF EMERGENCY MEDICINE
VIRGINIA CHAPTER DIVISION OF THE
AMERICAN ACADEMY OF EMERGENCY MEDICINE

International Society Partners

EMERGENCY MEDICINE ASSOCIATION OF TURKEY
LEBANESE ACADEMY OF EMERGENCY MEDICINE
MEDITERRANEAN ACADEMY OF EMERGENCY MEDICINE

NORWEGIAN SOCIETY FOR EMERGENCY MEDICINE
SOCIEDAD ARGENTINA DE EMERGENCIAS

SOCIEDAD CHILENO MEDICINA URGENCIA
THAI ASSOCIATION FOR EMERGENCY MEDICINE

To become a *WestJEM* departmental sponsor, waive article processing fee, receive print and copies for all faculty and electronic for faculty/residents, and free CME and faculty/fellow position advertisement space, please go to <http://westjem.com/subscribe> or contact:

Stephanie Burmeister
WestJEM Staff Liaison
Phone: 1-800-884-2236
Email: sales@westjem.org

Western Journal of Emergency Medicine:

Integrating Emergency Care with Population Health

Indexed in MEDLINE, PubMed, and Clarivate Web of Science, Science Citation Index Expanded

This open access publication would not be possible without the generous and continual financial support of our society sponsors, department and chapter subscribers.

Professional Society Sponsors

AMERICAN COLLEGE OF OSTEOPATHIC EMERGENCY PHYSICIANS
CALIFORNIA ACEP

CALIFORNIA CHAPTER DIVISION OF
AMERICAN ACADEMY OF EMERGENCY MEDICINE

Academic Department of Emergency Medicine Subscriber

The University of Texas Medical
Branch
Galveston, TX

University of Chicago
Chicago, IL

University of Southern California
Los Angeles, CA
University of Vermont Medical Center
Burlington, VA

UT Health Houston McGovern
Medical School
Houston, TX

University of Cincinnati Medical Center/
College of Medicine
Cincinnati, OH

University of Virginia Health
Charlottesville, VA

Touro University College of
Osteopathic Medicine
Vallejo, CA

University of Colorado Denver
Denver, CO

University of Washington - Harborview
Medical Center
Seattle, WA

Trinity Health Muskegon Hospital
Muskegon, MI
UMass Memorial Health
Worcester, MA

University of Florida, Jacksonville
Jacksonville, FL

University of Wisconsin Hospitals
and Clinics
Madison, WI

University at Buffalo Program
Buffalo, NY

University of Iowa Hospitals and Clinics
Iowa City, IA

UT Southwestern Medical Center
Dallas, TX

University of Alabama, Birmingham
Birmingham, AL

University of Kansas Health System
Kansas City, IA

Franciscan Health Olympia Fields
Phoenix, AZ

University of Arizona College of
Medicine-Tucson
Little Rock, AR

University of Louisville
Louisville, KY

WellSpan York Hospital
York, PA

University of Arkansas for Medical
Sciences
Galveston, TX

University of Maryland School of
Medicine
Baltimore, MD

West Virginia University
Morgantown, WV

University of California, Davis
Medical Center
Sacramento, CA

University of Miami Jackson Health
System
Miami, FL

Wright State University Boonshoft
School of Medicine
Fairborn, OH

University of California San Francisco
General Hospital
San Francisco, CA

University of Michigan
Ann Arbor, MI

Yale School of Medicine
New Haven, CT

University of California San Francisco
Fresno
Fresno, CA

University of North Dakota School of
Medicine and Health Sciences
Grand Forks, ND

University of Southern Alabama
Mobile, AL

State Chapter Subscriber

ARIZONA CHAPTER DIVISION OF THE
AMERICAN ACADEMY OF EMERGENCY MEDICINE
CALIFORNIA CHAPTER DIVISION OF THE
AMERICAN ACADEMY OF EMERGENCY MEDICINE
FLORIDA CHAPTER DIVISION OF THE
AMERICAN ACADEMY OF EMERGENCY MEDICINE

GREAT LAKES CHAPTER DIVISION OF THE
AMERICAN ACADEMY OF EMERGENCY
MEDICINE

TENNESSEE CHAPTER DIVISION OF THE

AMERICAN ACADEMY OF EMERGENCY MEDICINE
UNIFORMED SERVICES CHAPTER DIVISION OF THE
AMERICAN ACADEMY OF EMERGENCY MEDICINE

VIRGINIA CHAPTER DIVISION OF THE
AMERICAN ACADEMY OF EMERGENCY MEDICINE

International Society Partners

EMERGENCY MEDICINE ASSOCIATION OF TURKEY
LEBANESE ACADEMY OF EMERGENCY MEDICINE
MEDITERRANEAN ACADEMY OF EMERGENCY MEDICINE

NORWEGIAN SOCIETY FOR EMERGENCY MEDICINE
SOCIEDAD ARGENTINA DE EMERGENCIAS

SOCIEDAD CHILENO MEDICINA URGENCIA
THAI ASSOCIATION FOR EMERGENCY
MEDICINE

To become a *WestJEM* departmental sponsor, waive article processing fee, receive print and copies for all faculty and electronic for faculty/residents, and free CME and faculty/fellow position advertisement space, please go to <http://westjem.com/subscribe> or contact:

Stephanie Burmeister
WestJEM Staff Liaison
Phone: 1-800-884-2236
Email: sales@westjem.org

Program Signaling in Emergency Medicine: The 2022–2023 Program Director Experience

Alexis E. Pelletier-Bui, MD*
 Timothy Fallon, MD[†]
 Liza Smith, MD[‡]
 Tania Strout, PhD, MS[†]
 Michelle Fischer, MD, MPH[§]
 Mark Olaf, DO^{||}
 Erin McDonough, MD[¶]
 Brian Barbas, MD[#]
 Michael Cirone, MD^{**}
 Elizabeth Barrall Werley, MD[§]

*Cooper University Hospital/Cooper Medical School of Rowan University, Department of Emergency Medicine, Camden, New Jersey
[†]Maine Medical Center, Tufts University School of Medicine, Department of Emergency Medicine, Portland, Maine
[‡]University of Massachusetts Chan School of Medicine, Baystate Medical Center, Department of Emergency Medicine, Springfield, Massachusetts
[§]Penn State College of Medicine/Penn State Health Milton S. Hershey Medical Center, Department of Emergency Medicine, Hershey, Pennsylvania
^{||}Geisinger Commonwealth School of Medicine, Department of Emergency Medicine, Scranton, Pennsylvania
[¶]University of Cincinnati College of Medicine, Department of Emergency Medicine, Cincinnati, Ohio
[#]Loyola University Chicago, Stritch School of Medicine, Loyola University Medical Center, Department of Emergency Medicine, Maywood, Illinois
^{**}University of Illinois, Department of Emergency Medicine, Chicago, Illinois

Section Editors: Paul Logan Weygandt, MD and Chris Merritt, MD

Submission history: Submitted February 2, 2024; Revision received April 22, 2024; Accepted June 3, 2024

Electronically published August 27, 2024

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.19392

Introduction: Program signaling (PS), which enables residency applicants to signal their preference for a specific program, was introduced in emergency medicine (EM) in the 2022–2023 residency application cycle. In this study we evaluated EM program directors' (PD) utilization of PS in application review and ranking. This study also explores the relationship between program characteristics and number of signals received as well as the relative importance and utilization of signals related to the number of signals received.

Methods: This is an institutional review board-approved, cross-sectional study of PDs at Accreditation Council for Graduate Medical Education-accredited EM residency programs. We used descriptive statistics to describe the characteristics of residency programs and practices around PS. Measures of central tendency and dispersion summarized continuous variables. We used chi-square analysis or the Fisher exact test for comparisons between groups for categorical variables. Comparisons for continuous variables were made using the *t*-test for independent samples or analysis of variance.

Results: The response rate was 41% ($n = 113/277$ EM programs). Most programs participated in PS ($n = 261/277$ EM programs, 94.2%). Mean number of signals received was 60 (range 2–203). Signals received varied based on program characteristics including geographic location and program type, duration, environment, and longevity. Most used PS in holistic review (52.2%), but other uses varied by proportion of applications that were signaled. The importance of PS in application review (mean 2.9; 1–5 scale, 1 = not important, 5 = extremely important) and rank list preparation (2.1) was relatively low compared to other application elements such as standardized letters of evaluation (4.97 for review, 4.90 for ranking).

Conclusion: The study provides insights into PS utilization in EM's inaugural year. We have identified patterns of signal use based on program characteristics and number of signals received that can inform signal allocation and utilization on an individual applicant and program level. A more nuanced understanding of signal use can provide valuable insight as the specialty of EM grapples with fluctuations in its applicant numbers and shifting demographics of its applicant pool. [West J Emerg Med. 2025;26(1.2)1–10.]

INTRODUCTION

Program signaling (PS) was introduced into the residency application process in response to the increasing number of applications received by programs, exacerbating the challenge of comprehensive holistic review.¹ Subsequently, EM has experienced drastic fluctuations in the number of applicants pursuing EM and specialty Match rates, as well as unprecedented changes to the demographics of its application pool over the last several years.² Even with variability in the number of applications to emergency medicine (EM) in recent years, EM application numbers remain significantly above what they were 10 years ago.^{2,3} Program signaling allows applicants to assign signals to their most desired training programs, so that programs may focus their holistic efforts toward high-yield interview candidates, potentially benefiting both applicants and programs.

Program signaling was implemented in EM via the Electronic Residency Application Service (ERAS) in the 2022–2023 residency application cycle, allowing applicants to send five signals at the time of their residency application submission with instruction to not signal their home or away-rotation institutions.⁴ The Association of American Medical Colleges (AAMC) published generic guidance for programs regarding the use of PS only during the interview-offer phase and programs attested to a code of conduct regarding signal usage when opting into the process, including guidance not to use PS in rank order list (ROL) decisions.⁵ While data was evaluated by ERAS across all participating specialties, and other specialties have reported their own specialty-specific data, opportunities remained to further investigate questions specific to PS within EM.^{6–18} The unique challenges facing EM created an appetite and underscored the need for specialty-specific guidance.

To provide evidence-based guidance, the ERAS Application Working Group, a subset of the Council of Residency Directors in EM (CORD EM) Application Process Improvement Committee, created a survey to address more nuanced EM-specific questions not asked or answered by the AAMC survey. Our objective in this study was to determine how EM program directors (PD) used PS in their application review and ranking practices during the 2022–2023 application cycle, particularly in relation to the proportion of signaled applications received. To our knowledge, no other specialties participating in PS have reported PS utilization data in this manner. We also explored the relationship between program characteristics and the number of signals received, including characteristics not previously studied by the AAMC such as geographic location, program length of training, program environment, and program longevity. Lastly, we investigated the relative importance and utilization of signals in comparison to other residency application elements and in relation to the number of signals received.

Population Health Research Capsule

What do we already know about this issue?
Program signaling (PS) was introduced into the emergency medicine (EM) residency application process in 2022–2023 via the Electronic Residency Application Service.

What was the research question?
How did EM program directors use PS in application review and ranking?

What was the major finding of the study?
52.2% of program directors used PS in holistic review. Other uses varied by proportion of signaled applications.

How does this improve population health?
Understanding PS usage patterns helps inform PS allocation and usage on an individual applicant and program level.

METHODS

Study Design

We used a cross-sectional study design. Participants were PDs in Accreditation Council for Graduation Medical Education (ACGME)-accredited EM residency programs participating in the 2023 National Resident Matching Program Match. The CORD member directory, cross-referenced with the ACGME Accreditation Data System public search website, was used to compile the email distribution list. We edited the list to reflect new PDs when possible (277). The survey was created following a thorough literature review and synthesis of background information. Questions were iteratively reviewed by experts in EM medical education. The survey was further refined after conducting two cognitive interviews with EM residency program leaders and then piloted by several EM educators to assess for clarity of the questions. Data was primarily quantitative. No identifying information was collected. The study was designed to take about 10 minutes to complete. Our survey tool is included in [Appendix 1](#). This study was approved by the institutional review board at the institution of authors TF and TS.

Data Collection

The survey link was distributed via email. We collected data using a confidential and secure web-based (Qualtrics, Provo, UT) survey of EM residency PDs or their designees. Anonymous links were created for each potential respondent and distributed via Qualtrics. As described by Dillman and

colleagues, one week prior to distribution of the survey link, PDs received a brief email introducing the study and informing them that they would receive the study link in the coming week.¹⁹ Participants then received a message containing the survey link. Non-responders received up to three reminder messages over five weeks.

Data Analysis

Data was downloaded from REDCap, hosted at Maine Medical Center, directly into SPSS for Windows v 27 (IBM SPSS Statistics for Windows, IBM Corp., Armonk, NY) statistical software for analysis. We used descriptive statistics to describe the characteristics of study participants' residency training programs. Program practices and experiences around PS were described using numbers and percentages for each categorical variable. We summarized continuous variables using measures of central tendency (mean or median) and dispersion (standard deviation, interquartile range [IQR]). Comparisons between groups for categorical variables were made using chi-square analysis or the Fisher exact test. Comparisons for continuous variables were made using the *t*-test for independent samples or analysis of variance. We accepted a *P*-value of <0.05 as significant. We also computed differences between groups and their associated 95% confidence intervals (CI) and created visual data displays to aid in interpretation.

RESULTS

Program Characteristics

We received 113/277 surveys (response rate 41%). Participants represented diverse geographic regions, with the largest numbers from the Middle Atlantic, East North Central Midwest, and South Atlantic regions (Table 1). Programs represented were most commonly urban, university-based, and three years length of residency training. Faculty at participating programs were largely university or hospital employees, and most programs reporting being founded more than 15 years.

Program Signaling Participation and Applications Received

The majority of respondents participated in the PS component of the ERAS supplemental application during the 2022–2023 residency application cycle (106, 94%). Reasons for non-participation included not signing up in time (three, 2.7%), feeling that it would not contribute to applicant review or interview offer decisions (two, 1.8%), and being a newly approved program (1, 0.9%). Programs interviewed to fill a mean and median of 12 postgraduate year (PGY)-1 spots (range 6–26 spots, IQR 8–15). The number of signals received by participating programs ranged from 2–203, with a mean of 60 and median of 50 (IQR 23–86). Programs reported receipt of between 283–1,400 applications (mean 768, median 772, IQR 600–926). The proportion of applications that were signaled ranged

from 0.7% to 26.5% (mean 7.3%, median 6.5%, IQR 3.9–10.1%).

There was a moderate, positive correlation between the number of signals and the number of applications received ($r = 0.581$, $P < 0.001$) and the proportion of signals received increased based upon the number of applications received ($P < 0.001$) as well as the proportion of applications that were signaled ($P < 0.001$). The number of signals received increased as the number of PGY-1 positions increased ($P < 0.001$). Four quartiles were determined for the number of program signals received, the number of applications received, and the proportion of applications signaled (Supplemental Table 1) to allow for further comparison of data as subsequently detailed.

Signals Received by Program Characteristics

The number of signals received differed significantly based on several key characteristics: geographic location, with greater numbers of signals received in coastal regions ($P < 0.01$); program duration, with four-year receiving more than three-year programs ($P < 0.01$); program type, with urban programs receiving the most ($P < 0.01$); program environment, with university-based programs receiving the most ($P < 0.01$); and longevity of programs with programs with existence >15 years receiving the most ($P < 0.01$). Additional detail is provided in Figures 1 and 2 and Supplemental Figure 1.

Signal Utilization

Programs most commonly endorsed using PS as one component of holistic review (59, 52.2%). Additional specific ways that signals were used include the following: as a tiebreaker between two equally qualified candidates (45, 39.8%); as a screening tool (44, 38.9%); to help prioritize the program's wait list or wait list order (31, 27.4%); and to send an interview invitation to every applicant who signaled the program (19, 16.8%). The proportion of applications that were signaled appeared to affect the frequency with which programs endorsed using signals to prioritize the wait list ($P < 0.001$), serve as a tiebreaker ($P < 0.001$), and to send interview invitations to every signaling applicant ($P = 0.03$) (Figure 3). Participants anticipated using PS in the 2023–2024 cycle similarly to their reported use in the 2022–2023 cycle, and similar differences were also noted for anticipated use based on the proportion of applications that were signaled.

Signal Importance

Participants rated the importance of various application elements when considering interview invitations and preparing their program's rank order list (ROL) using a 5-point scale (1 = not important at all, 5 = extremely important) (Table 2). Participants rated the standardized letter of evaluation (SLOE) as the most important element

Table 1. Characteristics of participating residency programs and survey respondents.

Characteristic	% (n)	Comparison to existing program data (percentage of programs)
Professional role		
*Program director	100 (113)	
Geographic region		
Middle Atlantic	24.8 (28)	23.7 ^a
East North Central Midwest	20.4 (23)	20.5 ^a
South Atlantic	17.7 (20)	19.1 ^a
Pacific West	11.5 (13)	10.6 ^a
West South Central	11.5 (13)	9.9 ^a
New England	5.3 (6)	4.2 ^a
Mountain West	4.4 (5)	3.9 ^a
West North Central Midwest	2.7 (3)	3.9 ^a
East South Central	1.8 (2)	4.2 ^a
Program length		
Three years	77.0 (87)	80.6 ^b
Four years	23.0 (26)	19.4 ^b
Program environment		
Urban	63.7 (72)	Not available
Suburban	30.1 (34)	Not available
Rural	6.2 (7)	Not available
Program type		
University-based	47.8 (54)	35.4 ^a
Community-based, university-affiliated	36.3 (41)	46.2 ^a
Community-based	15.9 (18)	18.4 ^a
Faculty employment model		
University or hospital	73.5 (83)	Not available
Contract management group	18.6 (21)	Not available
Democratic physician-led group	8.0 (9)	Not available
Program longevity		
<5 years	17.7 (20)	Not available
5–10 years	8.0 (9)	Not available
10–15 years	10.6 (12)	Not available
>15 years	63.7 (72)	Not available

*261/277 EM programs participated in PS for 2022–2023. All 277 programs surveyed.


Middle Atlantic = NJ, NY, PA; East North Central Midwest = IL, IN, MI, OH, WI; South Atlantic = DC, DE, GA, FL, MD, NC, SC, VA, WV, PR; Pacific West = AK, CA, HI, OR, WA; West South Central = AR, LA, OK, TX; New England = CT, MA, ME, NH, RI, VT; Mountain West = AZ, CO, ID, MT, NM, NV, UT, WY; West North Central Midwest = IA, KS, MN, MO, ND, NE, SD; East South Central = AL, MS, KY, TN.

^aFellowship and Residency Electronic Interactive Database (FREIDA), <https://freida.ama-assn.org>


^bEmergency Medicine Residents' Association (EMRA) Match Database, <https://match.emra.org/>

when reviewing applications (mean 4.97, 95% CI 4.93–5.00). The SLOEs (mean 4.90, 95% CI 4.83–4.97) and interview day performance (mean 4.81, 95% CI 4.72–4.89) were most important when preparing the ROL. Importance of the presence or absence of a program signal when reviewing applications was a mean of 2.9 (95% CI 2.67–3.13) and median of 3 (2,4). Importance of the presence or absence of a

program signal when preparing a ROL was a mean of 2.1 (95% CI 1.87–2.32) and median of 2 (1–3). About 30% of participants (28) endorsed the presence or absence of a program signal as very or extremely important when reviewing applications while 11% (10) rated program signals as being equally important to ROL development.



(A)



(B)

Figure 1. (A) Mean number of signals received by geographic region. (B) Median number of signals received by geographic region. Geographic regions include: East North Central Midwest (IL, IN, MI, OH, WI), East South Central (AL, MS, KY, TN), Middle Atlantic (NJ, NY, PA), Mountain West (AZ, CO, ID, MT, NM, NV, UT, WY), New England (CT, MA, ME, NH, RI, VT), Pacific West (AK, CA, HI, OR, WA), South Atlantic (DC, DE, GA, FL, MD, NC, SC, VA, WV, PR), West North Central Midwest (IA, KS, MN, MO, ND, NE, SD), and West South Central (AR, LA, OK, TX).

We assessed for differences in PDs' relative assessments of various application elements based on the proportion of applications that were signaled ([Supplemental Figure 2](#)). As the proportion of applications signaled increased, the

proportion of programs endorsing board scores as "extremely important" decreased ($P < 0.01$). As the proportion of applications signaled increased, the proportion of programs endorsing communication before the interview




Figure 2. Mean number of signals received by program characteristics. (A) Mean number of signals received by program duration. (B) Mean number of signals received by environment type. (C) Mean number of signals received by program type. (D) Mean number of signals received by program longevity.




Figure 3. Program signal use in the 2022–2023 academic year by the proportion of applicants signaled.*

*The AAMC Code of Conduct, which programs attest to when signing up to participate in program signaling (PS), specifically prohibits the use of PS in rank-order list discussion and preparation.

Table 2. Importance of application elements.

Application element	Importance when reviewing applications	Importance when preparing rank order list
	Mean (95% CI)	Mean (95% CI)
SLOEs	4.97 (4.93–5.00)	4.90 (4.83–4.97)
Interview day interactions	N/A	4.81 (4.72–4.89)
Prior work or life experiences	3.61 (3.42–3.80)	3.52 (3.32–3.72)
Board scores	3.47 (3.27–3.66)	3.14 (2.93–3.35)
MSPE	3.44 (3.24–3.65)	3.32 (3.12–3.53)
Extracurricular involvement	3.36 (3.17–3.54)	3.25 (3.05–3.45)
Presence or absence of a program signal	2.90 (2.67–3.13)	2.10 (1.87–2.32)
Communication before interview	2.64 (2.42–2.87)	2.89 (2.65–3.13)
Research experience	2.46 (2.27–2.64)	2.43 (2.24–2.62)
Letters of recommendation	2.40 (2.22–2.58)	2.33 (2.15–2.52)

*5 point scale where 5 = extremely important and 1 = not important at all.

CI, confidence interval; SLOE, standardized letter of evaluation; MSPE, medical student performance evaluation.

as “not important at all” increased while the proportion rating this factor “very important” decreased ($P < 0.01$). Extracurricular involvement increased in importance as the number of applications signaled increased, with a larger proportion of participants rating this aspect of the application “extremely important” as the proportion of applications signaled increased ($P = 0.04$). Programs with the lowest proportion of signaling applicants were more likely to rate research experience as “not important at all” than those who had a larger proportion of applications signaled ($P = 0.02$).

DISCUSSION

Responses to our survey appear to be appropriately representative of programs nationwide with regard to geographic distribution, program length, and program type (Table 1).^{20,21} Ranges and median numbers for applications and PS data are similar to ERAS data, again demonstrating that our survey respondents reflected a representative sample of EM programs that participated in PS during the studied application cycle.⁶

For data analysis, we used quartiles based on the percentage of signaling applications a program received to correct for the differences in raw numbers based on program size. With the number of signals allocated to each EM applicant increasing from five to seven for the 2023–2024 academic year, it is reasonable to presume that the raw number and percentage of signaling applicants programs receive will also proportionally increase. This discrepancy may make it more difficult for a program to accurately identify with a given quartile based on this year’s application data, but these data should still serve as a rough guide by which programs can assess themselves.

Understanding the relationship between program characteristics and the number of received program signals can be helpful for both programs and applicants. Programs can determine their competitiveness within the context of similar programs, which can be particularly helpful in the current EM match environment with a changing applicant demographic pool and many programs going unmatched over the past few years.² Providing programs with a barometer against which to measure their own demographics and proportion of signaled applicants early in the application cycle can help guide how they incorporate program signals into their approach and more effectively select applicants who will be highest yield for their programs. By understanding signaling trends as related to program characteristics, advisors and applicants may be able to strategically determine the best approach for allocating signals to maximize each signal’s impact.

In our study, we noted that the Pacific West and New England regions demonstrated the highest mean and median signal numbers. In contrast, programs in the East South Central, Mid-Atlantic, West South Central, and West North Central Midwest received fewer signals. It is reasonable to speculate that many of these patterns reflect overall population density patterns, suggesting local preferences that mirror the US population. This hypothesis aligns with our data, which showed that more urban (likely more population-dense) programs received a higher proportion of signals. The only region that does not fit this hypothesis is the Mid-Atlantic region, which is the most densely populated in the country, but we suspect the very high EM program density in this region likely contributed to program signal dilution, leading to lower signals per program.

On average, four-year programs received a higher proportion of signaling applicants than three-year programs. While program length itself may be a driver of this, it may also be due to other confounding features more commonly associated with four-year programs, including urban location, university affiliation, and program duration and stability. Ultimately, our data was unable to discern this difference. Programs with the lowest proportion of signaling applicants were more likely to be smaller, rural, and not academically affiliated. These programs were more likely to rate research experience as “not important at all.” We suspect that these smaller, more community-oriented programs may be less research-focused in their missions and, therefore, emphasize research less in their applicant selection. Applicants may be able to use this information to target their signals depending on their interests.

It seems intuitive that the proportion of signaling applicants a program receives would affect how that program values and uses the signal, but to our knowledge this is the first data to demonstrate that effect. When examining signaling use among programs separated into quartiles based on the proportion of signaling applicants, significant differences emerged. Programs that received lower proportions of signaling applicants were more likely to report offering interviews to all signaling applicants while those with the highest proportion of signaling applicants were more likely to incorporate signals as a screening tool or to help prioritize the program’s wait list or wait-list order.

By asking programs to rate the importance of various application elements, we hoped to gain an understanding of the relative importance of PS in relation to interview offers and ROL creation. Receiving a program signal in orthopedics was ranked among the most important factors in resident selection for interview.¹³ While a successful sub-internship at the PD’s institution and letters of recommendation were the highest-ranked criteria for resident selection for interview at urology programs, 81% of urology PDs reported that a lack of a signal would negatively impact interview offer chances for an applicant.¹⁸ In our study, program signals were not shown to hold as much weight as in orthopedics or urology. Program signals were only rated as more important than narrative letters of recommendation, pre-interview communication, and research experience.

How an applicant performs clinically (SLOEs, Medical Student Performance Evaluation) is understandably most important, with PS intended to be only one small part of the holistic application review.²² Students can be reassured that the traditionally valued portions of the EM application retain their importance well above the value of a program signal, and programs across all quartiles are interviewing and ranking students who did not send them a signal.

Analyzing this data in a more granular fashion, we did observe some significant differences in the relative importance of residency application elements between quartiles. As the proportion of signaling applicants increased, the proportion of participants endorsing board scores as “extremely important” decreased. This discrepancy may speak to the intended ability of PS to mitigate the use of filtering behavior. Programs with smaller proportions of signaling applicants may continue to seek out strategies to stratify their applicant pool to better allocate their holistic review efforts, such as using board score filters. Programs with a higher proportion of signaling applicants, on the other hand, may not feel this same pressure. Alternatively, it is possible that having been prompted by the introduction of PS to investigate programs before applying, applicants may strategically have chosen to target their signals to programs that advertised a lack of board score cutoffs because their score fell below stated cutoffs at other programs or because they valued programs that do not emphasize standardized test scores.

Our data also demonstrates that as the proportion of signaling applicants increased, the proportion of respondents rating pre-interview communication as “extremely important” decreased and the proportion of respondents rating pre-interview communication as “not important at all” increased. This trend suggests that the signal is serving its intended purpose of allowing the applicant to meaningfully express interest, obviating the need for additional, extra-application communication, lessening the burden for both applicants and programs. It also suggests that PS reduces the impact of other communication from applicants.

The AAMC guidance was consistent in its messaging that program signals were only to be used during the application review and interview-offer portion of the application cycle. It is worth noting that despite all programs having attested in the code of conduct not to use PS in the consideration of ROL placement, 11% of programs reported program signals to be very important to the ROL development process. The 2022–23 AAMC PD survey found similar results among PD respondents from all specialties.⁶ Program directors may be extrapolating that a student who signaled is likely to be a higher probability match than a student who did not send a signal. This use presumes that student preference will not be significantly affected by their experiences engaging with programs throughout the interview season and is at risk of being flawed logic. However, it is important that applicants be aware that signals may be used by PDs in this manner and should take this into consideration when choosing where to signal.

Participation of EM programs in PS remained robust for the 2023–2024 cycle, with 278 of 279 programs participating and 97.5% of applicants participating (email communication from AAMC ERAS Pilot Administration Director, Jayme Bograd, January 2024).²⁴ We hope that

this data helps inform programs and applicants on a more nuanced approach to PS in the EM residency application process.

LIMITATIONS

Respondents (113) compared to the total number of ACGME-accredited EM residency programs (277) was limited. The PDs who chose to respond may differ from those who did not concerning their PS experience. Forty-six percent of EM programs did not fill in the 2023 Main Residency Match.²⁵ Our survey was distributed in the weeks that followed. The PDs experiencing a difficult Match cycle may have been more or less inclined to fill out a survey regarding the residency application process. University-based programs were over-represented. Community-based, university-affiliated programs were under-represented. The 11% of programs that reported using signals as part of their ROL discussions may be an underestimate as other programs may not have been comfortable disclosing behavior that was knowingly in violation of the code of conduct.

CONCLUSION

This study provides detailed data and patterns of signal use yielding insights into program signaling in EM's inaugural year for both programs and applicants. Our data provides a more nuanced understanding of signal utilization across a spectrum of EM programs in a way that allows individual programs to go beyond the general AAMC recommendations and compare their approach to that of programs with similar characteristics. Identifying patterns of signal use based on program characteristics can also inform advising for students deciding on how to best allocate their signals. As EM continues to navigate fluctuations in its applicant numbers and shifting demographics of its applicant pool, providing insight to guide signal use and utilization can help pave a path forward for the specialty toward the goal of more efficiently finding the right applicant for the right program.

Address for Correspondence: Alexis Pelletier-Bui, MD, Cooper Medical School of Rowan University / Cooper University Hospital, Department of Emergency Medicine, 401 Haddon Ave., Education & Research Building, 2nd Floor, Camden NJ 08103.
Email: pelletier-bui-alexis@cooperhealth.edu

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. No author has professional or financial relationships with any companies that are relevant to this study. There are no conflicts of interest or sources of funding to declare.

Copyright: © 2024 Pelletier-Bui et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. Pelletier-Bui AE, Schnapp BH, Smith LG, et al. Making our preference known: preference signaling in the emergency medicine residency application. *West J Emerg Med.* 2021;23(1):72–5.
2. Association of American Medical Colleges. ERAS preliminary data as of January 3 each season. 2024. Available at: <https://www.aamc.org/media/6231/download?attachment>. Accessed April 20, 2024.
3. Ramsay N. EM match 2020 by the numbers. 2020. Available at: <https://www.emra.org/students/newsletter-articles/em-match-2020-by-the-numbers/>. Accessed April 20, 2024.
4. Pelletier-Bui A, Werley E, Camejo M. EMRA hangouts - preference signaling 101. 2022. Available at: <https://www.emra.org/be-involved/events-activities/emra-hangouts/20220721-preference-signaling>. Accessed April 20, 2024.
5. Association of American Medical Colleges. Supplemental ERAS® application: guide for residency programs. 2022. Available at: <https://www.aamc.org/media/56451/download/>. Accessed April 20, 2024.
6. Association of American Medical Colleges. Supplemental ERAS® 2022–2023 application cycle: evaluation of program signaling. 2023. Available at: <https://www.aamc.org/media/64591/download>. Accessed April 20, 2024.
7. Association of American Medical Colleges. Supplemental ERAS® 2022–2023 application cycle: results of the program director reaction survey. 2023. Available at: <https://www.aamc.org/media/64996/download>. Accessed April 20, 2024.
8. LaFemina J, Rosman IS, Wallach SL, et al. The relationship between program and applicant characteristics with applicant program signals in the 2022 residency recruitment cycle: findings from 3 specialties. *Acad Med.* 2024;99(4):430–6.
9. Rosenblatt AE, LaFemina J, Sood L, et al. Impact of preference signals on interview selection across multiple residency specialties and programs. *J Grad Med Educ.* 2023;15(6):702–10.
10. Banks E, Winkel AF, Morgan HK, et al. Program signaling in obstetrics and gynecology residency applications. *Obstet Gynecol.* 2024;143(2):281–3.
11. Benjamin WJ, Lenze NR, Bohm LA, et al. Impact of applicants' characteristics and geographic connections to residency programs on preference signaling outcomes in the match. *Acad Med.* 2024;99(4):437–44.
12. Sergesketter AR, Song E, Shammass RL, et al. Preference signaling and the integrated plastic surgery match: a national survey study. *J Surg Educ.* 2024;81(5):662–70.
13. Suresh KV, Covarrubias O, Mun F, et al. Preference signaling survey of program directors-after the match. *J Am Acad Orthop Surg.* 2024;32(5):220–7.

14. Kotlier JL, Mihalic AP, Petrigliano FA, et al. Understanding the match: the effect of signaling, demographics, and applicant characteristics on match success in the orthopaedic residency application process. *J Am Acad Orthop Surg*. 2024;32(5):e231–9.
15. Cai F, Southworth E, Santiago S, et al. The golden tickets: impact of preference signaling on obstetrics and gynecology residency applicants. *Am J Obstet Gynecol*. 2024;230(2):262.e1–9.
16. Grauer R, Ranti D, Greene K, et al. Characterization of applicant preference signals, invitations for interviews, and inclusion on match lists for residency positions in urology [published correction appears in *JAMA Netw Open*. 2023;6(2):e233305]. *JAMA Netw Open*. 2023;6(1):e2250974.
17. Chang CWD, Thorne MC, Malekzadeh S, et al. Two-year interview and match outcomes of otolaryngology preference signaling. *Otolaryngol Head Neck Surg*. 2023;168(3):377–83.
18. Rodriguez-Alvarez JS, Munoz-Lopez C, Harwood S Jr., et al. Urology residency applicant selection: program directors' new criteria. *Urology*. 2024;S0090–4295(24):00141–9.
19. Dillman DA. *Mail and Internet surveys: The Tailored Design Method: 2007 with New Internet, Visual and Mixed-Mode Guide*. 2nd ed. Hoboken, NJ: John Wiley & Sons, 2007.
20. American Medical Association. Fellowship and residency electronic interactive database (FREIDA™). Available at: <https://freida.ama-assn.org>. Accessed April 20, 2024.
21. Emergency Medicine Residents' Association (EMRA). EMRA match database. Available at: <https://match.emra.org/>. Accessed April 20, 2024.
22. National Resident Matching Program. Results of the 2021 NRMP program director survey. 2021. Available at: <https://www.nrmp.org/wp-content/uploads/2021/11/2021-PD-Survey-Report-for-WWW.pdf>. Accessed April 20, 2024.
23. Association of American Medical Colleges. Residency programs participating in program signaling. 2023. Available at: <https://students-residents.aamc.org/applying-residencies-eras/residency-programs-participating-program-signaling>. Accessed April 20, 2024.
24. Association of American Medical Colleges. Exploring the relationship between program signaling & interview invitations across specialties: 2024 ERAS preliminary analysis. 2024. Available at: <https://www.aamc.org/media/74811/download?attachment>. Accessed April 20, 2024.
25. National Resident Matching Program. Results and data: 2023 main residency match. 2023. Available at: <https://www.nrmp.org/wp-content/uploads/2023/05/2023-Main-Match-Results-and-Data-Book-FINAL.pdf>. Accessed April 20, 2024.

Emergency Medicine Milestones Final Ratings Are Often Subpar

Diane L. Gorgas, MD*
 Kevin B. Joldersma, PhD[†]
 Felix K. Ankel, MD[‡]
 Wallace A. Carter, MD[§]
 Melissa A. Barton, MD[†]
 Earl J. Reisdorff, MD[†]

*Ohio State University Wexner Medical Center, Department of Emergency Medicine, Columbus, Ohio

[†]American Board of Emergency Medicine, East Lansing, Michigan

[‡]Regions Hospital, Department of Emergency Medicine, St. Paul, Minnesota

[§]Weill Cornell Medicine, Department of Emergency Medicine, New York, New York

Section Editors: Kendra Parekh, MD and Abra Fant, MD

Submission history: Submitted January 16, 2024; Revision received June 14, 2024; Accepted July 8, 2024

Electronically published August 16, 2024

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: [10.5811/westjem.18703](https://doi.org/10.5811/westjem.18703)

Background: The emergency medicine (EM) milestones are objective behaviors that are categorized into thematic domains called “subcompetencies” (eg, emergency stabilization). The scale for rating milestones is predicated on the assumption that a rating (level) of 1.0 corresponds to an incoming EM-1 resident and a rating of 4.0 is the “target rating” (albeit not an expectation) for a graduating resident. Our aim in this study was to determine the frequency with which graduating residents received the target milestone ratings.

Methods: This retrospective, cross-sectional study was a secondary analysis of a dataset used in a prior study but was not reported previously. We analyzed milestone subcompetency ratings from April 25–June 24, 2022 for categorical EM residents in their final year of training. Ratings were dichotomized as meeting the expected level at the time of program completion (ratings of ≥ 3.5) and not meeting the expected level at the time of program completion (ratings of ≤ 3.0). We calculated the number of residents who did not achieve target ratings for each of the subcompetencies.

Results: In Spring 2022, of the 2,637 residents in the spring of their last year of training, 1,613 (61.2%) achieved a rating of ≥ 3.5 on every subcompetency and 1,024 (38.8%) failed to achieve that rating on at least one subcompetency. There were 250 residents (9.5%) who failed to achieve half of their expected subcompetency ratings and 105 (4.0%) who failed to achieve the expected rating (ie, rating was ≤ 3.0) on every subcompetency.

Conclusion: When using an EM milestone rating threshold of 3.5, only 61.2% of physicians achieved the target ratings for program graduation; 4.0% of physicians failed to achieve target ratings for any milestone subcompetency; and 9.5% of physicians failed to achieve the target ratings for graduating residents in half of the subcompetencies. [West J Emerg Med. 2025;26(1.2)11–14.]

INTRODUCTION

With the advent of the Next Accreditation System (NAS), the Accreditation Council for Graduate Medical Education (ACGME) introduced a new assessment process called the “milestones.”¹ The milestones are objective behaviors that reflect elements of the major competencies (eg, patient care, systems-based practice) in thematic domains called “subcompetencies” (eg, emergency stabilization, patient- and family-centered communication). The milestone scale

uses nine ratings from 1.0, 1.5, 2.0, 2.5, etc, to 5.0. The scale is predicated on the assumption that a rating (level) of 1.0 corresponds to an incoming emergency medicine (EM)-1 resident and a rating of 4.0 is the graduation “target,” albeit not a graduation expectation or requirement. According to the ACGME: “Level 4 is designed as a graduation *goal* but *does not* represent a graduation *requirement*.”² The EM milestones have been used exclusively as a formative assessment by the ACGME. Likewise, a physician’s EM

milestone ratings are not considered when determining the eligibility of a physician to take the American Board of Emergency Medicine (ABEM) written qualifying examination.

The EM milestones were introduced in 2012, and the first ratings were reported in 2013.³ The EM milestones were revised in 2021, resulting in 22 subcompetencies. Since 2012, substantial validity evidence for the EM milestones has been accumulated.⁴⁻¹⁰ A resident's milestone ratings are usually assigned by clinical competency committees (CCCs). Some subcompetency ratings are below target levels. Often, the subcompetency ratings assigned by the CCCs are lower than the ratings that residents give themselves.¹¹ The milestones were initially designed to have a rating of 4.0 as the target for a resident completing an EM residency.⁹ Aggregate EM milestones are reported annually by the ACGME.¹² These data and other reports suggest that a substantial number of graduating residents are not achieving a level 4 rating in many milestone subcompetencies.

We undertook this study to determine the frequency with which graduating residents received the target milestone rating.

METHODS

Study Design

This retrospective cross-sectional study was a secondary analysis of an already de-identified dataset used in a prior study.¹³ Our current study was deemed exempt from human subject research by the Western-Copernicus Group Institutional Review Board. The dataset available to the investigators did not include physician or program characteristics that would allow a more detailed analysis.

Study Setting and Population

We analyzed milestone subcompetency ratings from Spring 2022 for categorical EM residents in their final year of training. These milestone ratings were submitted between April 25–June 24. This ratings report used EM Milestones 2.0, which included 22 subcompetencies. The dataset had been provided earlier to ABEM by the ACGME as part of the routine EM milestones secure data-sharing process.

Measurements or Key Outcome Measures

The primary measure was the number of subcompetencies for which physicians failed to achieve a target rating of 3.5 at the time that the Spring milestone ratings were submitted to the ACGME. Because the ratings were submitted between April and June prior to residency completion, and the CCC could have determined the ratings even earlier than that, an expected rating for purposes of the study was modified to be 3.5 rather than 4.0. Doing so assumed that the resident would achieve a rating of 4.0 over the remaining weeks to months of residency training. We determined the number of physicians

who did not achieve the target rating for the subcompetencies (from 0 subcompetencies to all 22 subcompetencies).

Data Analysis

Ratings were dichotomized as meeting the target level at the time of program completion (≥ 3.5) and not meeting the target level at the time of program completion (≤ 3.0). We calculated the number of competencies for which a target rating was not achieved.

RESULTS

In Spring 2022, there were milestone ratings for 2,637 residents in the Spring of their last year of training in 279 EM residencies. There were 1,613 residents (61.2%) who achieved a rating of ≥ 3.5 on every subcompetency and 1,024 residents (38.8%) who failed to achieve a rating of ≥ 3.5 on at least one subcompetency. There were 250 physicians (9.5%) who failed to meet half of their target subcompetency ratings. There were 105 residents (4.0%) who failed to meet the target rating (ie, rating was ≤ 3.0) on every subcompetency (Table).

Table. The frequency of emergency medicine residents receiving target milestone ratings lower than 3.5 in Spring 2022 (n = 2,637).

Number of ratings lower than 3.5	Number of terminal-year residents	Percent of total
0	1613	61.2
1	235	8.9
2	155	5.9
3	97	3.7
4	77	2.9
5	68	2.6
6	39	1.5
7	35	1.3
8	21	0.8
9	22	0.8
10	15	0.6
11	10	0.4
12	15	0.6
13	16	0.6
14	19	0.7
15	12	0.5
16	9	0.3
17	15	0.6
18	11	0.4
19	19	0.7
20	14	0.5
21	15	0.6
22	105	4.0

LIMITATIONS

First, the actual level of subcompetency achievement at graduation was imprecisely known. We chose a rating of ≥ 3.5 to represent the performance target, given that the milestone ratings were provided prior to the completion of the program. Using a rating of 4.0 to be assigned two months prior to graduation would likely underestimate subcompetency achievement and a score of 3.5 at two months prior to program completion would likely overestimate subcompetency achievement. Anticipating that all residents with a rating of 3.5 would achieve a rating of 4.0 within weeks was a benevolent assumption. Second, demographic data on residents (eg, gender) and program characteristics (eg, duration of training) were unavailable to the investigators. Although this lack of additional information limited our ability to determine factors associated with the ratings, we believe that the findings are sufficiently significant on their merit and warrant additional investigation.

Third, we did not correlate poor subcompetency ratings with program extension or remediation, thus limiting the opportunity to gather any evidence of predictive or consequential validity. It is possible that nearly every physician who did not achieve a rating of ≥ 3.5 on nearly half of the milestone subcompetencies underwent remediation. Fourth, the ratings are assigned by CCCs. The structures of, and information used by CCCs, vary by EM residency.^{14,15} We did not attempt to determine the reliability or accuracy of the individual ratings. Moreover, we did not examine the potential impact of bias in the ratings. Prior studies suggested that women were assigned lower performance ratings.^{16,17} Sixth, the ratings used for this study were from the first year of the EM Milestones 2.0. Although there was a degree of acclimation in developing facility with the EM Milestones 1.0, it is likely that the same degree of unfamiliarity would be less with the most recent version. The degree to which the continued use of EM Milestones 2.0 will change rating trends is unknown.

DISCUSSION

This study is the first in EM to demonstrate the degree to which physicians completing EM residencies are not achieving target subcompetency ratings. These data showed that of the 2,637 residents in their last year of training, nearly one in ten failed to meet target ratings for half of the EM subcompetencies. A similar finding was reported for physicians completing pediatric EM fellowships.¹⁸ However, that report used a target rating of 4.0, not 3.5 as in our study. Consequently, 67% of pediatric EM fellows did not attain a rating of at least 4.0 for at least one subcompetency.

A physician should be able to graduate from residency without scoring 4.0 on all 22 subcompetencies. In fact, all 4.0 ratings (a straight-line score) would be highly improbable.¹⁹ Consider the hypothetical situation that would result from

the milestones being used in a summative manner to determine ABEM board eligibility. If residents were required to have no more than six subpar (ie, <3.5) milestone ratings (more than one-fourth of the subcompetencies), then 353 residents (13.4%) in their final year of training would not be eligible to take the ABEM written qualifying examination. Given the intent of the milestones as a formative instrument, ABEM maintains the position that the milestones should not be used as a summative determinant of board eligibility.

The rate of program extension by physicians beyond a scheduled graduation date has been reported to be approximately 8.5%.¹³ These extensions include physicians undergoing academic remediation, as well as program extensions due to a personal leave of absence. The prevalence of physicians not meeting half of the target subcompetency ratings was 9.5%. Based on these findings, there were physicians who failed to meet at least half of the EM milestone subcompetencies yet were deemed competent to practice autonomously as attested by the program director. This likelihood does not challenge the construct validity of the milestones, nor does it suggest that the target is too high. In a fact, a prior validity study by Korte et al used program director survey data to verify the appropriateness of the target ratings.⁹

In this study we did not analyze the impact of training length (EM1-3 vs EM1-4). However, a review of mean scores was undertaken in a prior investigation that used the same study period.¹³ The scores suggest that residents in EM1-3 programs tended to have higher scores through the postgraduate years (PGY) 1–3. For example, in the PGY-3 year, residents from EM1-3 programs had a mean rating of 3.51 (95% confidence interval [CI] 3.50–3.53) and residents from EM1-4 programs had a mean rating of 3.07 (95% CI 3.05–3.09), while EM4 residents had a mean rating of 3.67 (95% CI 3.65–3.69).

This analysis is an initial exploration into a more thorough investigation of the final milestones rating that an EM resident receives. The current study does not identify variable impact within demographic groups, nor does it provide any indices of predictive validity. Given the findings of this analysis, a more thorough analysis of the milestones should be undertaken to determine their psychometric qualities and subsequent utility in the field. Given the use of the milestones as a formative evaluation system, it should not be used to make summative decisions such as the determination of ABEM board eligibility. A more structured, valid, and reliable process for making the summative determination that a physician has demonstrated the necessary competencies to practice safely and independently is advisable. Moreover, such a detailed summative process could also be used to make a confident determination that a physician is eligible for board certification. This process would be easily accommodated in a model of competency-based medical education.

CONCLUSIONS

Many physicians complete an EM residency without meeting a target rating for a graduating resident in up to half of the EM milestones. Some residents (4%) did not meet a target rating in any milestone. These findings support the continued use of the milestones as a formative instrument, rather than a tool to determine board eligibility.

Address for Correspondence: Diane L. Gorgas, MD, American Board of Emergency Medicine, 3000 Coolidge Road, East Lansing, MI 48823. Email: diane.gorgas@osumc.edu

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. Drs. Gorgas, Ankel, and Carter are members of the American Board of Emergency Medicine (ABEM), Board of Directors. Drs. Joldersma, Barton, and Reisdorff are employees of ABEM. ABEM receives no revenue from Emergency Medicine Milestones reporting. There are no other conflicts of interest or sources of funding to declare.

Copyright: © 2024 Gorgas et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- Nasca TJ, Philibert I, Brigham T, et al. The next GME accreditation system—rationale and benefits. *N Engl J Med*. 2012;366(11):1051–6.
- Emergency Medicine Milestones. The Accreditation Council for Graduate Medical Education. 2021. Available at: <https://www.acgme.org/globalassets/PDFs/milestones/emergencymedicinemilestones.pdf>. Accessed August 4, 2023.
- Beeson MS, Carter WA, Christopher TA, et al. Emergency medicine milestones. *J Grad Med Educ*. 2013;5(1 Suppl 1):5–13.
- Beeson MS, Carter WA, Christopher TA, et al. The development of the emergency medicine milestones. *Acad Emerg Med*. 2013;20(7):724–9.
- Beeson MS, Holmboe ES, Korte RC, et al. Initial validity analysis of the emergency medicine milestones. *Acad Emerg Med*. 2015;22(7):838–44.
- Hamstra SJ, Cuddy MM, Jurich D, et al. Exploring the association between USMLE scores and ACGME milestone ratings: a validity study using national data from emergency medicine. *Acad Med*. 2021;96(9):1324–31.
- Hamstra SJ, Yamazaki K, Barton MA, et al. A national study of longitudinal consistency in ACGME milestone ratings by clinical competency committees: exploring an aspect of validity in the assessment of residents' competence. *Acad Med*. 2019;94(10):1522–31.
- Holmboe ES, Yamazaki K, Nasca TJ, et al. Using longitudinal milestones data and learning analytics to facilitate the professional development of residents: early lessons from three specialties. *Acad Med*. 2020;95(1):97–103.
- Korte RC, Beeson MS, Russ CM, et al. The emergency medicine milestones: a validation study. *Acad Emerg Med*. 2013;20(7):730–5.
- Peck TC, Dubosh N, Rosen C, et al. Practicing emergency physicians report performing well on most emergency medicine milestones. *J Emerg Med*. 2014;47(4):432–40.
- Goldflam K, Bod J, Della-Giustina D, et al. Emergency medicine residents consistently rate themselves higher than attending assessments on ACGME milestones. *West J Emerg Med*. 2015;16(6):931–5.
- Edgar Laura, Hogan SO, Holmboe ES, et al. Milestones: National Report. 2022. <https://www.acgme.org/globalassets/pdfs/milestones/1160-ACGME-Milestones-Report-2022-R22.pdf>. Accessed August 4, 2023.
- Beeson MS, Barton MA, Reisdorff EJ, et al. Comparison of performance data between emergency medicine 1–3 and 1–4 program formats. *J Am Coll Emerg Physicians. Open* 2023;4(3):e12991.
- Doty CI, Roppolo LP, Asher S, et al. How do emergency medicine residency programs structure their clinical competency committees? A survey. *Acad Emerg Med*. 2015;22(11):1351–4.
- Goyal N, Folt J, Jaskulka B, et al. Assessment methods and resource requirements for milestone reporting by an emergency medicine clinical competency committee. *Med Educ Online*. 2018;23(1):1538925.
- Santen SA, Yamazaki K, Holmboe ES, et al. Comparison of male and female resident milestone assessments during emergency medicine residency training: A national study. *Acad Med*. 2020;95(2):263–8.
- Dayal A, O'Connor DM, Qadri U, et al. Comparison of male vs female resident milestone evaluations by faculty during emergency medicine residency training. *JAMA Intern Med*. 2017;177(5):651–7.
- Roskind CG, Leonard K, Baghdassarian A, et al. Pediatric emergency medicine fellows' milestone evaluations: do they all meet the targets for graduation? *AEM Educ Train*. 2021;5(3):e10620.
- Beeson MS, Hamstra SJ, Barton MA, et al. Straight line scoring by clinical competency committees using emergency medicine milestones. *J Grad Med Educ*. 2017;9(6):716–20.

Emergency Department Slit Lamp Interdisciplinary Training Via Longitudinal Assessment in Medical Practice

Samara Hamou, BA*
 Shayan Ghiaee, MD, MS[†]
 Christine Chung, MD[‡]
 Maureen Lloyd, MD[‡]
 Kelly Khem, MD[§]
 Xiao Chi Zhang, MD, MS[§]

*Sidney Kimmel Medical College, Philadelphia, Pennsylvania
[†]Department of Emergency Medicine, Emory University, Atlanta, Georgia
[‡]Department of Ophthalmology, Wills Eye Hospital, Philadelphia, Pennsylvania
[§]Department of Emergency Medicine, Thomas Jefferson University, Philadelphia, Pennsylvania

Section Editors: Jules Jung, MD and Andrew Ketterer, MD

Submission history: Submitted October 20, 2023; Revision received March 21, 2024; Accepted June 7, 2024

Electronically published August 16, 2024

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.18514

Introduction: Eye emergencies make up nearly 3% of US emergency department (ED) visits. While emergency physicians (EP) should diagnose and treat these ophthalmologic emergencies, many trainees report limited ocular exposure and insufficient training throughout their residency to confidently conduct a thorough slit-lamp exam.

Methods: We created an interdisciplinary, simulation-based mastery learning (SBML) curriculum to teach emergency attending physicians how to operate the slit lamp with multimodal learning methodology at a tertiary academic center. The EPs first demonstrate their initial slit-lamp competency with a 20-item checklist, and they then review the necessary curricular content to pass their independent readiness test before completing their in-person teaching and demonstration session with an ophthalmology attending to demonstrate procedural mastery (minimal passing score >90%).

Results: Fifteen EPs were enrolled; all completed the final exam of the curriculum. The pre- and post-curriculum checklist scores increased by an average of seven points ($P = .002$); 86.7% of EPs felt confident in completing a slit-lamp exam after the curriculum, compared to 20% at the beginning. Five of 15 reported teaching learners within the two-month post-curricular period, ranging from 5–30 students. The hands-on teaching was the most positively reviewed element of the curriculum.

Conclusion: The SBML program successfully trained EPs on performing a comprehensive slit-lamp exam with promising results of downstream education to junior learners. We encourage other institutions to leverage SBML as a teaching modality for procedural-based training and advocate cross-discipline education initiatives. [West J Emerg Med. 2025;26(1.2)15–24.]

INTRODUCTION

The slit-lamp¹ (Figure 1A) is a microscope that allows for a detailed examination of the anterior eye segment using light beam manipulation. The slit-lamp enables physicians to diagnose anterior ophthalmic pathologies such as corneal injuries, iritis, hyphema, hypopyon, and foreign bodies²; furthermore, it is essential for performing detailed ophthalmologic exam techniques such as lid eversion, fluorescein examination, and foreign body removal.³ The

Wood's lamp⁴ (Figure 1B), in contrast, is a handheld device often used to characterize skin pigmentation, dermal infections, and macroscopic infections with a built-in magnifying lens and ultraviolet (UV) light. The UV capabilities can highlight fluorescein staining during external ocular exams to assess corneal pathologies at lower magnification. While the Wood's lamp offers a less detailed examination than the slit lamp, it is a more portable diagnostic tool for larger ocular lesions, foreign bodies, or

specific reaction to fluorescein staining and meets the needs of the emergency physician (EP) under certain situations.

Eye emergencies make up nearly 3% of US emergency department (ED) visits, the most common of which are traumatic.^{5,6} The most common eye injury evaluated in the ED is corneal abrasion (superficial injury to the cornea) and eyelid laceration. Such injuries are best viewed under high-field magnified viewing using the slit lamp to assess for concomitant injuries or co-infections such as corneal ulcers, hypopyon/endophthalmitis, retained foreign body, full thickness corneal laceration, globe ruptures, and seidel testing.⁷ Ocular emergencies such as traumatic globe rupture, ocular foreign body, closed-angle glaucoma, and endophthalmitis are visible only using the slit lamp, and fall within the EP's scope of practice for diagnosis, triaging, and management.⁸ Mismanaged ophthalmic emergencies can result in inappropriate consultation, excessive testing, financial burden, and even irreversible vision loss.⁹ Despite the significance and frequency of ocular emergencies across the US, many EPs are not confident performing a detailed ophthalmic exam.¹⁰

Previous literature has found EPs receive fewer than 10 hours of ophthalmic education during residency with low confidence in performing a comprehensive ophthalmic slit-lamp exam.¹¹ Ophthalmic education through clerkships and didactics in medical school is also in decline, leading to the unpreparedness of incoming residents before any formal residency training.^{11,12} However, it is important that EPs be confident in using the slit lamp to appropriately triage and manage ocular emergencies as part of the Accreditation Council for Graduate Medical Education (ACGME) Emergency Medicine (EM) Milestones Patient Care domain (PC8) – General Approach to Procedures, which designates a set of sequential milestones for overall procedural competency, not focusing on a specific list of procedures.¹³

The optimal learning environment for adult learners to perform a technically challenging procedure should incorporate elements from both the mastery learning model and rapid cycle deliberate practice (RCDP). The mastery

learning model ensures that students can master a topic if they receive unlimited time and support in learning and reviewing material until mastery proficiency is reached. Meanwhile, the RCDP model ensures learners can practice skills repetitively while receiving brief, interspersed feedback to achieve a designated proficiency level before proceeding to the next task.^{14,15-17} Within medical education, simulation-based mastery learning (SBML) models have been successfully implemented across various specialties, such as emergency medicine, general surgery, critical care, and gastroenterology.^{18,19,20} In light of successful, smaller scaled studies on the effectiveness of slit-lamp training within undergraduate medical education, we propose a SBML procedural training curriculum that can enable adult learners to conduct deliberate performances of intended cognitive or psychomotor skills in sequential order with a repetitive skills assessment.^{15,21,22} Specific, informative feedback will enable sustained performance improvement to achieve slit-lamp mastery.²³ Our goal was to design a pilot interdisciplinary course that could teach EPs to complete a comprehensive slit-lamp exam in diagnosing common anterior eye pathology.

METHODS

Our study, Emergency Department Slit Lamp Interdisciplinary Training via Longitudinal Assessment in Medical Practice (ED SLIT LAMP), is a multicentered, collaborative project that leverages the conceptual frameworks of the mastery learning model and RCDP to ensure proficiency in conducting a comprehensive slit lamp exam. It also serves as a scaffold for deconstructing barriers in traditional siloed medical practices and leads to improved patient care, knowledge synthesis, and resource utilization of our consulting services. The study was conducted at Thomas Jefferson University (TJUH) and the Wills Eye Hospital (WEH) from 2021–2023. The hospitals with their respective EDs, are 0.2 miles apart, with staff from each institution working as consultants at the other; WEH residents function as ophthalmology consultation for the TJUH ED, while TJUH EPs function as overnight medical emergency




Figure 1. Slit lamp (A) and Wood's lamp (B).

consultants at the WEH ED. The geographic and relationship proximity created ideal conditions to develop and pilot a procedural skill competence SBML curriculum.

Emergency physicians were selected as ideal learners due to their level of training and unique teaching responsibilities. Using the TJUH ED listserv we recruited eligible participants and offered staggered financial incentives. For this pilot study, we required a minimum of 12 participants to meet 5% type 1 error and 80% power based on score improvement from baseline testing to post-testing, as referenced by Miller et al.²⁴ The ED SLIT LAMP study leveraged talents from content and education experts from both institutions to create an interdisciplinary procedural teaching curriculum. The success of a traditional SBML curriculum is linked to the learners' skill acquisition. Our study expands this measure to include interdisciplinary collaboration, demonstrating the successful alignment between educational and patient-centered goals that benefit both departments. To evaluate the curriculum, we employed all four levels of the Kirkpatrick model. Using pre- and post-test Likert scale questionnaires, our measurement of success included improved learner confidence (level 1), knowledge acquisition (level 2), willingness of learners to incorporate their skillset in clinical practice (level 3), and dissemination of this knowledge to junior learners (level 4). Any curricular feedback and improvements were extracted for future curricular iterations.

A needs-based analysis conducted at TJUH ED revealed EPs desired hands-on slit-lamp education and training on identifying anterior segment ophthalmic complaints. Since ophthalmology is a recognized component of the American Board of Emergency Medicine exam content, we constructed the pre-test clinical content based on critical and common ocular diagnoses, the most common WEH ED ophthalmology discharge diagnoses, and clinical identifications deemed "can't miss" by the ED and ophthalmology department.

All curricular contents (lecture materials, video recording, pre-post assessments, study surveys, mastery learning checklist) were created by the principal investigator [XCZ] with ophthalmology co-investigators consultation [CC, MEL] based on targeted needs assessment. These materials underwent sequential review by select experts at WEH and were modified sequentially until a consensus was reached. The minimal passing checklist score was determined to be 90%, based on combined determination from ophthalmologist experts at WEH and similar threshold determined by Miller et al.²⁴ Each curriculum assessment (Appendix A) was constructed to mirror the natural knowledge, skills, and attitude progression from the ACGME EM Milestones Patient Care Domain (PC8). Due to the multifaceted nature of EM, there is no specific procedural milestone for performing a slit-lamp exam, as described in detail in the ACGME Ophthalmology PC1:

Data Acquisition - Basic Ophthalmology Exam and Testing (Level 1).¹³ However, the EM PC8 milestones provide structured language applicable to many ED procedures and advanced device-assisted medical examinations (ie, slit-lamp exam). Please see Table 1 for the correlation between the EM milestone and ED SLIT LAMP assessments.

The longitudinal curriculum included four unique time points (Time 0–3) of intervention staggered over six months (Appendix A, Appendix B). At Time 0, participants completed an in-person baseline slit-lamp exam that was video-recorded and reviewed by two independent investigators [XCZ] [MEL]. At Time 1, the participants gained access to an asynchronous learning packet that consisted of a PowerPoint presentation on common ED eye complaints, digital library links to the WEH Manual, slit-lamp checklist, and a video recording of a comprehensive slit-lamp examination.²⁵ The participants also gained access to an independent readiness assessment (IRAT), which was required to be completed within 30 days with a minimum score of 90% before proceeding to the next in-person phase of the study (Appendix A).

Upon achieving the passing IRAT score, they were invited to participate in the Time 2 (in-person) SBML portion of the study where they were to complete an in-person demonstration of a comprehensive slit-lamp exam by a board-certified ophthalmologist [CC] on a standardized patient volunteer. Following the demonstration, participants were given unlimited time for RCDP with brief, interspersed feedback under the observation and teaching from the ophthalmologist. Participants were required to complete a minimum 18 of 20 checklist items to achieve mastery (Appendix B). Upon completing the final checklist, the participants were asked to complete a course evaluation and learner confidence survey (Appendix C) with Likert scaling, subjective commentary, and a validated 5-item Critical Incidence Questionnaire (CIQ) for curricular improvement. Given the unpredictability nature of the "unlimited attempts" at Time 2, all participants were scheduled at two-hour intervals to allow for device preparation, one to two re-attempts, debriefing, survey completion, and general troubleshooting. At Time 3, participants completed a 60-day post-examination survey, assessing their ocular knowledge, slit-lamp confidence, clinical teaching opportunities, and relevant interprofessional relationships.

We used a Wilcoxon signed-rank test to differentiate the checklist scores between the curricular intervention by incorporating collected paired data before and after the training, median and interquartile range values of subtotal scores at two-time points.²⁶ We used McNemar's test to comparing each categorical sub-score (Yes/No) by time points and corresponding *P*-value within the same population.²⁷ The descriptive summaries of survey questions at Time 0, Time 2, and three-month follow-up were analyzed using Bonferroni adjusted *P*-values (multiplying *P*-value

Table 1. Corresponding emergency department slit-lamp assessments to ACGME EM* milestone general approach to procedures.

ACGME EM milestone PC8	Bolded PC8 elements relatable to performing a slit lamp exam	Correlating ED SLIT LAMP assessments
Level 1	Identifies indications for a procedure and pertinent anatomy and physiology. Performs basic therapeutic procedures (eg, suturing, splinting)	Appendix A—Part II (clinical image examination)
Level 2	Assesses indications, risks, benefits, and alternatives and obtains informed consent in low- to moderate-risk situations. Performs and interprets basic procedures, with assistance. Recognizes common complications	Appendix B—Part I (slit lamp technical) Appendix B (final checklist)
Level 3	Assesses indications, risks, and benefits and weighs alternatives in high-risk situations. Performs and interprets advanced procedures, with guidance. Manages common complications	Appendix A—Part III (ophthalmology exam mix-n-match)
Level 4	Acts to mitigate modifiable risk factors in high-risk situations. Independently performs and interprets advanced procedures. Independently recognizes and manages complex and uncommon complications	Appendix B (final checklist)
Level 5	Teaches advanced procedures and independently performs rare, time-sensitive procedures. Performs procedural peer review	Appendix C—ED SLIT LAMP surveys

*ACGME EM, Accreditation Council for Graduate Medical Education Emergency Medicine; PC, patient care; ED SLIT LAMP, Emergency Department Slit Lamp Interdisciplinary Training.

from Wilcoxon signed-rank test by the number of multiple tests, doubling the P -values), which was directly compared to the pre-specified 5% significance level. All statistical analyses were performed using R 4.1.2 (R Foundation for Statistical Computing, Vienna, Austria).²⁸

This study was approved by the institutional review board at Thomas Jefferson University Hospital (TJUH) in Philadelphia, PA. Informed consent was obtained from participating physicians. This study was funded by the Center for Faculty Development and Nexus Learning Pedagogy Grant at Thomas Jefferson University.

RESULTS

Fifteen EPs (six females and nine males) were enrolled in ED SLIT LAMP during the two-year period; none were lost to follow-up. All participants were board-certified EPs with an average clinical experience of 7.8 years post-residency graduation. All EPs completed the final exam of the curriculum in one attempt and all under 60 minutes.

Table 2 lists the 20 steps of the slit-lamp exam curriculum checklist, comparing participant results from recorded slit-lamp attempts (Time 0) to the final in-person assessment (Time 2). The intra-class correlation in test scores between EPs and ophthalmologists at Time 0 (2 raters) was 0.98. We found a significant increase between the checklist scores before and after the education initiative, 12.0 to 19.0, $P = 0.002$.

The most notable differences between the pre- and post-curricular intervention were as follows: 1) instructing the patient to close their eyes while powering up and positioning

the patient in the slit lamp with the forehead touching the horizontal bar and chin in the chinrest ($P < 0.001$); 2) adjusting the microscope 90 degrees to facial plane with illumination set at a 45-degree angle ($P = 0.008$); 3) performing an anterior chamber evaluation ($P = 0.002$); 4) looking for cells and flare ($P = 0.021$); and 5) placing fluorescein in the inferior fornix of the eye ($P = 0.031$). The most missed steps at the baseline exam were: 1) applying a transparent face shield (26.7%); 2) instructing patients to close their eyes when the machine was turned on (26.7%); 3) looking for cells and flare (26.7%).

Figures 2 and 3 illustrate learners' confidence in performing and teaching the slit-lamp exam at the beginning of the study (Time 0), immediately after achieving procedural mastery (Time 2), and two months later (Time 3). Figure 4 illustrates the learners' likelihood in teaching the slit-lamp exam at Time 0 and Time 2. Before participating in the slit-lamp curriculum, 73% of EPs also reported rarely or never performing a slit-lamp exam, while 80% of EPs reported sometimes or often using a Wood's lamp for ocular complaints. Only 20% of EPs reported feeling confident in performing and teaching a comprehensive slit-lamp exam, while 67% of EPs reported feeling confident in using and teaching Wood's lamp for ocular examination.

After completing the slit-lamp curriculum (Time 2), 86.7% of EPs reported feeling confident performing a comprehensive slit-lamp exam for ocular complaints, and 73.3% were more confident in teaching residents how to perform a slit-lamp exam. Most EPs strongly agreed that the ED SLIT LAMP curriculum helped them perform an

Table 2. Descriptive summary of checklist evaluation at pre- and post-curricular and comparison between time points.

Checklist item	Performed	Time 0, N(%) (N = 15)	Time 2, N(%) (N = 15)	P-value from exact McNemar's test
1 - Identify slit lamp anatomy.	Yes	13 (86.7%)	15 (100%)	0.50
2 - Apply transparent face shield over the slit lamp (COVID).	Yes	4 (26.7%)	15 (100%)	<0.001
3 - Sanitize forehead and chin rest for the patient.	Yes	5 (33.3%)	14 (93.3%)	0.004
4 - Apply topical tetracaine/proparacaine on patient's eyes.	Yes	8 (53.3%)	12 (80.0%)	0.22
5 - Unlock instrument base and shift by pulling toward you.	Yes	15 (100%)	15 (100%)	NA
6 - Adjust eye pieces for your interpupillary distance and refractive error.	Yes	10 (66.7%)	14 (93.3%)	0.22
7 - Adjust table height and/or chair(s) - neither patient nor examiner should be hunched over.	Yes	12 (80.0%)	14 (93.3%)	0.50
8 - Instruct patient to close eyes while you power up by turning on the light source at low voltage setting and focus on right eyelid. Position patient in slit lamp with forehead touching the horizontal bar and chin in the chin rest.	Yes	4 (26.7%)	15 (100%)	<0.001
9 - Set magnification on lowest settings (10x to 12x), illumination at largest aperture and widest slit beam.	Yes	12 (80.0%)	15 (100%)	0.25
10 - Adjust chin rest so the patient is sitting comfortably with their chin on the chinrest and their forehead against the headrest.	Yes	12 (80.0%)	15 (100%)	0.25
11 - Practice macro and micro adjustments of the sliding base with joystick.	Yes	14 (93.3%)	15 (100%)	1.00
12 - Adjust microscope 90° to facial plane with illumination set at 45° angle (angle left for patient's right eye, and right for left eye).	Yes	7 (46.7%)	15 (100%)	0.008
13 - Perform outer structure evaluation.	Yes	14 (93.3%)	15 (100%)	1.00
14 - Perform anterior chamber evaluation.	Yes	5 (33.3%)	15 (100%)	0.002
15 - Look for cells and flare.	Yes	4 (26.7%)	12 (80.0%)	0.02
16 - Place a drop of tetracaine/proparacaine on a sterile fluorescein strip.	Yes	15 (100%)	15 (100%)	NA
17 - Place the fluorescein in the inferior fornix of the eye by pulling down on the lower lid and gently touching the bulbar conjunctiva with the fluorescein strip.	Yes	9 (60.0%)	15 (100%)	0.03
18 - Adjust cobalt blue filter on diaphragm wheel at maximum beam height and medium width slit setting for fluorescein evaluation.	Yes	14 (93.3%)	15 (100%)	1.00
19 - Focus the slit beam at 9:00 position on limbus. Move across the cornea to the 3:00 position by tilting joystick laterally.	Yes	12 (80.0%)	15 (100%)	0.25
20 - Pull instrument base toward you when finished and lock in position. Turn off.	Yes	4 (26.7%)	13 (86.7%)	0.004
		Time 0, median [IQR]	Time 2, median [IQR]	P-value from Wilcoxon signed rank test
Subtotal score		12.0 [10, 16]	19.0 [19, 20]	0.002

IQR, interquartile range.

independent slit-lamp exam and identify critical findings for common ocular complaints (80%), enhancing their learning more than traditional lectures and reading alone (86.7%). Of the asynchronous materials, the video demonstration was the most used (53% used it “a lot” or a “great deal”); the

PowerPoint lecture and WEH Manual were the least used. At two months post-ED SLIT LAMP (Time 3), 73% and 67% of participants expressed extreme confidence in performing and teaching a resident how to perform a slit-lamp exam. Five of 15 EPs reported teaching learners within the two-month




Figure 2. Learner confidence in performing the slit-lamp exam at Time 0 (pre-curricular), Time 2 (immediate post-SBML curriculum), and Time 3 (2-month post-SBML curriculum).




Figure 3. Learner confidence in teaching the slit-lamp exam at Time 0 (pre-curricular), Time 2 (immediate post-SBML curriculum), Time 3 (2-months post-SBML curriculum).

post-curricular period, ranging from 5–30 students per EP participant.

Table 4 summarizes the statistically significant findings from the survey responses based on the three timeframes. There was a statistically significant increase in self-reported confidence in 1) performing a comprehensive slit lamp exam and 2) teaching residents to perform this exam between Time 0 to Time 2 and Time 0 to Time 3 ($P < 0.001$). There was no difference in reliance on ophthalmology consultation to modify or reinforce a treatment plan for ocular complaints when comparing Time 0 to Time 3 ($P = 0.70$, $P = 0.814$). There was also no statistical difference in the number of patients with ocular complaints evaluated by the study

participants at the TJUH ED and WEH ED throughout the study ($P = 0.14$, $P = 1.00$).

DISCUSSION

The ED SLIT LAMP curriculum allowed EPs to increase their use and confidence in performing slit-lamp exams in the ED. The impetus for the project arose from EPs' intrinsic motivation to provide better patient care. Our participant population consisted primarily of junior faculty who were initially uncomfortable performing or teaching slit-lamp exams and preferred using the Wood's lamp. Upon completing the curriculum, the EPs noted a significant increase in self-reported confidence in using slit-lamps




Figure 4. Learner likelihood in teaching the slit-lamp exam at Time 0 (pre-curricular) and Time 2 (immediate post-SBML curriculum).

and were teaching multiple junior learners during their study enrollment.

The improvement between the pre-and post-curricular procedural competency also demonstrates the importance of understanding the technical nuances of the slit-lamp exam and practicing critical device movement, such as careful patient positioning, adjusting of the chin straps, changing the microscope angulation, and adjusting varying slit-lamp beam lengths and widths for diagnosing a wide range of anterior ocular pathologies. These skills are drastically different than those required to operate a Wood's lamp, which acts primarily as a magnifying glass with UV capabilities.

Our curriculum achieved three of the four Kirkpatrick goals. The majority of the participants (over 80%) reported positive reaction to the curriculum (the curriculum helped them perform a slit-lamp exam, evaluate for common pathologies, and offered more than traditional lectures) (Level 1); all of the participants demonstrated procedural mastery at Time 2 (Level 2); upwards of 50 learners received instructions from the study participants on how to use the slit lamp at Time 3 (Level 3). While the reliance on ophthalmology consultation did not reveal statistically significant changes, we posit that improved procedural acumen resulted in more targeted consultation questioning and improved rapport between the medical disciplines.

Since our participants were board-certified EPs with limited availabilities, the most valued component of the curriculum was the in-person RCDP session with the ophthalmologist (Time 2). This was reflected in almost every CIQ item, with specific mention of direct guidance in positioning the beam to look for cells and flare. The most surprising element to many participants was how many ocular diagnoses required the slit-lamp exam and that learning the procedure was not as complicated as they had

initially anticipated. In contrast, many of the participants felt most distanced or removed from the curriculum in reviewing the asynchronous learning materials.

We were unsurprised to see the confidence levels in using Wood's lamp unchanged between the three different time frames. While the slit lamp offers a superior and in-depth evaluation of the anterior segment of the eye, we acknowledge that a comprehensive slit-lamp exam is time- and resource-consuming and may not affect the clinician's management if the suspected pathology involves larger lesions, foreign bodies, or specific reaction to fluorescein staining. The Wood's lamp remains an easier and more portable diagnostic tool for some ocular pathologies, and its use in the clinical arena is still acceptable in certain situations.

LIMITATIONS

This study was conducted at a single, large, tertiary academic center with an affiliated ophthalmology hospital and supported with internal grant funding. While the results were positive, multiple factors could prevent this study from being replicated, especially at community sites without a close relationship with ophthalmology. One of the most significant challenges is scheduling in-person evaluations in the pre-curricular session, as well as the final in-person training and examination. We encountered significant logistical challenges in creating a schedule that was amenable to the ophthalmologists, EPs (with unpredictable shift schedules), and research investigators, as well as finding a consistent space in the WEH and WEH ED that had access to an attached-observer scope to ensure the participants were focusing on the correct anatomic structure during their procedural demonstration. This was further exacerbated when accounting for the "unlimited attempts" for RCDP. As this was our pilot study with advanced learners, we

Table 3. Statistical analysis of survey questions between the three different study timeframes.

Survey question	Time 0 median [IQR] ^a	Time 2 median [IQR] ^b	Time 3 median [IQR] ^c	Bonferroni adjusted <i>P</i> -value from Wilcoxon signed rank test time 0 vs. time 2	Bonferroni adjusted <i>P</i> -value from Wilcoxon signed rank test time 0 vs. time 3
Slit lamp					
Based on your current practice patterns: how confident are you in: performing a comprehensive slit lamp exam for ocular complaints?	1 [1, 2]	4 [3, 4.5]	3 [2.5, 4]	<0.001	<0.001
Based on your current practice patterns, how confident are you in: teaching residents to perform a comprehensive slit lamp exam for ocular complaints	1 [1, 2]	3 [2.5, 4.5]	3 [2, 4]	<0.001	0.004
How often do you: perform an independent slit lamp exam for ocular complaints?	2 [1, 2.5]	n/a*	3 [3, 3]	n/a*	0.064
Wood's lamp					
Based on your current practice patterns, how confident are you in: performing a comprehensive Wood's lamp exam for ocular complaints?	4 [2, 4]	4 [4, 5]	4 [3, 5]	0.016	0.03
Based on your current practice patterns, how confident are you in: teaching residents to perform a comprehensive Wood's lamp exam (with access to a slit lamp) for ocular complaints?	4 [2, 4]	4 [4, 5]	4 [3, 5]	0.03	0.08
How often do you: use a wood lamp (with access to a slit lamp) for ocular complaints?	3 [3, 4]	n/a*	3 [3, 3]	n/a*	1.00
Ophthalmology consultation habits					
How confident are you in identifying common ocular pathology seen in your main work site (CC, MHD, Urgent Care)?	2 [2, 3]	n/a*	3 [3, 4]	n/a*	0.018
On average, how many eye pathologies do you see at the main work site?	10 [4, 15]	n/a*	5 [3, 12.5]	n/a*	0.14
On average, how many eye pathologies do you see at other facilities?	12 [0, 40]	n/a*	37.5 [13.5, 50]	n/a*	1.00
How often do you rely on ophthalmology consultation to: help modify your treatment plan for ocular complaints?	3 [3, 3]	n/a*	3 [2.5, 3]	n/a*	0.70
How often do you rely on ophthalmology consultation to: reinforce your treatment and plan for ocular complaints?	3 [2, 3]	n/a*	3 [2, 3]	n/a*	0.814
How often do you rely on ophthalmology consultation to: provide additional information and guidance to your treatment and plan for ocular complaints?	3 [3, 4]	n/a*	3 [3, 3.5]	n/a*	1.00

Confidence levels: 1 = Not at all confident, 5 = Extremely confident

Frequency levels: 1 = Never, 5 = Always

^aTime 0 = pre-curricular evaluation.

^bTime 2 = immediate post SBML exam. Frequency of slit lamp and Wood's lamp use were intentionally omitted for Time 2 due to the close proximity between Time 0 and Time 2, thus resulting in 'n/a' for some calculations.

^cTime 3 = three months after SBML exam.

CC, Jefferson Hospital in Center City Philadelphia; MHD, Jefferson Methodist Hospital; IQR, interquartile range.

over-budgeted a two-hour template for each learner, which drastically limited the number of participants we could schedule for the final in-person exam.

Due to the longitudinal nature of this study and several in-person components, maintaining participant recruitment and engagement was also difficult. Of the 50 eligible board-certified TJUH EPs, only 15 EPs volunteered to participate. The primary deterrent, when discussed with non-participants, was time restraints and commuting into the city for in-person evaluations and examinations. We suggest implementing dedicated teaching days (ie, conference days or faculty meetings) for larger participant recruitment and subsequent follow-up and examination.

This study was funded by an internal grant that provided minor financial incentives for the participants and standardized patient volunteers. While our needs-based analysis revealed participants were more focused on promoting better patient care, many of the participants expressed appreciation for the staggered gift cards, which also incentivized them to complete each timeline-specific survey. All other investigators' efforts, in contrast, were in-kind and required dedicated non-academic and non-clinical time to enroll participants, record all the interactions, and provide unrestricted time availabilities for the final mastery assessment. This study was also unanimously supported by both departmental leaderships to promote a better collegial relationship and interdisciplinary education opportunity between organizations with the two principal investigators holding unique leadership positions, ophthalmology consulting director [CC] and EM clerkship director [XCZ]. We suspect that also positively affected our recruitment process and the success of this interdisciplinary training curriculum. As this study was conducted at an academic hospital in an urban setting, it has been suggested that academic centers likely overestimate EP comfort and confidence in the diagnosis and management of ophthalmic emergencies.⁹ Furthermore, the proximity between both EDs may skew the data, as these EPs are likely exposed to fewer ophthalmic emergencies than hospitals without a nearby eye-focused ED.

Ultimately, the biggest limitation to this pilot study was the lack of in-person skill assessment at the 60-day follow-up due to limited staffing and scheduling challenges. In lieu of an objective competency score, we leveraged self-reported confidence at the 60-day mark as an approximate measurement of the skill retention. We recognize that learners are poor at gauging their own abilities, both over- and underestimating their skills based on a variety of factors. It is notable that 80% of our learners were initially "not confident" in completing a comprehensive slit-lamp exam prior to the SBML curriculum and scored an average checklist score of 60%. At Time 2, almost 87% of responders were "confident" in completing a comprehensive slit-lamp exam after receiving an average checklist score of 95%.

Unfortunately, there is no association between learners' confidence and passing rate (score >18) at Time 0 (Pearson chi-square 3.46, $P = 0.17$) and Time 2 (Pearson chi-square 0.833, $P = 0.66$), respectively. While we are unable to predict how these learners would have performed on their slit-lamp exam test at day 60, we are encouraged to see the number of study participants who continued to teach slit-lamp exam for junior learners. We posit these participants will likely have improved sustained competence and decreased skill decay by actively teaching others. Future studies should be considered to add a final examination (procedure or multiple-choice question) to validate our results.

CONCLUSION

Emergency physicians are expected to diagnose and manage ocular complaints as part of their training and clinical practice. Our primary focus was to create a rigorous methodologic training curriculum (slit-lamp exam) for a specialty-focused skillset that could result in downstream teaching. This project highlighted a significant need for slit-lamp exam training within our institution that led to a successful transdisciplinary simulation-based mastery learning curriculum and improved our EPs' confidence in performing and teaching slit-lamp exams to future clinicians. Furthermore, this study demonstrates that adult learners, especially attending physician value direct interaction with clinical instructors when learning a new skillset and are intrinsically motivated to hone their skillset and teach it to future learners when they have achieved this mastery. We encourage other institutions to leverage SBML as a teaching modality for procedural-based training and advocate cross-discipline education initiatives. Future investigation could include creating a multicenter study to implement this curriculum at other academic institutions and potentially include it in EM residency training.

ACKNOWLEDGMENTS

We want to thank the Thomas Jefferson University Center for Faculty Development and Nexus Learning (CFDNL) for their generous support and grant funding in completing this study.

Address for Correspondence: Xiao Chi Zhang, MD, Thomas Jefferson University, Department of Emergency Medicine, 1020 Sansom St., Thompson Bldg, Suite 239, Philadelphia, PA 19107. Email: Xiaochi.zhang@jefferson.edu

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. This study received grant funding from the Center for Faculty Development and Nexus Learning (CFDNL) Pedagogy, Thomas Jefferson University. There are no other conflicts of interest or sources of funding to declare.

Copyright: © 2024 Hamou et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- User: OKJaguar. File: slit lamp and binocular microscope. 2019. Available at: https://commons.wikimedia.org/wiki/File:Slit_lamp_and_binocular_microscope.jpg. Accessed August 26, 2023.
- Knoop K. Ophthalmologic procedures in the emergency department—part III: slit lamp use and foreign bodies. *Acad Emerg Med*. 1995;2(3):224–30.
- Seol SH, Cho J, Lee WJ, Choi SC. Use of a slit-lamp microscope for treating impacted facial foreign bodies in the emergency department. *Clin Exp Emerg Med*. 2015;2(3):188–92.
- User: Seawind60. File: Wood's UV lamp. 2013. Available at: https://commons.wikimedia.org/wiki/File:Wood%27s_UV_lamp.JPG. Accessed August 26, 2023.
- Babineau M and Sanchez L. Ophthalmologic procedures in the emergency department. *Emerg Med Clin North Am*. 2008;26(1):17–34.
- Cheung CA, Rogers-Martel M, Golas L, et al. Hospital-based ocular emergencies: epidemiology, treatment, and visual outcomes. *Am J Emerg Med*. 2014;32(3):221–4.
- Owens PL and Mutter R. (2011). Statistical brief #112: Emergency department visits related to eye injuries, 2008. In *Healthcare cost and utilization project (HCUP) statistical briefs*. Rockville, MD: Agency for Healthcare Research and Quality.
- The Wills Eye Emergency Department. All eyes, all the time. *Eye Level*. 2021. Available at: https://www.willseye.org/wp-content/uploads/2021/05/WEB_3009-Eye-Level-Spring-2021.pdf. Accessed October 17, 2023.
- Uhr JH, Governatori NJ, Zhang QE, et al. Training in and comfort with diagnosis and management of ophthalmic emergencies among emergency medicine physicians in the United States. *Eye*. 2020;34(9):1504–11.
- Druck J, Valley MA, Lowenstein SR. Procedural skills training during emergency medicine residency: are we teaching the right things? *West J Emerg Med*. 2009;10(3):152–6.
- Gelston CD and Patnaik JL. Ophthalmology training and competency levels in care of patients with ophthalmic complaints in United States internal medicine, emergency medicine and family medicine residents. *J Educ Eval Health Prof*. 2019;16:25.
- Graubart EB, Waxman EL, Forster SH, et al. Ophthalmology objectives for medical students: revisiting what every graduating medical student should know. *Ophthalmology*. 2018;125(12):1842–3.
- Accreditation Council for Graduate Medical Education (ACGME). Emergency medicine milestones. 2023. Available at: <https://www.acgme-i.org/globalassets/acgme-international/milestones-2.0/emergencymedicinemilestones2.0int.pdf>. Accessed October 17, 2023.
- Griswold-Theodorson S, Ponnuru S, Dong C, et al. Beyond the simulation laboratory: a realist synthesis review of clinical outcomes of simulation-based mastery learning. *Acad Med*. 2015;90(11):1553–60.
- Chancey RJ, Sampayo EM, Lemke DS, et al. Learners' experiences during rapid cycle deliberate practice simulations: a qualitative analysis. *Simul Healthc*. 2019;14(1):18–28.
- Lemke DS, Fielder EK, Hsu DC, et al. Improved team performance during pediatric resuscitations after rapid cycle deliberate practice compared with traditional debriefing: a pilot study. *Pediatr Emerg Care*. 2019;35(7):480–6.
- Mavis SC, Kreofsky BL, Ouk MY, et al. Training fellows in neonatal tele-resuscitation using a simulation-based mastery learning model. *Resuscitation Plus*. 2021;8:100172.
- Zendejas B, Cook DA, Bingener J, et al. Simulation-based mastery learning improves patient outcomes in laparoscopic inguinal hernia repair: a randomized controlled trial. *Ann Surg*. 2011;254(3):502–9; discussion 509–11.
- Franklin BR, Placek SB, Gardner AK, et al. Preparing for the American Board of Surgery flexible endoscopy curriculum: development of multi-institutional proficiency-based training standards and pilot testing of a simulation-based mastery learning curriculum for the endoscopy training system. *Am J Surg*. 2018;216(1):167–73.
- Barsuk JH, McGaghie WC, Cohen ER, et al. Use of simulation-based mastery learning to improve the quality of central venous catheter placement in a medical intensive care unit. *J Hosp Med*. 2009;4(7):397–403.
- Qureshi N. International perspective from Saudi Arabia on “procedural skills training during emergency medicine residency: are we teaching the right things?” *West J Emerg Med*. 2009;10(3):157–8.
- Hoonpongsimanont W, Nguyen K, Deng W, et al. Effectiveness of a 40-minute ophthalmologic examination teaching session on medical student learning. *West J Emerg Med*. 2015;16(5):721–6.
- Siddaiah-Subramanya M, Smith S, Lonie J. Mastery learning: how is it helpful? An analytical review. *Adv Med Educ Pract*. 2017;8:269–75.
- Miller DT, Zaidi HQ, Sista P, et al. Creation and implementation of a mastery learning curriculum for emergency department thoracotomy. *West J Emerg Med*. 2020;21(5):1258–65.
- Rutledge BK. The wills eye manual: Office and emergency room diagnosis and treatment of eye disease. *Arch Ophthalmol*. 1996;114(1):112.
- McNemar Q. Note on the sampling error of the difference between correlated proportions or percentages. *Psychometrika*. 1947;12(2):153–7.
- Conover WJ. *Practical Nonparametric Statistics*. Hoboken, NJ: John Wiley & Sons. 1999.
- R: The R Project for Statistical Computing. June 11, 2023. Accessed August 8, 2023. <https://www.r-project.org/>.

Teaching the New Ways: Improving Resident Documentation for the New 2023 Coding Requirements

Nathan Zapolsky, MD*
Annemarie Cardell, MD[†]
Riddhi Desai, DO*
Stacey Frisch, MD, MEd[‡]
Nicholas Jobeun, DO, MEd[§]
Daniel Novak, DO*
Michael Silver, MS*
Arlene S. Chung, MD, MACM^{||}

*Maimonides Medical Center, Department of Emergency Medicine, Brooklyn, New York
[†]Emory University Hospital, Department of Emergency Medicine, Atlanta, Georgia
[‡]New York University, Grossman School of Medicine, New York, New York
[§]Icahn School of Medicine at Mount Sinai, New York, New York
^{||}University of Vermont Medical Center, Burlington, Vermont

Section Editor: Benjamin Holden Schnapp, MD

Submission history: Submitted June 1, 2024; Revision received June 12, 2024; Accepted July 8, 2024

Electronically published September 19, 2024

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.21183

BACKGROUND

In January 2023, significant changes to the structure of the Current Procedural Terminology CPT(R) evaluation and management (E/M) codes (here forward called the 2023 E/M changes) were implemented for emergency department (ED) encounters. These modifications aim to lessen administrative workload and accurately match coding with contemporary patient care practices.¹ They are anticipated to impact roughly 85% of the relative value units of ED care² and, thus, also have significant financial implications for EDs. Residents provide front-line care and documentation for millions of patients seen in United States EDs annually. The Model of Clinical Practice of Emergency Medicine³ identifies financial principles, such as billing and coding, to be required core content for board certification. Furthermore, the Accreditation Council for Graduate Medical Education (ACGME) includes quality clinical documentation to be one of the milestones that determine advancement in residency training.⁴ Interventions that alleviate documentation burden are also associated with improved physician well-being per the existing literature.⁵ However, research suggests that most emergency medicine (EM) residents do not receive formal training in billing and coding and have knowledge gaps in this area.⁶⁻⁸

Historically, documentation of encounters in the ED focused on the number of elements within history of present illness (HPI), review of systems (ROS), physical exam (PE), and medical decision-making (MDM). These new coding guidelines shift the focus almost entirely to MDM. They emphasize documentation of differential diagnoses; independent interpretation of medical testing; justification of testing not pursued; social determinants of health; chronic diseases; histories; communication with consultants,

ancillary staff, and primary care; and review of external records.

OBJECTIVES

We sought to improve resident understanding of and compliance with the 2023 E/M changes. Objectives included identification of the key elements required at each E/M level, charting and receiving feedback of sample encounters, and appreciation for the importance of accurate and high-quality documentation. We secondarily sought to investigate whether our intervention improved resident wellness specifically via benefits in confidence to perform accurate and expeditious documentation and completion of charting in a timely manner.

CURRICULAR DESIGN

Our curriculum was developed using Kern's six-step approach to curriculum design⁹ as a part of the educational quality improvement process at a single EM residency program based at a single, large, tertiary-care, urban hospital with an approximate annual ED patient census of 110,000 from October 1, 2022–February 28, 2023. Prior to study initiation a needs assessment was performed. Key stakeholders in departmental billing and coding were identified and interviewed, and relevant literature was reviewed.¹⁻¹⁰ This included the hospital chief medical information officer, ED vice chair, and billing and coding leadership. The interviews revealed a shared opinion that often the documentation to reach the appropriate expected level of service (4 or 5) was lacking to support that level of billing and most of that documentation should be captured in the MDM portion of the note. Thus, the MDM portion of the note was targeted for the intervention.

Our educational methods primarily used in-person, flipped-classroom sessions. We decided to use a flipped-classroom approach for several reasons: 1) to allow residents to gain exposure to the new billing criteria prior to the in-person sessions; 2) as a mechanism to assess resident understanding and skills, both individually with homework responses as well as in a group setting; and 3) to use faculty's in-person time for oversight and feedback.¹¹ We also applied a spaced learning approach to maximize knowledge acquisition and retention.¹² The sessions were held on December 14, 21, and 28, 2022.

For pre-session homework each week, all residents were provided a sample patient HPI, ROS, and PE components. All learners were provided the same case, and cases were changed each week. Cases were formulated to include elements that could be expanded upon in the MDM. Residents were also provided with the "CPT Evaluation and Management (E/M) Code and Guideline Changes" document.¹⁰ They were then asked to create an MDM section in accordance with the above document. Homework responses were reviewed by faculty, and feedback was given individually via email. Written feedback for residents was generated using a template based largely on the 2023 E/M guidelines changes.¹⁰ An ideally documented sample MDM section was also supplied for reference (Supplement 1).¹³

During each 30-minute session, residents were divided into small groups of four and provided an example patient case, which included only the HPI, ROS, and PE components. Residents then collaboratively wrote an MDM section for the case. All groups were provided the same case, and cases were changed each week to focus on different aspects of the MDM section. Each small group of residents shared their response with the larger group and were provided peer feedback under the guidance of a faculty facilitator selected for their advanced knowledge in either education or operations. Facilitators were provided in advance with an example of an ideally documented MDM section, which residents were also provided with at the conclusion of the exercise.

IMPACT/EFFECTIVENESS

We employed a pre-post interventional study design using a convenience sample of residents, in which group assignment was based on the number of trainings each resident could attend due to scheduling factors outside the scope of this study. This study was determined to be exempt by the institutional review board of Maimonides Medical Center. Participation was voluntary and anonymous. We evaluated the impact of our brief educational intervention on subjective measures of EM resident knowledge, skills, and attitudes via survey and on objective measures of skills and behaviors by assessing aggregate chart data.

Surveys were developed through a group iterative process that included one author (ASC) with expertise in survey

design methodology. RedCap,^{14,15} hosted at [Maimonides Medical Center] was used to anonymously distribute both pre- and post-intervention surveys. Both surveys consisted of six Likert-scale questions, three regarding their reported use of documentation shortcuts, and three assessing attitudes about their own understanding of and predicted skill with the new E/M coding changes. Six additional multiple-choice questions assessed knowledge about documentation rules. A final question was for feedback and requested ideas for other E/M billing and coding education. The pre-intervention survey, distributed November 30, 2022, differed from the post-intervention survey of December 28, 2022, only in asking the self-reported number of sessions attended. (Supplement 2).

Resident skills were assessed using actual clinical documentation. Resident aggregate E/M levels were assessed across three months pre-intervention (October 1–December 31, 2022) and two months post-intervention (January 1–February 28, 2023). Due to variation in resident clinical schedules, we chose the above time periods to capture the greatest proportion of the ED encounters documented by residents. We used the Kirkpatrick model to evaluate our intervention's impact.¹⁶ Surveys were used to assess resident subjective reactions, and objective knowledge by identification of billable elements in a provided sample MDM. We used actual clinical documentation to assess changes in behavior. Specifically, we assessed whether trainees had a statistically significant increase ($P < 0.05$) in the proportion of E/M level 5 charts (99285) and likewise a significant decrease in level 1, 2, 3, and 4 charts (99281, 99282, 99283, 99284).

We used descriptive statistics and comparison of means with the Mann-Whitney U test stratified by number of educational sessions attended to analyze significant differences in knowledge and attitudes before and after the intervention. For knowledge, these calculations were summarized with median and interquartile range (IQR) and compared across time periods using an exact Wilcoxon signed-rank test. A Bonferroni correction for the significance of the intervention changes the alpha to 0.01667. For each chart level (99281–99285), we created logistic regression models using generalized estimating equations for individual repeated measures to account for personal variability. The number of attended flipped-classroom sessions was treated as the independent variable. Zero trainings were considered to be the pre-period for analysis. All analyses considered alpha ≤ 0.05 to be statistically significant and were conducted using SPSS v 28.0¹⁵ (SPSS Statistics, IBM Corp, Armonk, NY).

Forty-six of the 54 EM residents (85%) eligible for the study completed both pre- and post-intervention surveys. All 54 residents participated in at least one survey. Due to clinical schedules, some residents were not present at one or more of the three offered sessions. The first live session was attended




Figure. Proportion of each Current Procedural Terminology evaluation and management level by number of educational sessions attended.

by 33 (61%) residents, the second by 38 (70%), and the third by 40 (74%). Six (11%) residents attended one session, 15 (28%) attended two sessions, and 25 (46%) attended three sessions. Eight (15%) did not attend any sessions.

Residents demonstrated a significant improvement in knowledge regarding which elements are the key to the MDM within the 2023 E/M changes [6 (5–6.5) to 8 (7.5–8) $P < 0.001$], and by correctly identifying the number and complexity of problems, complexity of data, risk level, and the overall complexity of a sample encounter. There was no statistically significant improvement in identification of the important coding elements (4 [3–5] to 5 [3.5–5], $P = 0.38$). Residents also endorsed greater confidence in their ability to describe (2 [1–3] to 4 [3–4], $P < 0.005$), accurately document (3 [2–3] to 4 [3–4], $P < 0.005$), and bill (2 [2–3] to 3 [2–3] $P < 0.005$). There were no significant changes in their opinion of their ability to complete their charts in a timely manner ($P < 0.19$, CI 0.165–0.215) in the decision to use dictation software ($P = 1$), shortcuts ($P = 1$), or custom prepared text phrases ($P = 1$) following the intervention. Residents participating in any number of flipped-classroom sessions showed significant changes in their skills, including an increase in E/M Level 5 coded charts, and a significant decrease in Level 1, 2, and 3 coded charts ($P < 0.005$). The increase in Level 5 charts and decrease in Level 3 charts were significant after just one session (Figure). No significant change was observed in Level 4 charts.

To the best of our knowledge, this is the first study to date to describe the impact of an educational intervention on EM resident documentation knowledge, skills, and attitudes within the framework of the 2023 E/M changes. Naturally, our experience and results at our single EM residency

program based at a large, urban, tertiary-care hospital may not be generalizable. This intervention data is single center and preliminary, and the intervention should undergo repetition and comparison before firm conclusions can be drawn.

We chose to collect data during a time range tied to the same illness season to keep the case acuity mix and attending/resident staffing comparable. We otherwise could have compared to the same months of the previous year for pre-intervention data, to best match the illness season, or alternatively, post-intervention data could have been drawn instead from the following year (2024) to help mitigate recency bias in the intervention group. That being said, the major differences in resident and attending staffing between times a year apart could also have confounded results.

We considered faculty supplemental documentation and its effect on documentation outcomes during our study design and took a pragmatic approach. For the duration of this study the attending population was stable, no significant changes to attending education were performed during this period, and attending staffing remained at baseline with no changes to ratios, shift durations, or standard distributions of encounters throughout the ED care areas. To further address this concern we attempted analysis of the attending distribution between these various groups. No attending had a greater than 1.4% change in their billing from pre- to post-intervention, and their small contributions to the overall billing for each intervention group was, therefore, unlikely to have biased the large differences seen between groups. However, the differences in distribution of attending shifts between the groups varied statistically significantly, and bias

cannot be assessed without patient-level billing records. This could be considered in future studies.

Our program may have implications regarding wellness as well. Residency training must prepare emergency physicians for all aspects of their eventual professional expectations. Residents receiving education expressed greater confidence in their ability to describe, accurately document, and bill for care provided. Business literature frequently notes how a lack of clear expectations increases work stress and harms employee wellness and productivity.¹⁸ However, whether this association applies to emergency physicians deserves further study.

CONCLUSION

Overall, we observed significant improvements in resident knowledge, attitudes, skills, and behaviors regarding clinical documentation. We hope to apply these successes and lessons learned to the formation of enduring education materials at our own institution for documentation improvement for both residents and attendings.

Address for Correspondence: Nathan Zapolsky, MD, Maimonides Medical Center, Department of Emergency Medicine, 4802 Tenth Ave., Brooklyn, 11219 NY. Email: nizapo@gmail.com

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. No author has professional or financial relationships with any companies that are relevant to this study. There are no conflicts of interest or sources of funding to declare.

Copyright: © 2024 Zapolsky et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- Thought Leadership Team. 2023 E/M coding changes. 2023. Available at: <https://www.aapc.com/resources/evaluation-management-coding-changes-2023>. Accessed August 18, 2023.
- McKenzie DA and Granovsky MA. 2023 documentation guideline changes for ED E/M codes 99281–99285. 2022. Available at: www.acepnow.com/article/2023-documentation-guideline-changes-for-ed-e-m-codes-99281-99285/. Accessed January 25, 2023.
- Beeson MS, Bhat R, Broder JS, et al. The 2022 model of the clinical practice of emergency medicine. *J Emerg Med*. 2023;64(6):659–95.
- Cooney RR, Murano T, Ring H, et al. The emergency medicine milestones 2.0: setting the stage for 2025 and beyond. *AEM Educ Train*. 2021;5(3):e10640.
- Nguyen O, Jenkins N, Khanna N, et al. A systematic review of contributing factors of and solutions to electronic health record-related impacts on physician well-being. *J Am Med Inform Assoc*. 2021;28(5):974–84.
- Romans B, Nguyen E, Biddle M, et al. Billing and coding: disparities in healthcare provider training. *J Allied Health*. 2022;51(1):43–6.
- Dawson B, Carter K, Brewer K, et al. Chart smart: a need for documentation and billing education among emergency medicine residents? *West J Emerg Med*. 2010;11(2):116–9.
- Chin S, Li A, Boulet M, Howse K, et al. Resident and family physician perspectives on billing: an exploratory study. *Perspect Health Inf Manag*. 2022;19(4):1g.
- Thomas PA, Kern DE, Hughes MT, et al. *Curriculum Development for Medical Education: A Six-step Approach*, 3rd ed. Baltimore, MD: The Johns Hopkins University Press, 2015.
- American Medical Association. CPT® evaluation and management (E/M) code and guideline changes. 2022. Available at: <https://www.ama-assn.org/practice-management/cpt/cpt-evaluation-and-management>. Accessed March 24, 2024.
- Rotellar C and Cain J. Research, perspectives, and recommendations on implementing the flipped classroom. *Am J Pharm Educ*. 2016;80(2):34.
- Weidman J and Baker K. The cognitive science of learning: concepts and strategies for the educator and learner. *Anesth Analg*. 2015;121(6):1586–99.
- Allen A. Medicine clerkship (inpatient) H&P examples: H&P5. 2018. Available at: <https://www.med.unc.edu/medclerk/wp-content/uploads/sites/877/2018/10/hp5.pdf>. Accessed February 24, 2024.
- Harris PA, Taylor R, Thielke R, et al. Research electronic data capture (REDCap) – a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42(2):377–81.
- Harris PA, Taylor R, Minor BL, et al. The REDCap consortium: building an international community of software partners. *J Biomed Inform*. 2019;95:103208.
- Kirkpatrick DL. *Evaluating Training Programs: The Four Levels*. San Francisco, CA: Berrett-Koehler Publishers Inc, 1994.
- IBM Corp. IBM SPSS Statistics for Windows, Version 28.0. 2021. Armonk, NY: IBM Corp.
- Bornstein J. How a lack of clear expectations leaves employees anxious and adrift. 2023. Available at: <https://www.forbes.com/sites/forbescoachescouncil/2023/06/05/how-a-lack-of-clear-expectations-leaves-employees-anxious-and-adrift/?sh=70de0f8f5983>. Accessed April 23, 2024.

Telesimulation Use in Emergency Medicine Residency Programs: National Survey of Residency Simulation Leaders

Max Berger, MD, MEHP*
Jack Buckanavage, MD[†]
Jaime Jordan, MD, MAEd*
Steven Lai, MD*
Linda Regan, MD, MEEd[‡]

*University of California Los Angeles, David Geffen School of Medicine,
Department of Emergency Medicine, Los Angeles, California

[†]Mount Sinai School of Medicine, Department of Emergency Medicine,
New York, New York

[‡]Johns Hopkins University, Department of Emergency Medicine, Baltimore, Maryland

Section Editors: Kendra Parekh, MD and Benjamin Holden Schnapp, MD

Submission history: Submitted June 14, 2024; Revision received September 20, 2024; Accepted September 26, 2024

Electronically published October 22, 2024

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.24863

Introduction: Coronavirus 2019 (COVID-19) accelerated the need for virtual learning including telesimulation. Many emergency medicine (EM) programs halted in-person simulation and trialed telesimulation, but specifics on its utilization and plans for future use are unknown. Telesimulation has been defined as “a process by which telecommunication and simulation resources are utilized to provide education, training, and/or assessment to learners at an off-site location.” Our objective in this study was to describe the patterns of telesimulation usage in EM residency programs during COVID-19-induced learning restrictions as well as its anticipated future utility.

Methods: We identified EM simulation leaders via the EMRA Match website, institutional websites, or personal contact with residency coordinators and directors, and invited them to participate by email. Participants completed a confidential, web-based survey consisting of multiple-choice items and one free-response question, developed by our study team with consideration of survey research best practices and Messick’s validity framework. We collected data between January–February 2022. We calculated descriptive statistics for multiple-choice items and examined the free-response answers for common themes.

Results: We obtained contact information for simulation leaders at 139 EM residency programs. Survey response rate was 65% (91/139). During in-person restrictions, 62% (56/91) of programs used telesimulation. Assuming all restrictions lifted, 38% (34/90) of respondents planned to continue to use telesimulation, compared to 9% (8/91) using telesimulation before COVID-19. Most respondents planned to use telesimulation for medical knowledge (26/34, 76%) and communication/teamwork-focused cases (23/34, 68%). In response to the free-response question regarding experience with and plans for use, we identified three major themes: 1) telesimulation is a valuable alternative to in-person learning; 2) telesimulation is an option for learners unable to participate in person; and 3) telesimulation is challenging for procedural education.

Conclusion: Despite the relatively limited use of telesimulation in EM residencies prior to COVID-19, an increased number of programs have plans to continue incorporating telesimulation into their curricula. This plan for continued use opens opportunities for further innovation and scholarship within simulation education. [West J Emerg Med. 2025;26(1.2)42–47.]

INTRODUCTION

Restrictions imposed on in-person education during the coronavirus 2019 (COVID-19) pandemic accelerated the need for virtual learning, including telesimulation.^{1,2} Telesimulation has been defined as “a process by which telecommunication and simulation resources are utilized to provide education, training, and/or assessment to learners at an off-site location.”³ Initial applications were in lower resource settings such as developing countries where learners did not have access to simulation centers or instructors.^{4,5} Within telesimulation, different modalities have been described that vary in fidelity as well as location of the learners and instructors relative to each other and the simulation center.⁶⁻⁸

Several published articles since early 2020 have described different institutions’ approaches to telesimulation since the pandemic.^{1-2,9-13} Common themes include the need to modify learning objectives to virtual environments and to select the appropriate modality of telesimulation based on institutional needs and resources.⁹⁻¹² Different modalities of telesimulation have been described, including the following: 1) learners virtually observing and debriefing a live simulation⁷; 2) learners present with a manikin while instructors facilitate from a separate location⁶; 3) instructors present with a manikin while learners remotely participate⁷; and 4) completely remote option where learners and instructors both participate remotely from separate locations.^{10,11}

Limited data comparing telesimulation vs traditional simulation suggests that learner satisfaction with telesimulation or hybrid virtual and in-person simulation is similar, although this was not found in all studies.^{7,13,14} A scoping review from 2021 highlighted the mixed data on student perception of telesimulation, with some of the included studies indicating remote facilitation of simulation being perceived as equally or more effective than live facilitation, while others found remote facilitation to be inferior.¹⁴ Facilitator perception of telesimulation has not been well studied. Limited learning outcome data has suggested similar improvements between in-person simulation and telesimulation.^{8,14}

Our objective in this study was to describe the patterns of telesimulation usage in emergency medicine (EM) residency programs during COVID-19-induced, in-person learning restrictions as well as its anticipated utility moving forward. This information is crucial to understanding the value of telesimulation and its utility in medical education.

METHODS

Study Design, Setting, and Population

We conducted a cross-sectional survey study of faculty in charge of simulation for EM residency programs in the United States. We collected data from January–February 2022. After identifying EM residency programs and their websites from the EMRA Match database,¹⁵ we searched each website for contact information for the director of

Population Health Research Capsule

What do we already know about this issue?
COVID-19 accelerated the need for virtual learning including telesimulation.

What was the research question?
To what extent was telesimulation used by EM residencies during COVID-19, and what is its anticipated utility moving forward?

What was the major finding of the study?
Only 9% (8/91) of programs used telesimulation before COVID-19. During COVID restrictions, 62% (56/91) of programs used it, while after limitations were lifted, 38% (34/90) planned to continue telesimulation.

How does this improve population health?
As an adjunct to traditional in-person simulation curriculum, telesimulation is a viable option to improve medical knowledge and communication-based competencies.

simulation education. If there was no director designated, we emailed each residency’s program coordinator and/or director asking for contact information for the faculty in charge of residency simulation. Each program was allowed only one designated participant. This study was given exempt status by the University of California, Los Angeles Institutional Review Board (IRB#21-001336) and the Johns Hopkins University Homewood IRB (HIRB00013694).

Survey Development and Dissemination

Given the lack of any previously created survey applicable to this construct, the primary author (MB) developed a web-based survey tool with consideration of survey research best practices and Messick’s validity framework.¹⁶⁻¹⁹ For content validity evidence, we first performed a literature review, and the author group of expert simulation educators and medical education researchers reviewed the survey for clarity and relevance to the construct. We defined telesimulation as including any simulation activity where “telecommunication and simulation resources are utilized to provide education, training, and/or assessment to learners at an off-site location.”³ We piloted the survey with a group of simulation educators who were not included in the target sample to gather response process validity evidence. After piloting, we revised the survey for clarity. The final survey included multiple-choice items and one free-text response item (Appendix 1).

We invited participants by email and sent two targeted, follow-up invitations to non-responders at bi-weekly intervals. We administered the survey via Qualtrics (Qualtrics, LLC, Provo, UT). No individual identifying information was collected. To maximize response rate and minimize guessing, we did not require participants to complete all survey items. Participants were not compensated for participating in the study.

Data Analysis

We calculated and reported descriptive statistics for items with discrete answers. We conducted calculations using Qualtrics and Microsoft Excel for Mac (Microsoft Corp, Redmond, WA). We examined the answers to the free-text responses to identify common themes that would broaden the reader's understanding of the data. Successive wave analysis was performed to assess the extent of possible nonresponse bias.²⁰ We examined whether use of telesimulation during the pandemic, planned future use of telesimulation after in-person restrictions, and respondent program format (postgraduate years [PGY] 1–3 vs 1–4) differed by wave. Bivariate chi-square tests for each variable of interest by wave were performed using Microsoft Excel for Mac, and *P*-values less than 0.05 were considered statistically significant. We used the consensus-based checklist for reporting of survey studies (CROSS) as reporting guidelines (Appendix 2).²³

RESULTS

Of 139 simulation leaders we identified, 91 (65%) completed the survey with 87 (63%) completing all items. We report demographic data for survey respondents in Table 1, while respondents' experience and perceptions of telesimulation are shown in Table 2. Prior to the COVID-19 pandemic, 9% (8/91) of survey respondents were using telesimulation in their curricula. There was a wide variety of prior experiences with telesimulation, with the most common being that they had heard of telesimulation but never been involved (44%). Ninety-two percent (84/91) of respondents reported that their institution prohibited in-person learning activities at some point during the COVID-19 pandemic. During in-person learning restrictions, 62% (56/91) used telesimulation in some form.

When survey respondents were asked about what format(s) of telesimulation were used, 11% (10/90) stated that they only used a completely virtual oral-boards style format, while the rest of those who used telesimulation reported using a hybrid or virtual format involving a patient monitor and/or manikin. The largest percentage of survey respondents felt that medical knowledge and communication/teamwork-focused cases were best suited for telesimulation (72% and 47% respectively), while most felt that procedure training was not well suited for telesimulation (62%). Thirty-eight percent (34/90) of respondents stated they planned to use telesimulation in some form in their curriculum moving

Table 1. Survey respondent demographics.

	n (%)
Format of respondent's current residency program	
PGY 1–3	62/89 (70%)
PGY 1–4	27/89 (30%)
Size of respondent's current residency program (total number of residents in all years)	
≤20 residents	11/90 (12%)
21–40 residents	38/90 (43%)
41–60 residents	31/90 (34%)
≥60 residents	10/90 (11%)
Respondent's current residency program primary institution setting	
University-based	58/90 (64%)
Community-based	28/90 (31%)
County-based	13/90 (14%)
Prior simulation training of survey respondent	
Fellowship training in simulation	31/90 (34%)
Non-fellowship training in simulation	48/90 (53%)
No formal training in simulation	17/90 (19%)
Respondent years since residency graduation	
≤5	22/90 (24%)
6–10	33/90 (37%)
11–15	16/90 (18%)
16–20	8/90 (9%)
≥21	11/90 (12%)

PGY, postgraduate year.

forward, mostly for medical knowledge and communication/teamwork-focused cases (76% and 68%, respectively).

We received 14 free-text responses, and identified three major themes, described below with exemplar quotes.

1. Telesimulation is a valuable alternative to in-person learning:
 “It has been the ‘better than nothing’ option but accepted by learners when other options are not feasible.”
 “It has exceeded expectations in how helpful it has been.”
2. Telesimulation is an option for learners unable to participate in person:
 “We found that it's a great option for residents with families or who have other extenuating circumstances why they can't participate in person, ie, breastfeeding moms, new parents, elder care, etc. Many of our residents who are between nights or between mid-shifts will log on and participate.”
3. Telesimulation is challenging for procedural education:
 “Difficult to learn muscle memory for high acuity, low occurrence skills.”

Table 2. Key survey results.

	n (%)
EM residency program use of telesimulation	
Prior to COVID-19 pandemic	8/91 (9%)
During in-person learning restrictions	56/91 (62%)
Planned use after in-person restrictions lifted	34/90 (38%)
During any point in the COVID-19 pandemic, did your institution prohibit in-person learning activities?	
Yes	84/91 (92%)
No	7/91 (8%)
Experience with telesimulation prior to COVID-19	
Had never heard of telesimulation	17/91 (19%)
Heard of telesimulation but never involved	40/91 (44%)
Attended a presentation	20/91 (22%)
Participated as an instructor	19/91 (21%)
Participated as a learner	6/91 (7%)
Conducted a research project	5/91 (5%)
Read a paper about telesimulation	16/91 (18%)
Formats of telesimulation used during COVID-19 restrictions	
Completely virtual; utilizing real-time patient monitor and/or manikin	21/90 (23%)
Completely virtual; oral boards style cases	31/90 (35%)
Hybrid; instructor, learners and/or sim tech in sim center while others remote	31/90 (35%)
What simulation activities were best suited for telesimulation?	
Medical knowledge focused cases	65/90 (72%)
Communication/teamwork focused cases	42/90 (47%)
Procedure focused cases	5/90 (6%)
Dedicated procedure training	2/90 (2%)
Procedure training on homemade models	10/90 (11%)
What simulation activities were not well suited for telesimulation?	
Medical knowledge focused cases	0/87 (0%)
Communication/teamwork focused cases	18/87 (21%)
Procedure focused cases	52/87 (60%)
Dedicated procedure training	54/87 (62%)
Procedure training on homemade models	23/87 (26%)
Percent of future simulation curriculum involving telesimulation	
0% of the curriculum	56/90 (62%)
1–25% of the curriculum	30/90 (33%)
26–50% of the curriculum	3/90 (3%)
51–75% of the curriculum	1/90 (1%)
76–100% of the curriculum	0/90 (0%)

(Continued on next column)

Table 2. Continued.

	n (%)
Types of future simulation activities for those who plan to continue using telesimulation	
Medical knowledge-focused cases	26/34 (76%)
Communication/teamwork-focused cases	23/34 (68%)
Procedure focused cases	7/34 (21%)
Dedicated procedural training	5/34 (15%)
Procedure training on homemade models	5/34 (15%)

EM, emergency medicine.

“Procedural training was the most difficult to simulate via telesim.”

For the wave analysis, the study included 91 respondents, including 42 in wave 1 (46%), 21 in wave 2 (23%), and 28 in wave 3 (31%). Results of the examined survey questions did not statistically differ by wave with all *P*-values > 0.05. (See Supplemental Table.)

DISCUSSION

Despite relatively low use of telesimulation within EM programs prior to the COVID-19 pandemic, we found that many EM residency programs (62%) quickly adapted to in-person learning restrictions by using telesimulation. While not all programs that trialed telesimulation plan to continue its use, 38% of respondent programs do plan to continue to use telesimulation, compared to 9% of programs using telesimulation prior to COVID-19. This represents a large increase in the overall usage of telesimulation within EM residencies. Our study also sheds light on how telesimulation can benefit EM programs. Being able to increase learner participation to include residents with family obligations or between night shifts could allow for increased return on investment for simulation resources and faculty time. Most respondents who plan to continue to use telesimulation reported that they will use it as a small percentage of their overall simulation curriculum, which highlights that telesimulation is not replacing in-person simulation but augmenting the traditional curriculum. This could be in a hybrid format that allows for increased participation, or as part of separate telesimulation days that could reduce the travel burdens on learners and instructors.

There was large variation in how programs conducted telesimulation during in-person restrictions. This is in line with prior literature and likely reflects individual program needs, preferences, and available resources.^{1,2,8–11,13,22} Most described telesimulation as best suited for medical knowledge and communication/teamwork-focused cases, rather than for procedure teaching. This is interesting given that early descriptions of telesimulation in the literature mostly

involved procedural teaching.^{5,6} One possible explanation for this discrepancy is that those early studies involved duplicate simulators at remote locations, an expense that is likely not practical, or necessary, for a residency program given the ability to host procedure training as part of the in-person curriculum. While it is apparent that there are increased plans for the use of telesimulation compared to the pre-pandemic era, not all residency programs who used telesimulation during times of in-person restrictions are planning to continue to do so. The reasons for this are unknown but may relate to telesimulation resource availability or limited outcome data on its utility.

Based on our results, we believe that telesimulation can continue to be a valuable addition to the traditional in-person simulation curriculum, particularly in allowing for increased participation of learners and instructors, reducing resource costs such as simulation center and staff time, and allowing for a viable option to practice medical knowledge and communication-based competencies. Now that telesimulation has been established as an instructional strategy that will continue to be part of many EM residency curricula, it opens opportunities for future innovation and scholarship within simulation-based medical education. Additional investigation could compare different modalities of telesimulation on objective learning outcomes.²³ It would also be interesting to explore the role of virtual and augmented reality within telesimulation.^{24,25}

LIMITATIONS

Despite multiple attempts, we were not able to obtain contact information for a simulation leader from all EM programs. However, the breakdown of PGY 1–3 vs PGY 1–4 programs of survey respondents (70% PGY 1–3 vs 30% PGY 1–4), approximating the actual distribution of the EM residency programs (81% PGY 1–3 vs 19% PGY 1–4), suggests that the sample closely resembles that of the population.⁸ Given our response rate of 65%, it is possible non-response bias affected our results, with participants with less interest or familiarity with telesimulation being less likely to respond. However, the results of our successive wave analysis failed to detect non-response bias for the selected survey questions.

There may be other influences affecting a program's use of telesimulation that we were not able to capture, and in this survey study we examined only the opinions of faculty and not those of resident learners. Additionally, the literature-based definition of telesimulation we used may be overly broad and encompass more than what typical educators might consider telesimulation. Finally, we acknowledge that the survey was administered in 2022 with in-person learning restrictions just starting to be lifted, and how people are using telesimulation now may be changing. Future work could examine this evolving use of telesimulation within EM residency programs.

CONCLUSION

This study describes past and planned future use of telesimulation within EM residency programs. A large proportion of EM residencies trialed telesimulation during COVID-19-induced restrictions. Despite relatively low use of telesimulation prior to the pandemic, more EM programs plan to incorporate telesimulation moving forward as a limited portion of their overall simulation curriculum. Opportunities for further innovation and scholarship within this area of simulation education will be possible given this planned continued use.

ACKNOWLEDGMENTS

This project was completed in fulfillment of the requirements for the Johns Hopkins University Master of Education in the Health Professions.

Address for Correspondence: Max Berger, MD, MEHP, University of California Los Angeles, David Geffen School of Medicine, Department of Emergency Medicine, 1100 Glendon Ave., Suite 1200, Los Angeles, CA 90024. Email: MRBerger@mednet.ucla.edu

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. No author has professional or financial relationships with any companies that are relevant to this study. There are no conflicts of interest or sources of funding to declare.

Copyright: © 2024 Berger et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. Patel SM, Miller CR, Schiavi A, et al. The sim must go on: adapting resident education to the COVID-19 pandemic using telesimulation. *Adv Simul.* 2020;5:26.
2. Diaz MCG and Walsh BM. Telesimulation-based education during COVID-19. *Clin Teach.* 2021;18(2):121–5.
3. McCoy CE, Sayegh J, Alrabah R, et al. Telesimulation: an innovative tool for health professions education. *AEM Educ Train.* 2017;1(2):132–6.
4. Mikrogianakis A, Kam A, Silver S, et al. Telesimulation: an innovative and effective tool for teaching novel intraosseous insertion techniques in developing countries. *Acad Emerg Med.* 2011;18(4):420–7.
5. Okrainec A, Henao O, Azzie G. Telesimulation: an effective method for teaching the fundamentals of laparoscopic surgery in resource-restricted countries. *Surg Endosc.* 2010;24(2):417–22.
6. Hayden EM, Khatri A, Kelly HR, et al. Mannequin-based telesimulation: increasing access to simulation-based education. *Acad Emerg Med.* 2018;25(2):144–7.

7. McCoy CE, Sayegh J, Rahman A, et al. Prospective randomized crossover study of telesimulation vs standard simulation for teaching medical students the management of critically ill patients. *AEM Educ Train*. 2017;1(4):287–292.
8. Yasser NBM, Tan AJQ, Harder N, et al. Telesimulation in healthcare education: a scoping review. *Nurse Educ Today* 2023;126:105805.
9. Nadir NA, Kim J, Cassara M, et al. Simulation-based emergency medicine education in the era of physical distancing: lessons learned in the COVID-19 pandemic. *AEM Educ Train*. 2021;5(3):e10586.
10. Ray JM, Wong AH, Yang TJ, et al. Virtual telesimulation for medical students during the COVID-19 pandemic. *Acad Med*. 2021;96(10):1431–5.
11. Sanseau E, Lavoie M, Tay KY, et al. TeleSimBox: a perceived effective alternative for experiential learning for medical student education with social distancing requirements. *AEM Educ Train*. 2021;5(2):e10590.
12. Thomas A, Burns R, Sanseau E, et al. Tips for conducting telesimulation-based medical education. *Cureus*. 2021;13(1):e12479.
13. Austin A, Rudolf F, Fernandez J, et al. COVID-19 educational innovation: hybrid in-person and virtual simulation for emergency medicine trainees. *AEM Educ Train*. 2021;5(2):e10593.
14. Heffernan R, Brumpton K, Randles D, et al. Acceptability, technological feasibility and educational value of remotely facilitated simulation-based training: a scoping review. *Med Educ Online*. 2021;26(1):1972506.
15. Emergency Medicine Residents' Association. EMRA Match. Available at: <https://webapps.emra.org/utills/spa/match#/search/map>. Accessed July 21, 2021.
16. Rickards G, Magee C, Artino AR, Jr. You can't fix by analysis what you've spoiled by design: developing survey instruments and collecting validity evidence. *J Grad Med Educ*. 2012;4(4):407–10.
17. Gehlbach H and Artino AR, Jr. The survey checklist (manifesto). *Acad Med*. 2018;93(3):360–66.
18. Artino AR, Jr., Durning SJ, Sklar DP. Guidelines for reporting survey-based research submitted to academic medicine. *Acad Med*. 2018;93(3):337–40.
19. Messick S. Validity. In: Linn R.L, editor. *Educational Measurement*. 3rd ed. New York, NY: Macmillan; 1989:13–103.
20. Duszynski TJ, Fadel W, Dixon BE, et al. Successive wave analysis to assess nonresponse bias in a statewide random sample testing study for SARS-CoV-2. *J Public Health Manag Pract*. 2022;28(4):E685–91.
21. Sharma A, Minh Duc NT, Luu Lam Thang T, et al. A consensus-based checklist for reporting of survey studies (CROSS). *J Gen Intern Med*. 2021;36(10):3179–87.
22. Levine M, Singh M, Restivo A, et al. Recommendations for optimizing virtual simulation: a trial and error process from the COVID-19 pandemic. *J Grad Med Educ*. 2022;14(1):18–21.
23. Kirkpatrick JD and Kirkpatrick WK. *Kirkpatrick's Four Levels of Training Evaluation*. Alexandria, VA: ATD Press; 2016.
24. Mistry D, Brock CA, Lindsey T. The present and future of virtual reality in medical education: a narrative review. *Cureus*. 2023;15(12):e51124.
25. Pallavicini F, Pepe A, Clerici M, et al. Virtual reality applications in medicine during the COVID-19 pandemic: systematic review. *JMIR Serious Games*. 2022 Oct 25;10(4):35000.

Substantial Variation Exists in Clinical Exposure to Chief Complaints Among Residents Within an Emergency Medicine Training Program

Corlin M. Jewell, MD*
Amy T. Hummel, MD*†
Dann J. Hekman, MS*
Benjamin H. Schnapp, MD, MEd*

*University of Wisconsin School of Medicine and Public Health, BerbeeWalsh
Department of Emergency Medicine, Madison, Wisconsin
†Emergency Medicine Specialists SC, Wauwatosa, Wisconsin

Section Editors: Doug Franzen, MD and Andrew Ketterer, MD

Submission history: Submitted February 25, 2024; Revision received September 27, 2024; Accepted October 11, 2024

Electronically published November 19, 2024

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.20281

Introduction: While many aspects of emergency medicine (EM) residency training are standardized among residents within a single residency program, there is no standard for the distribution of chief complaints (CC) that residents should see over the course of residency. This could result in substantial variability in each resident's clinical exposure. Our objective in this study was to explore EM residents' clinical exposure to CCs to determine whether substantial variation exists. If such variation exists, this could suggest the need for curricular reform to address gaps in resident clinical exposure during training.

Methods: This was a retrospective observational study of EM residents who graduated in the years 2016–2021 at a single, university-affiliated emergency department (ED) in the midwestern United States. All patient encounters where a CC was logged were included and categorized into 1 of 20 clinical domains based on the 2016 American Board of Emergency Medicine Model of Clinical Practice. We calculated descriptive statistics for the top 10 most encountered domains for comparison among residents.

Results: We included a total of 228,916 patient encounters from 69 residents in the analysis. Residents were involved in an average of 3,323 distinct patient encounters during the study period. The overall interquartile range for patient encounters was 523. The three CC domains with the broadest interquartile variation were abdominal and gastrointestinal disorders (116), musculoskeletal disorders (nontraumatic) (93), and traumatic disorders (86).

Conclusion: Within a single, three-year academic EM program, substantial variation existed among residents with regard to the variety of patient CCs seen during their residency training. [West J Emerg Med. 2025;26(1.2)48–53.]

INTRODUCTION

Medical residency training allows physicians to gain the cognitive and procedural skills necessary to practice independently. Based on experiential learning theory, patient encounters form the foundation upon which physicians in training begin to master the practice of medicine.¹ Additionally, the development of “illness scripts,” or mental models for the classification of patient presentations, is

crucial to the development of clinical skills and reasoning during residency training.² These models are developed over time by multiple exposures to presentations of similar disease states.^{3,4} Emergency medicine (EM) trainees must be exposed to a variety of patient chief complaints (CC) throughout the course of residency to develop these scripts and become ready to begin independent practice.

Educators within EM have worked to define many aspects of EM residency training, including optimum number of shifts, on-shift educational goals/practices, and didactic content.⁵ Despite this, the clinical experience of an individual resident may be highly variable and may be partially driven by self-selection of patients by the resident. Studies in pediatric EM suggest that there is significant variation in the overall number of patients and range in acuity among individual residents.^{6,7} However, there is little adult EM literature that explores the variation in clinical experience seen by residents within a modern EM program. The literature that does exist in adult EM suggests there is substantial variation in clinical exposures among residents.⁸ A study from 2006 found that the number of cases seen overall correlated with improved performance on a standardized test designed to assess clinical competence. However, the effect plateaued at around 200 cases.⁹ Prior work by our group has shown that case volume in an individual domain did not correspond to performance within that domain on corresponding questions on the in-training exam.¹⁰

These studies suggest that individuals within a single training program may be gaining variable experience with certain types of patient presentations and lacking exposure (and therefore opportunities to develop mastery) to other complaints and pathology. However, this variability in clinical exposure during training has not been shown in adult EM for over three decades.⁸ Since then, the number of annual visits to the ED as well as the complexity of medical care provided have substantially increased.^{11,12} We, therefore, hypothesized that substantial differences in clinical exposure still exist among residents at the time of graduation. Understanding these differences is of critical importance for residency programs as considerable variation could push some residents below a threshold to develop robust illness scripts suitable for independent practice.

METHODS

Study Design and Setting

We conducted this retrospective, observational study at a three-year EM residency program situated within an urban, academic emergency department (ED) in the Midwest. The ED for the primary clinical site has a total of 54 beds and sees an annual volume of approximately 60,000 patient visits. During the study period, the residency had 12 first postgraduate year one (PGY-1) positions available each year. The study ED divides its beds into two adult clinical areas and a pediatric clinical area. All three areas are physically connected on a single floor of the hospital. Residents from all three years are assigned to nine-hour shifts in each clinical area. Each shift includes 1–2 junior (PGY-1) residents, 1–2 senior (PGY-2 or PGY-3) residents, and one attending physician. Any resident can assign themselves to patients of any severity regardless of seniority. In Fall 2020,

Population Health Research Capsule

What do we already know about this issue?
Studies from 30 years ago reported variation in the distribution of chief complaints seen by emergency medicine residents during training.

What was the research question?
We hypothesized that substantial differences in clinical exposure still exist among residents at the time of graduation.

What was the major finding of the study?
The three chief complaint domains with the most variability between individual resident experience, as measured by the greatest 25–75 interquartile ranges were abdominal and gastrointestinal disorders (median 594 patients per resident, IQR 116), nontraumatic musculoskeletal disorders (median 314, IQR 92), and traumatic disorders (median 525, IQR 86).

How does this improve population health?
Understanding these differences is important, as substantial variation could mean that some residents do not develop robust illness scripts suitable for independent practice.

the study ED shifted from a “pod” model in which the two adult clinical areas would assign themselves predominately to patients in their clinical area to a “free-for-all” model in which either adult team could assign themselves to any adult patient regardless of the clinical area they were roomed in. During the study, physician assistants were employed in the ED and would occasionally take the place of a resident on shift (particularly during weekly resident didactics).

Data Acquisition

Residents were eligible for inclusion if they had completed residency within three consecutive years and graduated in the years 2016–2021 (therefore, the study period was from June 2013–June 2021). The electronic health record (EHR) was used to create a database of patient encounters; all encounters where eligible EM residents were the first resident assigned to the patient were analyzed. We used deidentified patient encounter data, listed by first CC. The CC was used to identify the nature of the patient encounter as this data was available at the time of patient presentation, often dictates the patient’s ED workup, and would not have been affected by information discovered during the later stages of a

patient's hospital course. This approach is consistent with prior literature.^{9,13} To maintain anonymity, only the senior author, a member of the residency leadership team, had access to each resident's individualized study identification number.

We excluded from analysis encounters where no CC was listed or no resident was assigned. In cases where multiple residents were assigned to a single encounter (e.g., a patient had been signed out to a different resident), we analyzed this encounter only for the initial resident assigned. This was done as they are typically the most involved in the cognitive workload of determining the patient's initial diagnostic and treatment plan. The CC for each encounter was selected and entered into the EHR by the primary nurse who cared for the patient in the ED initially. At our institution, this is nearly always selected from a list of common CCs, although it can be entered as free text. Encounters in which multiple CCs were listed were only coded into a single domain based on the first listed CC.

Data Analysis

A list of common CCs in EM has been categorized into a set of 20 content domains via a consensus process by two EM attendings using the 2016 American Board of Emergency Medicine (ABEM) Model of Clinical Practice as a framework.¹⁴ For CCs identified in our data that were not already categorized by a previously described method,¹³ we repeated the same categorization process in which each CC was assigned to a single domain by two board-certified EM attending physicians at our institution. Disagreements between the two reviewers were adjudicated by a third board-certified emergency physician. If a symptom was entered as the CC, such as "fever" (which could correspond to one of multiple domains), it was preferentially categorized into a domain based on what the coding physicians felt was the

most likely to dictate the ED workup, rather than the "signs, symptoms, and presentations" domain. We used Excel (Microsoft Corp, Redmond, WA) to calculate descriptive statistics and create plots and tables. The top 10 most encountered domains overall were analyzed. We excluded less common domains given the low number of total encounters in each area, which would have been more vulnerable to random fluctuations in when these patients present to the ED.

This project was deemed exempt quality improvement by the University of Wisconsin Health Sciences Institutional Review Board.

RESULTS

A total of 315,614 encounters were initially identified from the EHR. Of these encounters 198 were excluded as no CC was listed. After excluding residents whose clinical experience was outside the study period and those who had left the training program prior to graduation or had a prolonged leave of absence, a total of 228,916 patient encounters from 69 residents were included in the analysis. Each resident was assigned to an average of 3,323 distinct patient encounters. Assessment of the top 10 most common clinical exposure domains showed wide ranges in the case numbers of individual residents. The [Table](#) lists the mean, minimum, maximum, interquartile range (IQR) and 25th and 75th percentile for the 10 most common content domains. The [Figure](#) shows the range of exposure to the 10 most common domains in box-and-whisker format.

DISCUSSION

Our data suggests that residents within a single training program have substantial variation in their clinical experiences as measured by the variation in ABEM content

Table. Mean, 25th–75th percentile ranges, interquartile range, and minimum/maximum encounters for the 10 most encountered domains per resident.

	Mean	Median	25 th , 75 th percentile	IQR	Minimum, maximum
Total encounters	3323		3086, 3609	523	2595, 4053
Abdominal and gastrointestinal disorders	583	594	528, 644	116	416, 721
Traumatic disorders	529	525	484, 570	86	370, 725
Cardiovascular disorders	327	330	302, 356	54	233, 429
Nervous system disorders	319	319	301, 340	39	226, 402
Musculoskeletal disorders (non-traumatic)	314	314	269, 361	92	179, 460
Thoracic-respiratory disorders	280	281	246, 313	67	178, 383
Systemic infectious disorders	165	169	149, 179	30	115, 219
Head, ear, eye, nose, and throat disorders	150	151	136, 165	29	96, 196
Signs, symptoms, and presentations	129	130	120, 142	22	88, 170
Psycho-behavioral disorders	126	128	106, 139	34	67, 211

IQR, interquartile range.




Figure. Top 10 most common clinical exposure domains seen by graduation per resident. Boxes illustrate the 25th–75th percentile of number of clinical exposures by residents in each domain, with whiskers representing the minima, maxima, and outliers.

domains seen by individual residents. This is similar to what was described by Langdorf et al. in 1990, despite the previous study being performed over three decades ago and the substantial subsequent differences in the utilization of the ED.⁸ We found wide interquartile ranges between the maximum and minimum number of encounters among residents, suggesting that some residents saw substantially more patients within particular domains than others.

The magnitude of the educational significance of the exposure variability of residents is unclear. It is possible that a resident who sees twice as many musculoskeletal chief complaints as another resident by graduation is significantly more competent in that domain. Alternatively, it is also possible that they have both attained the minimal level of exposure to competently manage musculoskeletal complaints independently. The effects of clinical exposure on clinical competence, including the minimal number of encounters required to demonstrate competency in a particular domain, is an open question and an avenue for further research. However, the formation of illness scripts is continually modified by subsequent patient encounters.^{3,4} Therefore, the identification of high degrees of variation among residents may prompt program leadership to institute changes in the curriculum or supplement clinical exposure with individualized learning plans. This is likely more important for domains that are encountered less frequently

overall, such as psycho-behavioral disorders, where larger relative differences in exposure could result in greater deficits in illness script formation.

In addition to prompting changes made by the program, identification of high variability in clinical exposure may enhance resident self-assessment. As demonstrated previously, self-assessment when done in isolation, is an imperfect means of driving improvement but can be enhanced greatly when informed by additional information from a variety of sources.¹⁵ Understanding the distribution of the patient encounters residents have during training, and the potential gaps in their clinical exposure, could be a potential means of allowing for informed self-assessment for a resident's clinical skills. This could be potentially further enhanced if facilitated under the supervision of faculty coaches within the program, a method that has become increasingly popular in medical education.^{16,17} Future work could follow a cohort of residents who are able to track their own patient volumes more regularly than was possible in the current study and compare themselves to their peers throughout training and evaluate whether any differences in clinical competence are identified. This could also allow programs to determine the perceived value of this information to residents. Finally, residents could use this data to drive their patient selection while working in the ED.

Beyond the potential for shaping resident self-assessments, clinical exposure data may have important implications for residency program leadership as we move toward an era of competency-based medical education (CBME). Two of the pillars of CBME, “teaching tailored to competencies” and “effective programmatic assessment,”¹⁸ lend themselves well to the identification of program clinical weaknesses as well as to the creation of new curricular experiences designed to address areas of limited clinical exposure identified by resident CC data. These experiences could potentially take the form of targeted readings or simulation sessions designed to supplement lower frequency clinical encounters.

LIMITATIONS

This was a single-center study in an urban, academic ED, and findings may not be generalizable to training programs in different environments. Additionally, the data was retrospective, making the educational utility of this information or any potential causes of variation difficult to determine.

Use of a CC to categorize each patient encounter into a clinical domain has an element of subjectivity and may have led to some encounters being miscategorized with respect to the workup done or final diagnosis. Some additional subjectivity may have been introduced by how we classified CCs that could potentially have been categorized into multiple different domains (such as “fever” or “ingestion”). This was done based on what was determined to be most likely to drive the initial workup in the department. For example, although a CC of “chest pain” could represent a cardiac or pulmonary etiology, in almost all cases, a cardiac etiology must be excluded. Therefore, it was felt that this would influence the formation and modification of the resident’s illness script most heavily. It is also possible that encounters were mischaracterized due to only using the first CC listed and not considering the others if multiple CCs were listed. Like the prior limitation, it was felt that the first CC was most likely to dictate the initial ED workup. Using discharge or final diagnoses instead was considered for this study, but it was felt that the CC is more likely to drive the initial differential and diagnostic workup for the patient.

Additionally, ABEM domains may be too broad to capture important differences in exposure (e.g., two residents with the same exposure to “respiratory disorders” could have seen large numbers of pneumonia patients or, alternatively, many patients with asthma). Training is inherently variable as the EM environment differs by clinical site, day, shift, or even season. Therefore, there may have been slight differences in when individual residents were in the ED clinically or the number/type of overall ED shifts worked. It is important to note that some of the included residents’ training occurred during the COVID-19 pandemic, which may have had an effect on both the variety and number of

clinical exposures seen by these residents. Future work could also explore exposure based on sub-domains from the ABEM model to get a more granular look at individual resident clinical experiences rather than relying on the relatively broad domains.

Other clinical variables may also have an effect on a resident’s clinical exposure, including the timing of months rotating in the ED. However, the ED did not undergo major changes in the staffing model of physicians (including residents) during this period. Also, while it is likely that more senior residents assign themselves to critically ill patients, this was felt to be unlikely to meaningfully impact our results given that data was obtained at the time of graduation. Therefore, each resident would have acted in a senior role for the same amount of time. Finally, our use of the EHR at the main clinical training site of the residency to generate the data did not capture the clinical experience at two other training sites for the residency that use a different EHR. This may have served to moderate or exacerbate the differences seen among residents. However, clinical experiences at these other sites comprised a total of only four months of the 36-month curriculum, and so it is likely that our overall findings would not have been substantially affected.

CONCLUSION

Within a single, three-year academic emergency medicine program, there was substantial variation among residents regarding the variety of patient chief complaints seen throughout residency when mapped to ABEM’s Model of Clinical Practice.

Address for Correspondence: Corlin M. Jewell, MD, University of Wisconsin School of Medicine and Public Health, BerbeeWalsh Department of Emergency Medicine, 800 University Bay Dr., Madison, WI 53705. Email: cmjewell@medicine.wisc.edu

Conflicts of Interest: By the *WestJEM* article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. No author has professional or financial relationships with any companies that are relevant to this study. There are no conflicts of interest or sources of funding to declare.

Copyright: © 2025 Jewell et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. Teunissen PW, Scheele F, Scherpbier AJ, et al. How residents learn: qualitative evidence for the pivotal role of clinical activities. *Med Educ.* 2007;41(8):763–70.
2. Bowen JL. Educational strategies to promote clinical diagnostic reasoning. *N Engl J Med.* 2006;355(21):2217–25.

3. Custers EJ, Regehr G, Norman GR. Mental representations of medical diagnostic knowledge: a review. *Acad Med*. 1996;71(10 Suppl):S55–61.
4. Hatala R, Norman GR, Brooks LR. Influence of a single example on subsequent electrocardiogram interpretation. *Teaching and Learning in Medicine*. 1999;11(2):110–7.
5. Stahmer S and Kuhn G. Optimizing resident training: results and recommendations of the 2009 Council of Residency Directors Consensus Conference. *Acad Emerg Med*. 2010;17 Suppl 2:S78–86.
6. Li J, Roosevelt G, McCabe K, et al. Pediatric case exposure during emergency medicine residency. *AEM Educ Train*. 2018;2(4):317–27.
7. Chen EH, Cho CS, Shofer FS, et al. Resident exposure to critical patients in a pediatric emergency department. *Pediatr Emerg Care*. 2007;23(11):774–8.
8. Langdorf MI, Strange G, Macneil P. Computerized tracking of emergency medicine resident clinical experience. *Ann Emerg Med*. 1990;19(7):764–73.
9. Kern MW, Jewell CM, Hekman DJ, et al. Number of patient encounters in emergency medicine residency does not correlate with in-training exam domain scores. *West J Emerg Med*. 2022;24(1):114–8.
10. Hayashino Y, Fukuhara S, Matsui K, et al. Quality of care associated with number of cases seen and self-reports of clinical competence for Japanese physicians-in-training in internal medicine. *BMC Med Educ*. 2006;6:33.
11. Strange GR and Chen EH. Use of emergency departments by elder patients: a five-year follow-up study. *Acad Emerg Med*. 1998;5(12):1157–62.
12. Cairns C and Kang K. National Hospital Ambulatory Medical Care Survey: 2021 emergency department summary tables. 2023. Available at: https://ftp.cdc.gov/pub/Health_Statistics/NCHS/Dataset_Documentation/NHAMCS/doc21-ed-508.pdf. Accessed September 21, 2024.
13. Bischof JJ, Emerson G, Mitzman J, et al. Does the emergency medicine in-training examination accurately reflect residents' clinical experiences? *AEM Educ Train*. 2019;3(4):317–22.
14. Counselman FL, Babu K, Edens MA, et al. The 2016 model of the clinical practice of emergency medicine. *J Emerg Med*. 2017;52(6):846–9.
15. Wolff M, Santen SA, Hopson LR, et al. What's the evidence: self-assessment implications for life-long learning in emergency medicine. *J Emerg Med*. 2017;53(1):116–20.
16. Deiorio NM, Moore M, Santen SA, et al. Coaching models, theories, and structures: an overview for teaching faculty in the emergency department and educators in the offices. *AEM Educ Train*. 2022;6(5):e10801.
17. Sargeant J, Lockyer J, Mann K, et al. Facilitated reflective performance feedback: developing an evidence- and theory-based model that builds relationship, explores reactions and content, and coaches for performance change (R2C2). *Acad Med*. 2015;90(12):1698–706.
18. Van Melle E, Frank JR, Holmboe ES, et al. A core components framework for evaluating implementation of competency-based medical education programs. *Acad Med*. 2019;94(7):1002–9.

The Effect of Hospital Boarding on Emergency Medicine Residency Productivity

Peter Moffett, MD
Al Best, PhD
Nathan Lewis, MD
Stephen Miller, DO
Grace Hickam, MD
Hannah Kissel-Smith, MD
Laura Barrera, MD
Scott Huang, MD
Joel Moll, MD

Department of Emergency Medicine, Virginia Commonwealth University School of Medicine, Richmond, Virginia

Section Editors: Abra Fant, MD and William Edward Soares, MD

Submission history: Submitted July 21, 2024; Revision received October 7, 2024; Accepted October 11, 2024

Electronically published November 27, 2024

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: [10.5811/westjem.31064](https://doi.org/10.5811/westjem.31064)

Introduction: Emergency department boarding has escalated to a crisis, impacting patient care, hospital finances, and physician burnout, and contributing to error. No prior studies have examined the effects of boarding hours on resident productivity. If boarding reduces productivity, it may have negative educational impacts. We investigated the effect of boarding on resident productivity as measured by patients per hour and hypothesized that increased boarding leads to decreased productivity.

Methods: This was a retrospective study at a quaternary, urban, academic Level I trauma center from 2017–2021 with a three-year emergency medicine residency of 10–12 residents per year and annual volumes of 80,000–101,000. Boarding was defined as the time between an admission order and the patient leaving the ED. We created a multivariable mixed model with fixed covariates for year, month, day of week, resident experience, shift duration, total daily ED patients, and with residents as repeated measures. The effect of boarding was estimated after covarying out all other factors.

Results: All variables included in the model were significantly associated with changes in productivity. Resident experience has the largest effect such that for each month of residency experience, a resident adds 0.012 patients per hour (95% confidence interval [CI] 0.010–0.014). Isolating the effect of boarding demonstrated that for every additional 100 hours of boarding, a resident's productivity decreased by 0.022 patients per hour (95% CI 0.016–0.028). In the study, the median daily boarding was 261 hours; if this were eliminated (assuming a resident completes 100 10-hour shifts annually), a resident could be expected to see 56.9 more patients per year (95% CI 40.7–73.1).

Conclusion: Hospital boarding significantly reduces resident productivity as measured by patients per hour. Further studies are warranted to determine the educational impact. [West J Emerg Med. 2025;26(1.2)54–62.]

INTRODUCTION

Emergency department (ED) boarding (defined as patients admitted to the hospital but remaining in the ED) has reached critical levels and has been declared a crisis by

the American College of Emergency Physicians.¹ The scope of the crisis is daunting with effects on patient care, errors, physician burnout, hospital economic stress, and ambulance diversion.² Increased ED boarding also leads to increases in

medication errors, time to antibiotics, time to percutaneous coronary intervention for patients with myocardial infarction, time to care for patients with acute stroke, patient mortality, and risk-adjusted hospital spending, and has effects on all levels of acuity.^{3–10}

Within the context of boarding, EDs must also provide sound educational training involving both quality and quantity of patient experiences. Residency programs seek to improve efficiency and productivity in their residents throughout their training. Many variables have been associated with resident productivity including time of shift, shift length, and resident experience.^{11–13} There are, however, few studies that evaluate the effect of ED crowding and boarding time on the effect of emergency medicine (EM) resident productivity.¹⁴ If boarding decreases the number of patients seen during a residency, there may be an impact on resident education.

In this study we aimed to investigate the effect of boarding on EM resident productivity as measured by patients per hour. We hypothesized that increased hospital boarding would result in decreased resident productivity.

METHODS

Study Design and Setting

This was a retrospective study conducted at the Virginia Commonwealth University Health System, the only comprehensive Level I trauma center in Richmond, VA. During the study period from January 2017–June 2021, the total patient volumes ranged from 80,000–101,000 per year. On average, 30% of patients were admitted to the hospital, of whom 5% went to the intensive care unit. Patients <18 years of age constituted 22% of the total volume. The department is staffed with board-certified emergency physicians, and during the study period 81% of patients were seen by a resident. The remaining non-resident cases were seen by advanced practice practitioners (APP) in a low-acuity area of the ED or by attending physicians and were not included in the study. Throughout the study there was no change in this staffing model such that APPs were never competing for the same patients as residents. The department has 76 beds with 35 in an acute area, 10 in trauma/resuscitation, 10 in a mid-track area, 16 in a pediatric department, and five in a fast-track zone.

Our residency program is three years in length, and class sizes ranged from 10 residents in 2017 to 12 residents in 2021. During postgraduate years (PGY)-1, 2, and 3, residents work in the ED for 26 weeks, 29 weeks, and 35 weeks, respectively. Resident shift lengths varied from 9–12 hours with the most typical shift being 10 hours. On average, each 24-hour period had a total of 137 hours of resident coverage in overlapping shifts. The EM residents saw patients in all Emergency Severity Index (ESI) categories and were the primary physicians for all emergent patients (ESI 1 and 2). Residents cared for patients in all areas of the ED other than the

Population Health Research Capsule

What do we already know about this issue?
Emergency department boarding negatively impacts patient care, hospital efficiency, and physician well-being.

What was the research question?
Does increased ED boarding reduce emergency medicine resident productivity, as measured by patients per hour?

What was the major finding of the study?
For every additional 100 hours of ED boarding, a resident's productivity decreased by 0.022 patients per hour (95% CI 0.016–0.028); a resident sees 57 fewer patients per year due to boarding.

How does this improve population health?
Understanding the negative effects of boarding on productivity may help policy makers find solutions to improve patient flow, patient care, resident education, and overall health outcomes.

low-acuity area. All residents staff patients directly with an attending physician without oversight by a more senior resident; therefore, the productivity numbers for residents in all three years of training are independent.

The study was granted exempt status by the Virginia Commonwealth University Institutional Review Board (HM20024717).

Selection of Participants

Data from all patients evaluated by an EM resident was captured in a database, and in conjunction with scheduling data it was used to determine the average number of patients per hour. Only EM residents were included. The study period was selected as this was the maximum amount of time for which data was available prior to the hospital switching to a new electronic health record. As the database was initially created to provide feedback to residents, certain data was removed and not available to us for analysis. Information from the first month of EM for each resident was not provided, and due to initial effects from the COVID-19 pandemic, data from April–July 2020 was not included.

Measurements

We combined three databases for analysis: the patient database of all ED encounters; the resident scheduling database; and the hospital boarding database.

During the study period, the EM residency program received monthly, system-generated reports listing the unique patient identifier, name of the resident assigned to care for the patient, the ESI acuity level, the date/time of first contact and check out, and the disposition. The resident assignment was derived from tracking board data, and in scenarios where multiple residents were assigned to a patient encounter, only the first resident assigned was credited for each unique patient encounter. The EM residents were scheduled for 9-hour, 10-hour, or 12-hour shifts during the study period. All non-EM residents and staff were excluded from the patient database.

Boarding data was reported daily from hospital analytics. The number of hours of boarding was defined as the time between an admission order and when the patient left the ED. Boarding hours was selected as this was the variable available to us from the hospital analytics database.

Outcomes

We designed a model to isolate the effects of ED boarding on resident productivity as measured by patients per hour. Patients per hour was defined as the total number of new patients seen during the shift divided by the duration of the shift in hours. The covariates were chosen from those found in previous studies to be related to resident productivity.^{11,13,15,16} These included year, month, day of the week, cumulative residency months in training, shift duration, total patients per day, and boarding. Months in training was chosen as a continuous covariate to delineate resident experience rather than the rough classification of PGY-1, -2, or -3 based on the observation that resident productivity begins low in the PGY-1 year, increases in the PGY-2 year, and then plateaus. This monthly experience variable was modeled using cubic regression.

Analysis

We described the data using counts and percentages. Patients per hour was modeled using a multivariable mixed model, with covariates defined as fixed effects and residents as repeated measures. We used an autoregressive (AR1) covariance structure to account for the dependence between repeated measures. The fixed effects were year (reference = 2019), month (reference = 12), day of the week (reference = Thursday), resident month in training (centered on 18), total patients per day/100, shift duration, and daily boarding hours/100. We chose the year 2019 as a reference as it was the last full year of data prior to the start of the COVID-19 pandemic. December was chosen as it aligns with the 18th month of residency, which is when productivity plateaued in our model. Thursday was selected as it is thought to represent the day with the most ideal flow since it avoids weekends, Monday, and Friday patient surges, as well as Wednesday morning didactics when EM residents are not working clinically. The total patients per day, shift duration, and

boarding hours were referenced at the median values in our dataset.

We estimated the effect of boarding from the marginal regression model after covarying out all other factors. Estimates are described using 95% confidence intervals. All data management and analysis were performed using SAS software (version 9.4 and JMP Pro version 17.2 (SAS Institute Inc, Cary, NC)).

RESULTS

Characteristics of Study Subjects

During the study period, 263,058 patients were seen in the ED by 601 clinicians including the 80 EM residents studied. During the 49 months studied between 2017–2021, EM residents were scheduled to 16,949 shifts and were assigned 188,685 patients (Table 1). Total daily patient volume varied considerably during this time (mean 177, SD 26, range

Table 1. Characteristics of the emergency department residents' shifts and patients evaluated (January 2017–June 2021).

Characteristic	Shifts N	Patients N	(%)
Total	16,949	188,685	
Year			
2017	3,496	44,119	(23)
2018	3,955	47,569	(25)
2019 (11 months)*	4,053	47,035	(25)
2020 (8 months)†	3,101	29,191	(15)
2021 (6 months)	2,344	20,771	(11)
Month			
1- January	1,909	21,052	(11)
2- February	1,576	18,004	(10)
3- March	1,680	18,901	(10)
4 [†] - April	1,302	15,229	(8)
5 [†] - May	1,371	16,385	(9)
6 [†] - June	1,337	15,191	(8)
7 [†] - July	820	10,129	(5)
8- August	1,560	15,543	(8)
9- September	1,376	14,741	(8)
10- October	1,431	15,299	(8)
11*- November	1,062	11,639	(6)
12- December	1,525	16,572	(9)
Day of week			
Sunday	2,249	25,887	(14)
Monday	2,679	29,099	(15)
Tuesday	2,756	29,504	(16)
Wednesday [‡]	1,989	21,970	(12)
Thursday	2,601	27,874	(15)

(Continued on next page)

Table 1. Continued.

Characteristic	Shifts N	Patients N	(%)
Friday	2,525	28,785	(15)
Saturday	2,150	25,566	(14)
Shift			
7 AM to 5 PM	1,688	16,332	(9)
7 AM to 7 PM	180	2,512	(1)
9 AM to 7 PM	2,546	28,306	(15)
12 PM to 10 PM	3,386	38,586	(20)
2 PM to 12 AM	2,470	28,631	(15)
3 PM to 12 AM	3,553	41,138	(22)
9 PM to 7 AM	3,126	33,180	(18)
PGY			
PGY-1 [§]	5,162	44,817	(24)
PGY-2	4,756	57,447	(30)
PGY-3	7,031	86,421	(46)
Disposition			
Admitted		74,663	(40)
Discharged		114,022	(60)

*November 2019 was excluded as the hospital information management system was down.

†April 2020 through July 2020 was excluded due to COVID-19 and hospital changes.

‡Wednesdays mornings are resident didactics.


§The first month of a residency was excluded (orientation month).
ESI, Emergency Severity Index; PGY, postgraduate year.

88–263). As indicated in the table, the ED experienced a patient count variability that changed across years, months, days of the week, shifts, and PGY level. Of all 188,167 patients seen by EM residents, 40% were admitted.

Boarding hours per day varied considerably (mean 281, SD 127, range 50.8–914.4; [Figure 1](#)). The hospital information system calculated boarding hours daily; however, across the 1,490 days studied, there were six impossible (negative) values and nine very low values. Low values were identified by large residuals in the multiple regression model. Rather than treating these as missing values, we used a multiple regression model to impute the 15 values in question.

Main Results

All the factors in the repeated-measures mixed-model were significant ($P < 0.001$). [Table 2](#) shows the estimated effect of each term in the model. The joint effect of all the factors on resident productivity is shown in [Figure 2](#). These profile plots show the marginal model predicted value of resident productivity on the vertical axis across all the covariates on the separate horizontal axes. The importance

**Figure 1.** Boarding across study years.

Line set at median boarding hours across the entire study period (261 hours/day).

Each box plot represents a month (line = median, box = 25th to 75th quartile, whiskers = typical extremes, circles = outliers).

Note: April 2020–July 2020 hours are not available as they correspond to the beginning of the COVID-19 pandemic.

of a factor is visualized by the steepness of the prediction trace.

Isolating the effect of boarding demonstrated that for every additional 100 hours of daily departmental boarding, individual resident productivity decreased by 0.022 patients per hour (95% confidence interval [CI] 0.016–0.028, [Table 2](#)). In the reference standard scenario, a resident could be expected to see 1.10 patients per hour with boarding at the daily median (261 hours) but could see 1.15 patients per hour if boarding were eliminated ([Figure 2](#), Panel C). [Table 3](#) shows how resident productivity was degraded by boarding across the range of values seen at our institution. A resident would see 1.14 patients per hour when boarding was at the lowest in the study compared to 0.95 patients per hour at the maximum level of boarding seen in the study, which is a difference of 0.19 patients per hour (95% CI 0.15–0.22). Assuming a resident completes approximately 100 shifts a year that are of 10 hours duration and boarding was eliminated, then a resident could be expected to see 56.9 more patients per year (95% CI 40.7–73.1). This would represent a 5% increase in patient volume per resident annually.

Resident experience has the largest effect on resident productivity. Resident productivity was low initially at 0.5 patients per hour (95% CI 0.46–0.54) by the second month of training ([Figure 2](#)). Improvement was initially rapid to 0.75 patients per hour at seven months, then plateaued near the 18-month point (1.10 patients per hour) to finally reach 1.12 patients per hour at the end of the 36 months (95% CI 1.08–1.17). When evaluating our data by PGY level, our

Table 2. Multiple regression results predicting new patients per hour per resident for each variable.

Effect	Estimated new patients per hour	Standard error	95% CI
Intercept	1.0957	0.0173	1.0618 to 1.1297
Year			
2017	0.1501	0.0122	0.1262 to 0.1740
2018	0.0837	0.0117	0.0608 to 0.1065
2019	[reference]		
2020	-0.0641	0.0137	-0.0909 to -0.0373
2021	-0.1682	0.0156	-0.1987 to -0.1377
Month			
1- January	0.0635	0.0172	0.0298 to 0.0972
2- February	0.0776	0.0182	0.0420 to 0.1133
3- March	0.0498	0.0181	0.0144 to 0.0852
4- April	0.0840	0.0197	0.0453 to 0.1227
5- May	0.0750	0.0196	0.0366 to 0.1133
6- June	0.0585	0.0201	0.0191 to 0.0979
7- July	-0.0077	0.0219	-0.0507 to 0.0353
8- August	0.0550	0.0185	0.0188 to 0.0912
9- September	0.0654	0.0187	0.0288 to 0.1021
10- October	0.0487	0.0184	0.0127 to 0.0847
11- November	0.0486	0.0199	0.0095 to 0.0876
12- December	[reference]		
Day of the week			
Sunday	0.0587	0.0118	0.0357 to 0.0818
Monday	-0.0312	0.0118	-0.0542 to -0.0082
Tuesday	0.0122	0.0110	0.0094 to 0.0338
Wednesday	0.1094	0.0123	0.0854 to 0.1334
Thursday	[reference]		
Friday	0.0475	0.0109	0.0261 to 0.0688
Saturday	0.1182	0.0120	0.0948 to 0.1417
Resident months (linear)*	0.0122	0.0010	0.0101 to 0.0142
(quadratic)	-0.0011	0.0000	-0.0012 to -0.0010
(cubic)	0.00003	0.00001	0.00002 to 0.00004
Total patients per day (per 100 patients)*	0.4021	0.0165	0.3697 to 0.4344
Shift duration*	-0.1277	0.0070	-0.1413 to -0.1140
Boarded (per 100 hours)*	-0.0218	0.0032	-0.0280 to -0.0156

The mixed-model also included resident as a repeated-effect with an AR(1) covariance structure.

*Continuous covariates were referenced to the median value. Median resident month = 18, total patients per day/100 = 1.77, shift duration = 10 hours, boarded hours/100 = 2.61.

CI, confidence interval.

PGY-1 residents saw 0.75 per hour, PGY-2 residents saw 1.10 patients per hour, and PGY-3 residents saw 1.12 patients per hour.

Total patients per day presenting to the ED was the next most important factor in resident productivity. For every 100 new patients presenting to the ED, an individual resident

would be expected to add 0.40 patients per hour (95% CI 0.37–0.43). The median value for daily total patient volume was 177 patients per day, but a low-volume day at the 10th percentile (143 total patients) resulted in a corresponding decrease in resident productivity to 0.96 patients per hour (95% CI 0.92–1.00). For a high-volume day at the 90th




Figure 2. Multiple regression results predicting new patients per hour per resident for each variable. All values (year, month, day of week, EM resident months, total patients, shift duration) in model held at reference standards with adjustments to boarding (last panel of each graph). Expected patients per hour in each scenario is indicated by the red number in the Y axis with 95% confidence intervals in blue. As boarded hours change (last panel of each graph) so do patients per hour (red number to left of each graph) in each of the three scenarios (A: Median boarding of 261 hours. B: Reducing boarding by 100 hours. C: Eliminating boarding hours.)

percentile (210 patients), resident productivity increased to 1.23 patients per hour (95% CI 1.19–1.26).

Resident productivity also changed based on the year, shift duration, and day of the week. Resident

productivity was highest in 2017 at 1.25 patients per hour (95% CI 1.21–1.28) and steadily decreased to the 0.93 patients per hour seen in 2021. Resident productivity for a nine-hour shift was predicted to be 1.21 patients per hour

Table 3. Estimated resident productivity by boarding hours.

Cutoff	Boarded (hours)	Estimated patients per hour	Standard error	95% CI
Maximum	914	0.954	0.027	0.900 to 1.007
75th percentile	351	1.076	0.018	1.042 to 1.111
Median	261	1.096	0.017	1.062 to 1.130
25th percentile	189	1.111	0.017	1.077 to 1.146
Minimum	51	1.141	0.018	1.105 to 1.178
No boarding	0	1.153	0.019	1.115 to 1.190

Marginal estimates from the mixed model with the following factors held constant: year = 2019, month = 12, day of the week = 5 (Thursday), resident month in training = 18, total patients per day/100 = 1.77, shift duration = 10 hours.

CI, confidence interval.

(95% CI 1.19–1.26), whereas for a 12-hour shift it was predicted to be 0.84 patients per hour (95% CI 0.80–0.89). Saturdays and Wednesdays averaged approximately 1.21 patients per hour, Sundays, and Fridays approximately 1.15 per hour, and Mondays, Tuesdays, and Thursdays 1.10 patients per hour.

Month-to-month variability had the smallest effect on resident productivity. Compared with the other months, July and December had lower resident productivity (1.09 vs 1.16 patients per hour).

DISCUSSION

To our knowledge, this is the first study to demonstrate that there is a significant reduction in resident productivity (measured as patients per hour) due to hospital boarding in the ED. In our model, this resulted in a decrease of 0.022 patients per hour (95% CI 0.016–0.028) for every 100 hours of daily boarding. While performed at a single institution, our dataset broadly aligns with multiple studies previously completed regarding resident productivity. In our study, we analyzed resident experience as the number of months in training rather than divided into PGY level. This was based on our observation that productivity rapidly increased during the PGY-1 year and then plateaued in the middle of the PGY-2 year.

When evaluating our data by PGY level, our PGY-1 residents saw 0.75 patients per hour, PGY-2 residents saw 1.10 patients per hour, and PGY-3 residents saw 1.12 patients per hour. Prior studies have demonstrated similar patterns with PGY-1 to -3 residents seeing between 0.79–0.81 patients per hour, 1.05–1.2 patients per hour, and 1.22–1.27 patients per hour, respectively.^{17–19} A study by Henning et al showed rapid progression from PGY-1 to PGY-2 year and then gradual progression in PGY-3 year but was based on patients per day.²⁰ Similarly, a study by Turner-Lawrence and Todd saw increasing productivity from 1.2 patients per hour to 1.5 patients per hour to 1.6 patients per hour by PGY-1 to -3 residents, respectively.¹³ While these productivity numbers are higher than those in our study, the authors did not adjust for additional variables.

In a more comparable study, Kirby et al reported the efficiency of EM residents during ED crowding.¹⁴ The authors used the National Emergency Department Overcrowding Study (NEDOCS) scoring system to categorize states in the ED as not crowded, crowded, and overcrowded. They found that resident productivity measured as new patients per hour increased initially in all year groups as the ED transitioned from not crowded to crowded, but then remained stable when transitioning from crowded to overcrowded. While the NEDOCS score uses a measure of ED boarding (the waiting time of the longest admitted patient), it does not include total patient boarding hours as in our study. Our study more directly examines the effect of boarding (one element of crowding) on resident

productivity. The paradoxical increase in resident productivity in the Kirby study may have been due to an increased number of patients presenting to the ED, which could have increased the NEDOCS score. Our study demonstrated that resident productivity increased with higher patient volumes, and including this in our model allowed us to better isolate the effect of boarding.

According to a study by the Academy of Administrators in Academic Emergency Medicine and the Association of Academic Chairs of Emergency annual benchmark survey, boarding times have dramatically increased since the COVID-19 pandemic.²¹ By the end of their study period, the median number of boarding hours per month was 11,480, which approximates to 382 hours of daily boarding. In our study, which includes a pre-pandemic period, the median daily boarding was 261 hours, suggesting that boarding is likely worsening over time and is a problem at many academic medical centers.

The educational impact of decreased patient volumes caused by boarding is uncertain. It is reasonable to expect that residents seeing fewer cases may lose valuable learning opportunities, but this has not been well studied and no firm numbers exist to suggest a threshold at which education suffers. Prior authors have surveyed residents regarding a perceived decrease in education during crowding.^{22,23} These studies concluded that residents did not perceive a difference in education during these times, but they used differing measures of crowding, were survey-based, and underpowered. Educators may switch to different models of teaching during periods of high boarding, leading to residents perceiving a less deleterious effect.²⁴

Others have postulated an educational Starling effect whereby some boarding allows supervising physicians more time to teach, but at some point there are diminished returns as fewer new patients become available to discuss.²⁵ A more recent study was conducted during the current boarding epidemic; the authors surveyed EM program directors regarding their perceptions of the impact of boarding on resident training.²⁶ In this study, 80% of the respondents felt that boarding negatively affected resident education, especially in the domains of managing department throughput and managing high volumes of patients per resident. While survey-based in nature, the study results broadly aligns with the prior studies in this area.

Theoretically, residents who see fewer cases may lose valuable learning opportunities. While the components of Bloom's domains of educational activities can be learned via different modalities of instructional techniques, clinical experience allows for the linking of knowledge to skills and then to attitudes/emotions.²⁷ By decreasing a learner's exposure to patients, one could argue that residents may lose valuable experiential learning opportunities. While some of these can be replicated in simulation or case-based discussion, other skills cannot and are best learned via hands-

on, experiential learning encounters. Experiential learning theory, as described by Kolb, highlights the importance of real-life experience and the influence this has on learning.²⁸ Unlike traditional learning and instructional methodology that focuses on rote memorization, experiential learning is an active process where residents are engaged in concept transformation through action as well as reflection on their experiences and patient encounters.

This learning theory also emphasizes principles of adult education in which prior learning experiences can be leveraged to create more meaningful and relevant educational experiences.²⁹ Additionally, decreasing patient interaction may also affect residents' application and translation of knowledge into practice. Behavioral learning theory emphasizes learning through interactions with the environment where reinforcement and feedback can encourage modification of behaviors. By incorporating behavioral learning strategies, medical education can foster not only technical competencies but also the development of professional habits such as effective communication between team members and patients.³⁰

LIMITATIONS

This study has several limitations. This was a single-center study that took place in a high acuity, quaternary-care center that also experiences high levels of boarding, which may limit generalizability to other centers. The database that captured the resident patient assignment was based on tracking board data and may have occasionally miscredited a resident with a patient encounter; however, as the dataset was large and involved multiple years with complete datasets for three full classes of residents this is unlikely to have greatly influenced the data. Our resident class size did increase during the 2021 year and thus could theoretically have decreased the number of patients available per resident. While we did not study that directly, it is unlikely to have impacted the data greatly as the additional residents allowed for the creation of an outside rotation at a free-standing emergency center and, therefore, resident staffing hours stayed generally consistent at the study site.

Our model did not include a measure of patient acuity as a covariate. While the ESI category and disposition were recorded for each patient, we did not feel there was a reliable way to convert this data into a meaningful measure of hourly acuity that influenced the amount of time a resident might dedicate toward patient care. For example, an ESI-1 patient who is admitted for an ST-segment elevation myocardial infarction may stay in the department for 15 minutes leaving the bed open for a new patient, while an ESI-3 patient requiring a workup for abdominal pain including imaging who is discharged may occupy a room and a resident for multiple hours. Since our dataset was large, it was assumed that all residents would be exposed equally to the same mix of acuities on individual shifts, by the end of their residency

and thus limit the effect on the data. Additionally, recent studies have called into question the accuracy of the ESI.^{26,27} A prior study on resident productivity did not show a correlation between ESI and clinician disposition times.¹⁴

Our study also included data from the COVID-19 pandemic, which affected patient volumes and ED boarding. The dataset we used was initially meant for reporting individual residents' productivity measures, so data from the first few months of the pandemic was not available for our current study. This likely served to decrease the effect of the initial pandemic response on our data. Just prior to the pandemic our ED had seen a growth in patient volumes from 87,000 patients per year to a peak of 101,000 patients per year, which was followed by a rapid decline to 83,000 a year in the 2021–2022 year. The volumes did slowly rise after the study period. This may have influenced some of the data from our later resident-year groups and served to decrease productivity.

Our measure of boarding may also have limitations. Total boarding hours per day was the variable available from our hospital analytics department. The number of boarded patients per day may have provided different data. For example, in our model a single behavioral health patient boarding for 20 hours from one day would be indistinguishable from 20 patients boarding in 20 individual rooms for a single hour each. As the dataset is large, and all residents were exposed to the same conditions throughout their time, it is unlikely any one resident's data (or the trend) would be affected based on these types of outliers.

CONCLUSION

We found a significant reduction in resident productivity as measured by patients per hour during periods of increased boarding. Further studies are warranted to determine the educational impact of these findings.

Address for Correspondence: Peter Moffett, MD, Virginia Commonwealth University, Department of Emergency Medicine, 1250 E Marshall Street, Suite 600, Richmond VA 23298. Email: peter.moffett@vcuhealth.org

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. No author has professional or financial relationships with any companies that are relevant to this study. There are no conflicts of interest or sources of funding to declare.

Copyright: © 2025 Moffett et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. American College of Emergency Physicians. Emergency department boarding and crowding. Available at: <https://www.acep.org/administration/crowding-boarding/>. Accessed December 29, 2022.
2. Kelen GD, Wolfe R, D'Onofrio G, et al. Emergency department crowding: the canary in the health care system. *NEJM Catalyst Innovations in Care Delivery*. 2021. Available at: <https://catalyst.nejm.org/doi/full/10.1056/CAT.21.0217>. Accessed April 4, 2022.
3. Kulstad EB, Sikka R, Sweis RT, et al. ED overcrowding is associated with an increased frequency of medication errors. *Am J Emerg Med*. 2010;28(3):304–9.
4. Chatterjee P, Cucchiara BL, Lazarciuc N, et al. Emergency department crowding and time to care in patients with acute stroke. *Stroke*. 2011;42(4):1074–80.
5. Kulstad EB and Kelley KM. Overcrowding is associated with delays in percutaneous coronary intervention for acute myocardial infarction. *Int J Emerg Med*. 2009;2(3):149–54.
6. Fee C, Weber EJ, Maak CA, et al. Effect of emergency department crowding on time to antibiotics in patients admitted with community-acquired pneumonia. *Ann Emerg Med*. 2007;50(5):501–9, 509.e1.
7. Baloescu C, Kinsman J, Ravi S, et al. The cost of waiting: association of ED boarding with hospitalization costs. *Am J Emerg Med*. 2021;40:169–72.
8. Richardson DB. Increase in patient mortality at 10 days associated with emergency department overcrowding. *Med J Austral*. 2006;184(5):213–6.
9. McCarthy ML, Zeger SL, Ding R, et al. Crowding delays treatment and lengthens emergency department length of stay, even among high-acuity patients. *Ann Emerg Med*. 2009;54(4):492–503.e4.
10. White BA, Biddinger PD, Chang Y, et al. Boarding inpatients in the emergency department increases discharged patient length of stay. *J Emerg Med*. 2013;44(1):230–5.
11. Joseph JW, Henning DJ, Strouse CS, et al. Modeling hourly resident productivity in the emergency department. *Ann Emerg Med*. 2017;70(2):185–90.e6.
12. Jeanmonod R, Jeanmonod D, Ngiam R. Resident productivity: does shift length matter? *Am J Emerg Med*. 2008;26(7):789–91.
13. Turner-Lawrence D and Todd BR. Monthly progression of emergency medicine resident efficiency: what can we expect of our residents throughout training? *J Emerg Med*. 2019;57(1):77–81.
14. Kirby R, Robinson RD, Dib S, et al. Emergency medicine resident efficiency and emergency department crowding. *AEM Educ Train*. 2019;3(3):209–17.
15. Jeanmonod R, Brook C, Winther M, et al. Resident productivity as a function of emergency department volume, shift time of day, and cumulative time in the emergency department. *Am J Emerg Med*. 2009;27(3):313–9.
16. Joseph JW, Davis S, Wilker EH, et al. Modelling attending physician productivity in the emergency department: a multicentre study. *Emerg Med J*. 2018;35(5):317–22.
17. DeBehnke D, O'Brien S, Leschke R. Emergency medicine resident work productivity in an academic emergency department. *Acad Emerg Med*. 2000;7(1):90–2.
18. Fredette J, Kim T, McHugh D, et al. A descriptive analysis of emergency medicine resident productivity over the course of training. *AEM Education and Training*. 2021;5(S1):S44–8.
19. Jeanmonod R, Damewood S, Brook C. Resident productivity: trends over consecutive shifts. *Int J Emerg Med*. 2009;2(2):107–10.
20. Henning DJ, McGillicuddy DC, Sanchez LD. Evaluating the effect of emergency residency training on productivity in the emergency department. *J Emerg Med*. 2013;45(3):414–8.
21. Kilaru AS, Scheulen JJ, Harbertson CA, et al. Boarding in US academic emergency departments during the COVID-19 pandemic. *Ann Emerg Med*. 2023;82(3):247–54.
22. Mahler SA, McCartney JR, Swoboda TK, et al. The impact of emergency department overcrowding on resident education. *J Emerg Med*. 2012;42(1):69–73.
23. Pines JM, Prabhu A, McCusker CM, et al. The effect of ED crowding on education. *Am J Emerg Med*. 2010;28(2):217–20.
24. Atzema C, Bandiera G, Schull MJ, et al. Emergency department crowding: the effect on resident education. *Ann Emerg Med*. 2005;45(3):276–81.
25. Shayne P, Lin M, Ufberg JW, et al. The effect of emergency department crowding on education: blessing or curse? *Acad Emerg Med*. 2009;16(1):76–82.
26. Goldflam K, Bradby C, Coughlin RF, et al. Is boarding compromising our residents' education? A national survey of emergency medicine program directors. *AEM Educ Train*. 2024;8(2):e10973.
27. Bloom BS, Engelhart MD, Furst EJ, et al. *Taxonomy of Educational Objectives: Cognitive and Affective Domains*. New York, New York: David McKay Company, Inc, 1956.
28. Kolb DA. *Experiential Learning: Experience as the Source of Learning and Development*. Englewood Cliffs, NJ: Prentice-Hall; 1984.
29. Schultz K, McEwen L, Griffiths J. Applying Kolb's learning cycle to competency-based residency education. *Acad Med*. 2014;91(2):284.
30. Merriam S. *Adult Learning: Linking Theory and Practice*. San Francisco, CA: Jossey-Bass; 2013.
31. Sax DR, Warton EM, Mark DG, et al. Evaluation of the emergency severity index in US emergency departments for the rate of mistriage. *JAMA Netw Open*. 2023;6(3):e233404.
32. Mistry B, Stewart De Ramirez S, Kelen G, et al. Accuracy and reliability of emergency department triage using the emergency severity index: an international multicenter assessment. *Ann Emerg Med*. 2018;71(5):581–7.e3.

Palliative Care Boot Camp Offers Skill Building for Emergency Medicine Residents

Julie Cooper, MD*
Jenna Fredette, MD†

*University of Pennsylvania, Department of Emergency Medicine, Philadelphia, Pennsylvania
†ChristianaCare, Department of Emergency Medicine, Newark, Delaware

Section Editors: Paul Logan Weygandt, MD and Benjamin Holden Schnapp, MD

Submission history: Submitted June 22, 2023; Revision received March 3, 2024; Accepted May 30, 2024

Electronically published September 6, 2024

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.18381

BACKGROUND

Emergency medicine (EM) residents routinely care for critically ill patients in both the emergency department (ED) and intensive care units.¹ Proficiency in primary palliative care skills is essential for all emergency clinicians.^{2,3} However, a significant number of residents lack exposure to formal education and training in palliative care.^{4,5} Moreover, education and training in palliative care encompasses several Accreditation Council for Graduate Medical Education (ACGME) competencies including system navigation for patient-centered care, understanding the physician's role in the healthcare system, patient- and family-centered communication, and interprofessional and team communication.⁶

Current curricula addressing primary palliative care skills in EM are notably limited.⁷⁻¹² Historically, our residency experienced inconsistencies in the teaching of primary palliative care skills. They were sporadically covered during regular conferences or left to develop organically over time. Furthermore, postgraduate year-2 (PGY-2) residents, who primarily manage seriously ill patients, found themselves engaging in challenging serious-illness conversations with patients and families with little to no training. Recognizing the imperative for more comprehensive education, we introduced a four-week, intensive primary palliative care curriculum specifically tailored for EM PGY-2 residents that was entitled "Palliative Care Bootcamp."

OBJECTIVES

The overall objective of the bootcamp was to introduce and strengthen primary palliative care skills among PGY-2 residents at an independent academic medical center. At the end of the curriculum, residents would be able to 1) define the scope of hospice and palliative medicine; 2) understand what primary palliative care skills are for non-specialty trained physicians; 3) recognize ED patients with palliative care needs; 4) implement a hospice evaluation; 5) understand how interdisciplinary teams are involved in the care of seriously ill patients; and 6) build communication skills for discussing goals of care (GOC).

CURRICULAR DESIGN

The curriculum and assessment were exempt from the institutional review board. Using Kern's six-step approach to curriculum development, we created an introductory primary palliative care curriculum. An EM faculty member with an interest in palliative care and residency leadership collaborated to develop the curriculum. The residency program endorsed the curriculum as it aligned with a curriculum redesign to include more PGY-specific education.

The curriculum was initially developed in 2017. The interdisciplinary palliative care team at the study institution served as content experts. The team performed a broad review of the residency curriculum and prioritized high-yield topics tailored to the local context. Sessions were scheduled during weekly conference and spanned four consecutive weeks. This schedule allowed for an intensive experience and allowed for rapid skill development. The curriculum is strategically delivered early in the PGY-2 year to leverage residents' existing experience in caring for seriously ill patients and facilitate meaningful reflection and inquiry.

The curriculum is structured in two phases (Table 1). The first phase spans three weeks and consists of three two-hour sessions. These sessions are dedicated to primary palliative care fundamentals such as an introduction to palliative care, prognosis and trajectory, and non-pain symptom management. Session facilitators included the EM faculty content expert as well as members of the institutional palliative care team, the director of chaplaincy who specialized in family support, the director of palliative care, and the palliative care fellow. Each session encompassed a didactic segment, interactive case-based discussions using scenarios prepared by facilitators or contributed by residents, and opportunities for resident questions.

In the final week, residents engaged in a four-hour session in the simulation center. This session was led by the EM content expert who is a trained facilitator with Vital Talk, a national non-profit that promotes evidence-based education in serious-illness communication.¹³ This session involves using a standardized patient. Residents are assigned to a

Table 1. The breakdown of palliative care bootcamp sessions by hour detailing the topic, learning objectives, mapping to ACGME* competencies, and the format of the session.

Hour	Topic	Objectives	ACGME competencies	Format/facilitator
1	Intro to primary palliative care in emergency medicine	Define primary palliative care and identify common ED presentations of patients with unmet palliative care needs. Define advance care planning, goals of care, code status and treatment limitations and describe how these are codified in legal and medical documents Interpret a POLST (Physician Orders for Life Sustaining Treatment) form and describe its use in acute care settings	<i>System navigation for patient centered care</i> <i>Physician role in healthcare systems</i>	Lecture – EM faculty content expert
2	Prognosis and trajectory	Describe four common trajectories of life-limiting illness Define prognosis and describe 3 strategies to assess prognosis in ED patients with serious illness	<i>Diagnosis, treatment, and clinical reasoning</i>	Case-based learning – EM content expert
3	Chaplain chat	Describe the role of the chaplain in the interdisciplinary care of seriously ill patients in the ED	<i>System navigation for patient-centered care</i> <i>Interprofessional and team communication</i>	Case-based learning – chaplain
4	Non-pain symptom management	Choose appropriate first- and second-line treatment for seriously ill patients experiencing nausea and vomiting in the ED Choose appropriate first- and second-line treatment for seriously ill patients experiencing dyspnea in the ED Choose appropriate first- and second-line treatment for seriously ill patients experiencing constipation in the ED	<i>Pharmacotherapy</i> <i>Diagnosis, treatment, and clinical reasoning</i>	Case-based learning – hospital palliative care specialist
5	Ask a consultant	Describe the role of the HPM clinician in the care of seriously ill patients in the hospital Understand the role of HPM consultation in the emergency department	<i>Interprofessional and team communication</i>	Case-based learning – hospital palliative care specialist
6	Intro to hospice	Describe the scope of hospice services and the settings where it can take place Identify patients who may qualify for hospice and how to get them evaluated Provide goal concordant care to patients enrolled in hospice who present to the ED	<i>System navigation for patient-centered care</i> <i>Physician role in healthcare systems</i>	Lecture – community hospice medical director
7–10	Serious illness communication workshop (VitalTalk)	Practice skills associated with goals of care conversations with a simulated patient.	<i>Patient- and family-centered communication</i>	Simulation and standardized patient skills-based practice – EM content expert

*ACGME, Accreditation Council for Graduate Medical Education; ED, emergency department; EM, emergency medicine; HPM, hospice and palliative medicine.

small group and they role-play delivering serious news with EM-based scenarios. This session builds skills around delivering serious news.

The curriculum underwent iterative adjustments informed by informal feedback from both facilitators and residents.

Modifications were made based on facilitator availability and interest, resulting in the inclusion or modification of topics, while certain subjects, such as opioid pain management, were removed due to redundancy in other educational settings.

SURVEY DEVELOPMENT

Before implementing the curriculum, we created a brief, pre-bootcamp survey to assess residents' prior exposure and familiarity with palliative care. Subsequently, two post-surveys were used to gauge residents' perceptions regarding the achievement of session-specific goals. We developed the first survey to evaluate the first three weeks of the bootcamp. The initial development collected all potential survey items that were refined through expert consultation. The survey used a five-point Likert scale ranging from 1 (strongly disagree) to 5 (strongly agree). The survey items had been pilot tested and refined in preceding years to ensure question clarity (Appendix 1).

A second survey, which was used for the simulation-based session, prompted residents to rate their self-assessed confidence surrounding the specific skills on conducting GOC conversations covered in the session (Appendix 2). The survey uses a five-point Likert scale ranging from 1 (not very confident) to 5 (very confident).

IMPACT/EFFECTIVENESS

The curriculum evaluation took place during the 2022 bootcamp. Each session had an average of 8–10 PGY-2 residents, of a total 17 potential participants. Attendance varied from week to week due to excused absences. Participation in both pre- and post-surveys was voluntary. Of the eligible residents, nine (52%) completed the pre-survey, revealing that all but one resident had prior exposure to a palliative care rotation during medical school, and 7 of 9 respondents (77%) reported previous communication skills training during their PGY-1 year.

Post-intervention surveys were collected after each session, with completion rates ranging from 25% (2/8 participants) to 70% (7/10 participants) per session. Notably, all respondents indicated agreement or strong agreement with the achievement of each session's objectives. For the simulation-based communication session, 88% (8/9) reported increased confidence overall, 88% (8/9) of residents reported increased confidence in responding to strong emotions, and 100% (9/9) reported enhanced confidence in eliciting patient goals and values.

TIPS FOR SUCCESS/CHALLENGES/LESSONS LEARNED

Several key themes emerged regarding the implementation of a bootcamp curriculum in primary palliative care for EM residents. One notable advantage of this curriculum is its longitudinal format, spanning four consecutive weeks with short intervals between sessions. This structure affords residents the opportunity to practice newly acquired skills while actively working in clinical settings, fostering continuous reflection and refinement of their abilities. Additionally, the curriculum is adaptable and

enables its implementation in programs lacking EM palliative care-trained faculty. Programs can use local resources such as institutional palliative specialists, interdisciplinary palliative teams, or several publicly available online resources.^{9,10,14}

However, despite its strengths, our curriculum faces several challenges. Notably, residents unable to attend sessions risk missing valuable educational opportunities, as the curriculum is not repeated during the academic year. Moreover, limited opportunities for ongoing skill acquisition and feedback outside scheduled sessions may hinder residents' ability to fully integrate palliative care principles into their practice. Furthermore, individual programs may be unwilling to invest 10 hours of curriculum to this specific topic and skillset. Lastly, while there was no cost for the simulation time and standardized patients at the study institution, there may be cost associated with this in other programs and this must be considered.

Furthermore, while participants expressed satisfaction with the curriculum, the outcomes data lack the rigor necessary to definitively establish its success. The impact of this curriculum on long-term knowledge or clinical behavior within the ED remains uncertain. It will be important to conduct more formal assessments of the curriculum objectives and to evaluate its application in the clinical setting.

CONCLUSION

As the role of primary palliative care in emergency medicine continues to evolve, there is a growing need to integrate these essential skills and concepts into all EM residencies. The bootcamp format has proven to be a valuable educational tool in our program, and its effectiveness warrants further exploration and dissemination within the broader EM community.

Address for Correspondence: Julie Cooper, MD, University of Pennsylvania, Department of Emergency Medicine, 3400 Spruce St., Philadelphia, PA 19104. Email: julie.cooper@penmedicine.upenn.edu

Conflicts of Interest: By the *WestJEM* article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. No author has professional or financial relationships with any companies that are relevant to this study. There are no conflicts of interest or sources of funding to declare.

Copyright: © 2024 Cooper et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. Smith AK, McCarthy E, Weber E, et al. Half of older Americans seen in emergency department in last month of life; most admitted to hospital, and many die there. *Health Affairs*. 2012;31(6):1277–85.
2. Shoenberger J, Lamba S, Goett R, et al. Development of hospice and palliative medicine knowledge and skills for emergency medicine residents: using the Accreditation Council for Graduate Medical Education milestone framework. *AEM Educ Train*. 2018;2(2):130–45.
3. Quill TE and Abernethy AP. Generalist plus specialist palliative care — creating a more sustainable model. *N Engl J Med*. 2013;368(13):1173–5.
4. Back AL, Fromme EK, Meier DE. Training clinicians with communication skills needed to match medical treatments to patient values. *J American Geriatrics Society*. 2019;67(S2):S435–41.
5. Baylis J, Harris DR, Chen C, et al. Palliative and end-of-life care education in Canadian emergency medicine residency programs: a national cross-sectional survey. *CJEM*. 2019;21(2):219–25.
6. Accreditation Council for Graduate Medical Education. Emergency medicine milestones. 2021. Available at: <https://www.acgme.org/specialties/emergency-medicine/milestones/>. Accessed February 7, 2024.
7. Loffredo AJ, Chan GK, Wang DH, et al. United States best practice guidelines for primary palliative care in the emergency department. *Ann Emerg Med*. 2021;78(5):658–69.
8. Kraus CK, Greenberg MR, Ray DE, et al. Palliative care education in emergency medicine residency training: a survey of program directors, associate program directors, and assistant program directors. *J Pain Symptom Manage*. 2016;51(5):898–906.
9. Northwestern Medicine. EPEC: Education in palliative and end-of-life-care. Available at: <https://www.bioethics.northwestern.edu/programs/epec/index.html>. Accessed February 7, 2024.
10. Nguyen D and Matese T. Assessing the effect of a hospice and palliative medicine curriculum on emergency medicine residents' knowledge. *Am J Hosp Palliat Care*. 2023;40(5):462–7.
11. Stanich J, Sunga K, Loprinzi-Brauer C, et al. Teaching palliative care to emergency medicine residents using gamified deliberate practice-based simulation: palliative gaming simulation study. *JMIR Med Educ*. 2023;9:e43710.
12. Benesch TD, Moore JE, Breyre AM, et al. Primary palliative care education in emergency medicine residency: a mixed-methods analysis of a yearlong, multimodal intervention. *AEM Educ Train*. 2022;6(6):e10823.
13. Vital Talk. VitalTalk. Available at: <https://www.vitaltalk.org/>. Accessed February 21, 2024.
14. Center to Advance Palliative Care. CAPC clinical training. Available at: <https://www.capc.org/clinical-training/>. Accessed February 7, 2024.

Making A Difference: Launching a Multimodal, Resident-Run Social Emergency Medicine Program

Naomi P. Newton, MD*
Christopher Freeman, MD†
Patricia Panakos, MD†

*Emory University School of Medicine, Department of Emergency Medicine, Atlanta, Georgia

†Jackson Health System, Department of Emergency Medicine, Miami, Florida

Section Editor: Paul Logan Weygandt, MD

Submission history: Submitted October 17, 2023; Revision received April 28, 2024; Accepted May 30, 2024

Electronically published August 16, 2024

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: [10.5811/westjem.18509](https://doi.org/10.5811/westjem.18509)

Introduction: Social medicine seeks to incorporate patients' social contexts into their medical care. Emergency physicians are uniquely positioned to address social determinants of health (SDoH) on the frontlines of the healthcare system. Miami-Dade County (MDC) is a diverse and socially vulnerable area. In 2020, the University of Miami-Jackson Health System (UM-JHS) emergency medicine (EM) residency program launched a multimodal, resident-led Social EM program to identify and address SDoH in the emergency department (ED).

Methods: We use a four-pillar approach to SDoH in the ED: Curriculum Integration; Community Outreach; Access to Care; and Social Justice. Residents graduate with a knowledge of Social EM principles through an 18-month curriculum, an elective, and a longitudinal track. We developed sustainable initiatives through interdepartmental and community-based partnerships, including a Narcan distribution initiative, an ED-based program linking uninsured patients to follow-up care, a human trafficking education initiative, and a quality improvement initiative for incarcerated patients.

Results: Given that the 18-month curriculum was launched in 2022, a full rotation of the curriculum had not been completed as of this writing, and data collection and analysis is an ongoing process. The initial pretest and post-test survey data show improvement in knowledge and confidence in managing Social EM topics. The Narcan initiative has screened 1,188 patients, of whom 144 have received Narcan. The ED-based patient navigation program has enrolled 31 patients to date, 18 of whom obtained outpatient care. Analysis of the impact/effectiveness of the program's other initiatives is ongoing.

Conclusion: To our knowledge, this is one of the most robust social EM programs to date, as many other programs primarily focus on service opportunities. Rooted in the revised principles of Bloom's taxonomy of cognitive learning, this program moves beyond understanding Social EM tenets to generating solutions to address SDoH in and outside the ED. [West J Emerg Med. 2025;26(1.2)5–13.]

BACKGROUND

Social medicine, or the incorporation of patients' social contexts into their medical care, has become a vibrant, interdisciplinary movement that has gained traction in medical schools, residencies, and at the national level. Social medicine emphasizes the importance of social determinants of health (SDoH), or "the conditions in the environments where people are born, live, learn, work, play, worship, and age that affect a wide range of health, functioning, and

quality-of-life outcomes and risks."¹ The US Department of Health and Human Services lists five core SDoH to consider during patient care: economic stability; education access and quality; healthcare access and quality; neighborhood and built environment; and social and community context.¹

Although SDoH can be applied to all specialties, they are perhaps most relevant to emergency medicine (EM). Passage of the Emergency Medical Treatment and Labor Act (EMTALA) in 1986² was acknowledgment that emergency

physicians are often the only link to the healthcare system for patients with financial limitations. Emergency physicians are estimated to provide two-thirds of acute care for all uninsured patients and half of acute care for all Medicaid patients.³ Whether they are rushing a patient to CT, leading their team during a resuscitation, or evaluating patients in a crowded hallway, emergency physicians are immersed in longstanding, complex social issues: trauma; poverty; homelessness; mental health disorders; etc. Therefore, recognizing the effects of SDoH on patient care is critical in the ED.

Jackson Memorial Hospital (JMH) is the primary training site for the University of Miami-Jackson Health System (UM-JHS) EM residency program. It is also the third largest public hospital in the country. The UM-JH Social EM program was launched in 2020 to improve the incorporation of patients' social contexts into their care.

Needs Assessment

When designing a Social EM program, keeping the residency's location and patient population in mind is important. Like most EDs across the nation, the JMH ED is a place of refuge for patients whose SDoH may prevent them from obtaining care elsewhere. As a safety-net hospital in the seventh most populous county in the nation,⁴ JMH serves a particularly diverse patient population with striking socioeconomic needs. The UM health system conducted formal needs assessments of Miami-Dade County (MDC) from 2019–2022, and the UM-JHS Social EM program was designed to reflect these needs.^{4,5}

The UM-JHS Social EM program was designed to ensure that all residents graduate with the ability to incorporate their patients' SDoH into ED care regardless of their ultimate practice locations. However, certain aspects of this program were designed to address the unique needs of MDC—a “minority-majority” community that experiences challenges with English proficiency, and in which 20% of the population lives below the poverty level.^{4,5}

PROGRAM GOALS

Bloom's taxonomy of cognitive learning objectives outlines six levels in the cognitive domain: knowledge; comprehension; application; analysis; synthesis; and evaluation.⁶ Over time, scholars have sought to revise this framework and, when taken as a whole, these revisions place less emphasis on a linear progression through each level.⁶ Instead, there is an increased focus on generating new hypotheses and developing projects that use and expand upon the acquired knowledge.⁶ Therefore, the UM-JHS Social EM program seeks to shift its participants from purely understanding SDoH as they pertain to EM, to generating effective solutions for addressing these SDoH in and outside the ED. The Social EM program outlines six goals for its residents, who are then tasked with generating effective

solutions and designing their own measurable outcomes for each goal. Upon successful completion of this program, residents should be able to:

1. Define and identify SDoH in the ED and apply these principles to bedside care.
2. Engage with MDC outside the ED and address its social and medical needs through longitudinal involvement in local outreach initiatives.
3. Solidify and share acquired knowledge through an 18-month, multimodal curriculum.
4. Identify and seek to address barriers to medical care experienced by patients who use the ED as their main source of healthcare.
5. Identify and seek to address recurrent social justice issues encountered in the ED.
6. Enact positive change through advocacy and quality improvement initiatives at hospital-wide, local, and/or national levels.

PROGRAM PARTICIPATION

Since its launch in 2020, the program has been divided into four pillars that address core areas within Social EM: Curriculum Integration; Community Outreach; Access to Care; and Social Justice (Figure 1). Initiatives within each pillar will be discussed in a separate section. Anyone affiliated with the UM-JHS ED can participate in initiatives across all four pillars. Many of these initiatives are longitudinal, allowing for varying levels of participation throughout residency. Additionally, this program also offers leadership, peer teaching, and scholarly opportunities that may count toward existing residency requirements.

The UM-JHS has a three-year EM residency program, and each of its classes (postgraduate years [PGY] 1–3) is comprised of 14–15 residents. EM residents are not required to participate in the Social EM program but are encouraged to do so. They may choose to serve as program leaders (Figure 1), participate in the longitudinal track and/or two-week elective (discussed in detail in subsequent sections below), or to participate in individual initiatives as their schedules allow. However, Social EM program leadership developed a formal curriculum to ensure that all residents graduate with a solid understanding of core Social EM principles, regardless of their level of involvement with the program; this will be discussed in a separate section.

CORE LEADERSHIP HIERARCHY

This program was designed to be executed by residents in collaboration with faculty, medical students, and staff. The program was structured into a core leadership hierarchy to appropriately divide the labor of designing and launching initiatives that pertain to each pillar, while ensuring that residents complete their existing clinical and academic




Figure 1. Social emergency medicine program organization and division of labor. Program directors consist of one faculty director and 1–4 resident directors (EM residents selected via a formal application process). Program directors oversee initiatives across all four pillars but spend additional time leading Curriculum Integration initiatives to ensure a seamless incorporation of Social EM principles into residency training. Pillar leaders are EM residents who are selected by program directors via a formal application process; they design and oversee initiatives in their assigned pillars. Faculty mentors are generally core faculty in the EM department with expertise in their assigned pillar. However, faculty in other specialties at UM-JHS may also serve as mentors if they currently oversee a community or hospital-based initiative that collaborates with the Social EM program. (For example, a faculty mentor from the family medicine department oversees the IDEA Needle Exchange Clinic.) Anyone affiliated with the EM department may serve as a team member. Team members work directly with their assigned resident leaders and divide the tasks required to launch and publicize initiatives.

requirements (Figure 1). This leadership hierarchy organizes, executes, and publicizes the program and its initiatives.

Directors

A faculty director and at least one resident director oversee the program together (Figure 1). The original directors, Patricia Panakos, MD, and Naomi Newton, MD, authored this paper and conceptualized the program together in Fall 2020. The collaboration between Drs. Panakos and Newton was borne from a shared passion for social medicine and a desire to implement an EM residency-based program to address the SDoH of patients in MDC. Dr. Panakos is the associate program director for the UM-JHS EM residency and has undergone formal training in curriculum development. She has also developed ED-based public health initiatives at JHS, such as a universal screening program for communicable diseases, including HIV, hepatitis C, and syphilis. Dr. Panakos continues her role as faculty director for the social EM program. Dr. Newton is an alumna of the UM-JHS EM residency and served as chief resident during her final year of training. She assumed the role of the social EM program’s resident director as a PGY-1 and transitioned her position upon her graduation in 2023. She has also collaborated with Dr. Panakos on public health

initiatives, including a universal HIV screening initiative in JHS’s pediatric ED. Dr. Newton is currently pursuing a two-year fellowship in health policy and advocacy at Emory University.

Given that there was no precedence for such a program at UM-JHS, Drs. Panakos and Newton worked almost daily to create the program and maintain its sustainability, while also completing their existing clinical and academic responsibilities. Drs. Panakos and Newton designed the program’s overall structure, created a formal selection process for pillar leaders, and identified community partners and faculty mentors with expertise in Social EM. They presented a formal proposal that was approved by both the chair of the ED at JMHS and the UM-JHS EM residency program director in October 2020. They also designed and launched the 18-month curriculum, two-week elective, and longitudinal track, which are described in subsequent sections of this paper. To account for continued program growth, the original directors selected four new resident directors for the 2023–2024 academic year via a formal application process (Figure 1).

Directors approve proposed initiatives across all pillars and work directly with pillar leaders to track progress and troubleshoot challenges. They check in remotely with pillar

leaders at least bi-monthly to ensure timely project completion. They also promote the program at a departmental and hospital-wide level and help pillar leaders identify faculty and community partners (Figure 1). Resident directors are recognized with a Social EM leadership award upon their graduation.

Resident Leaders

An average of two PGY-1 or PGY-2 EM residents lead each pillar. Interested residents apply for this position via a brief electronic application (Google Survey) at the start of the academic year and are selected by the directors. Residents generally do not serve as leaders of more than one pillar, as this position must be balanced with existing residency obligations. Resident leaders report directly to the directors and dedicate an average of two to four hours per week to their roles. As leaders progress through training, they may either remain in their leadership roles or transition their roles to incoming PGY-1s and PGY-2s. All resident leaders who have served for at least one year are recognized with a special award upon graduation from residency.

Leaders focus on designing initiatives that pertain to the goals of their assigned pillar. They identify appropriate partners within JHS and MDC to aid in developing and launching these initiatives (Figure 1). Partners include JHS faculty (including those in non-EM specialties), local outreach organizations (many of which already had established relationships with UM-JHS through medical student involvement), and other JHS-affiliated residency programs (eg, pediatrics, internal medicine, family medicine). Interdisciplinary collaboration prevents the Social EM program from “re-inventing the wheel” and helps initiatives achieve success with fewer funding, resource, and logistical restrictions. Resident leaders delegate day-to-day tasks to an interdisciplinary team to divide the labor of executing these initiatives. Leaders are required to check in remotely with their team members at least monthly to discuss progress on pillar initiatives.

Interdisciplinary Teams

Team members divide the tasks required to launch initiatives within their assigned pillar. They are required to dedicate a minimum of one to two hours per week on these tasks and check in regularly with their pillar leaders as previously discussed. Those who desire to do so may participate in more than one pillar team. Participation in a pillar team is open to anyone in the UM-JHS ED. However, during the first three years of the program, teams were primarily comprised of EM-bound UM medical and pharmacy students, JHS clinical pharmacy residents, and hospital staff (eg, nurses and social workers).

PROGRAM DESIGN: THE 4-PILLAR APPROACH

In the following section, we provide a broad overview of each pillar’s objectives and highlight several key initiatives within each pillar. When relevant, please see the corresponding appendices for additional details.

Curriculum Integration

This pillar incorporates the tenets of Social EM into residency training to empower future generations of emergency physicians to apply Social EM principles to their care. This is the only pillar that requires all EM residents to participate because its initiatives have been incorporated into the existing residency curriculum. Doing so ensures that all EM residents graduate with an understanding of SDoH and the principles of Social EM, regardless of their level of involvement in other pillars. Of note, approval from the institutional review board was not required for the development of this curriculum.

We developed and launched a multimodal, 18-month Social EM curriculum that has been incorporated into the existing 18-month residency didactic schedule (Appendix A). The curriculum covers 18 core social EM topics (Table 1) and includes journal clubs, simulation cases, lectures, problem-based learning, and interactive group discussions. The curriculum is led by faculty and residents with expertise or

Table 1. 18 core areas of study were chosen to be covered monthly during the 18-month Social EM curriculum. This curriculum is integrated into standard residency didactic schedule, which repeats every 18 months. Using a multimodal learning format, topics can be presented as traditional lectures, case-based discussions and journal clubs (“Cases”), or simulations. The initial modalities for each topic are listed below; the modalities used for each topic will change every 18 months (eg, the pediatric health lecture would be presented as either a case or simulation 18 months later). Additional details regarding logistics and implementation can be found in Appendix A.

Lectures	Cases	Simulations
1. Social determinants of health	7. Implicit bias/racism	13. Human trafficking and domestic violence
2. Healthcare coverage and access	8. Homelessness	14. Substance abuse and harm reduction
3. Financial stability	9. Health literacy	15. Caring for incarcerated patients
4. Frequent ED utilizers	10. Immigration	16. Highly communicable diseases/STI epidemics
5. Women’s health	11. Resource insecurity	17. Language and cultural barriers to healthcare
6. Pediatric health	12. Trauma-informed care	18. Gender identity

ED, emergency department; STI, sexually transmitted infection.

interest in the core topics. Social EM leadership assists presenters in identifying learning objectives for each session, selecting topics, and developing content. All conference attendees participate in pre- and post-surveys to assess their baseline knowledge and the effectiveness of each didactic session. Residents are also asked to evaluate the Social EM curriculum during the annual residency program evaluation. Surveys and results are discussed further in the Impact/ Effectiveness section of this manuscript.

In 2022, we launched the two-week Social EM elective for residents who desire a more in-depth experience with the program (Appendix B). This elective is open to PGY-2 EM residents during their elective block and is comprised of service opportunities, self-directed study, peer teaching, and initiative participation across all pillars. The PGY-2 rotation schedule is designed so that only one resident completes an elective in any given month. Therefore, the experience is personalized for each participating resident. Social EM directors work with the resident ahead of time to design an elective schedule that ensures participation across all pillars but allows them to engage more deeply within their pillar(s) of interest (Appendix B).

We also designed a longitudinal track that was launched the 2023–2024 academic year. Track participants engage in a

set number of outreach events, quality improvement initiatives, peer teaching, and self-directed study over 18 months. The requirements are based on a point system that ensures engagement with all pillars but allows for deeper exploration in areas of individual interest. Residents must reach a minimum of 30 points to complete the track (Figure 2). Requirements include a longitudinal scholarly activity that culminates in a presentation at the local, regional, or national levels (eg, developing a project to address food insecurity). They must also participate in the Social EM didactic curriculum through peer teaching, developing new elements to the curriculum, and mentoring medical students. Participants log their progress via an online form and must attend a minimum of nine monthly track meetings with the Social EM directors over an 18-month period. Upon graduation, residents who complete the track will receive a Distinction in Social EM.

Community Outreach

This pillar was designed to establish a meaningful presence in MDC beyond bedside care and to address social issues through partnerships with local organizations. For example, through a partnership with Miami Street Medicine, participants join an interdisciplinary team in providing

Activity	Hours Spent	Personal investment value	Total Points			
Caring with Compassion Module	4	0.5	2	All residents must participate in the following activities at a minimum		
Leadership of a Pillar	12	1.5	18	QTY	Activity	PTS
IDEA Clinic	4	0.5	2	1	CC Module	2
Miami Street Medicine	3	0.5	1.5	1	Curriculum Development	3
DOCS Fair	5	0.5	2.5	2	Community Outreach	1.5
Stop the Bleed Training Program	2	0.25	0.5	1	Access to Care	1
Stop the Bleed Lecturing	3	0.5	1.5	1	Social Justice	1
Coordinating care for 6 patients	4	0.25	1	1	JC Presentation	1
Writing a high utilizers brief	2	0.5	1	9	Track Meetings	2.25
Food Insecurity - Distribution Event	4	0.5	2			11.75
Human Trafficking Training Program	2	0.25	0.5			
Human Trafficking Lecturing	1	0.5	0.5		Minimum Points	30
Developing/Updating QI Protocol	4	1	4			
Attending advocacy event	6	1	6			
Developing a case/sim	4	0.75	3			
Developing a lecture	6	1	6			
Presenting article at JC	2	0.5	1			
Peer-reviewed publishing in Social EM	8	1.5	12			
Non-peer reviewed publishing in Social EM	4	1	4			
Developing a new Social EM project	12	1.5	18			
Social EM Elective	4	1	4			
Attending track meetings	1	0.25	0.25			
*Other pre-approved social EM activity	Varies	Varies	Varies			
		Total	91.25			
Legend:						

Figure 2. Point system for the 18-month longitudinal track. Note that opportunities in each pillar may vary over time. This figure lists opportunities from the fall of 2023.

regular medical care at locations commonly occupied by Miami's homeless population. Through a partnership with the Stop the Bleed Campaign, participants undergo formal training to serve as local instructors. Participants then lead workshops that teach non-medical community members to perform bystander cardiopulmonary resuscitation and stabilize victims of violence until first responders arrive. Participants may serve as instructors for Stop the Bleed events throughout MDC, as their schedules allow.

This pillar launched a Narcan program in July 2022, in partnership with the UM IDEA (Infectious Disease Elimination Act) Needle Exchange Clinic and the UM Michael Wolfson Department of Community Service (DOCS). This program seeks to address South Florida's opioid epidemic and is in keeping with the statewide Emergency Treatment for Suspected Opioid Overdose Act.⁷ At community health fairs, participants provide free opioid use disorder (OUD) screening, based on Diagnostic and Statistical Manual of Mental Disorders, 5th ed, criteria.⁸ Narcan is subsequently distributed to those identified to be at high risk for life-threatening overdoses, and additional OUD counseling and training on safely administering Narcan are provided.⁹

Access to Care

This pillar links chronically ill patients, high ED utilizers, and the uninsured to outpatient care. It also seeks to centralize existing social support resources within UM-JHS and efficiently address SDoH at the bedside. Through a partnership with DOCS, uninsured patients presenting to the ED with chronic complaints are paired with long-term patient navigators, who help them secure affordable outpatient care upon discharge.

The High Utilizers Initiative aims to streamline the care of patients who frequently visit the ED. Participants conduct chart reviews of individuals flagged as frequent utilizers in the electronic health record and create patient care briefs that auto-populate in their charts. These patients often present to the ED numerous times a week and receive care from different clinicians each time. By consolidating their pertinent medical information, these briefs allow for better, more streamlined patient care with less repetition of tests and procedures. The briefs also lessen the cognitive load of the clinician, decreasing the time spent on chart reviews and helping guide future care.

Many patients present to the ED with conditions that are exacerbated by a lack of basic resources. It is challenging to address these complex SDoH amidst the time constraints of ED care, and EDs can no longer rely heavily on social workers for assistance, due to the nationwide social worker shortage.¹⁰ This pillar partnered with Miami Street Medicine and the JMH Pharmacy Department to create resource guides for patients and clinicians in response to this need. Community resource guides (in English, Spanish, and

Haitian Creole) provide information for affordable outpatient clinics, prescriptions, mental health services, temporary housing, meal programs, and substance use treatment centers, as well as resources for pregnant patients and victims of domestic violence. Clinician resources include referral information for resident-run subspecialty clinics, instructions for initiating buprenorphine in the ED and referring patients to medication-assisted treatment clinics, and algorithms for human trafficking screening.

Social Justice

This pillar tackles health inequity and injustice issues through interdisciplinary education and quality improvement initiatives. The Human Trafficking Education Ambassador program, in partnership with JMH's Rape Treatment Center, teaches clinicians to screen for and treat victims of human trafficking. Florida has the third highest number of human trafficking cases in the nation, and MDC is, sadly, a known trafficking hub.¹¹ Trained residents lead interactive seminars, sharing HIPAA-compliant trafficking cases and teaching clinicians to identify and address red flags for trafficking.

This pillar also seeks to improve care for incarcerated patients in the ED, particularly concerning patient privacy and examinations in the presence of law enforcement. Initiatives include a recently published review on the barriers to caring for this population and recommendations to improve their delivery of care.¹² We also implemented a simulation session on caring for incarcerated patients into residency didactics.

IMPACT/EFFECTIVENESS

Curriculum Integration

Social EM leadership is in the process of completing a formal impact assessment of the curriculum integration pillar of the program via a single-group, pretest-posttest design.⁶ Brief pre- and post-didactic session surveys are designed for each Social EM topic in the 18-month curriculum. Surveys are designed to assess baseline knowledge of the topic and the changes in this baseline knowledge after the session. Survey questions also address relevant epidemiological statistics and useful community resources for addressing the topic in MDC. Each post-survey ends with a blank section for participants to write in any additional feedback, which Social EM program leadership uses for subsequent didactic sessions.

For convenience, these surveys are administered via electronic forms; conference attendees scan QR codes to the forms before and after the session. All residents, faculty, students, and staff in attendance are eligible for participation in the surveys. However, thus far, survey participation has generally been limited to resident attendees, as faculty, staff, and student attendance is less consistent. Hospital badge

numbers are used to compare individuals' changes in pre- and post-session responses.

Since the 18-month curriculum was launched in 2022, a full rotation of the curriculum has not been completed as of this writing, and data collection and analysis is ongoing. However, thus far, the curriculum topics have been well-received, with residents indicating an improved confidence in their ability to recognize and address these Social EM issues at the bedside. For example, [Figure 3](#) shows key results from the pre- and post-surveys administered during the first session of the formal curriculum in 2022—a simulation session on highly communicable diseases/sexually-transmitted infection (STI) epidemics ([Table 1](#)). These results suggest efficacy in improving baseline knowledge and confidence with the topic of acute HIV in the ED, including epidemiology, community resources, and initiating either highly active antiretroviral therapy or pre-exposure prophylaxis when indicated.

An impact assessment of the two-week elective is pending, as only one PGY-2 resident had completed at the time of this manuscript's development.

Community Outreach

It is challenging to concretely assess the impact of the Community Outreach pillar, as its service-driven initiatives are generally qualitative in nature. However, initial data

from the Narcan Initiative highlights its impact on MDC. As of May 2023, the program screened 1,188 patients across MDC, of whom 144 received Narcan. In recognition of the Narcan Initiative's current impact and continued growth, JMH's Department of Emergency Medicine received the 2023 University of Miami Mitchell Wolfson Sr. Department of Community Service award.

Access to Care

We are currently in the early stages of data collection to analyze the success of the Access to Care initiatives. Thus far, the patient navigation program has enrolled 31 ED patients. Of these patients, 18 were able to successfully complete their navigation goals and obtain outpatient care. This program has particularly benefitted non-English-speaking patients, whose language barriers can hinder their ability to navigate a complex system. For example, navigators were able to link a Spanish-speaking patient to outpatient oncologic care for her untreated gynecologic cancer. Recently, a homeless, uninsured patient living at the Miami Rescue Mission (MRM) was treated for an acute ulcerative colitis flare in the ED. After he was discharged, the navigators ensured that he obtained timely follow-up at an MRM-affiliated gastroenterology clinic, a student-run clinic staffed by UM faculty. We are continuing to publicize this program and encourage emergency clinicians

N = 30 residents	Pre-Acute HIV Sim	Post-Acute HIV Sim
% Correct	43%	92%




Figure 3. Comparison of key pre- and post-survey results after a simulation session on highly communicable diseases/STI epidemics; 30 residents completed the surveys. After the session, residents indicated an increased confidence in their ability to recognize acute HIV and initiate Highly Active Antiretroviral Therapy (HAART) or Pre-Exposure Prophylaxis (PrEP) treatment when indicated. Most residents found the session beneficial in learning about community resources for ED patients with HIV, as well as prescribing HAART and PrEP.

to enroll their patients during their shifts. We are still in the data collection phase of the High Utilizers initiative.

Social Justice

The Social Justice pillar initiatives experienced several launch delays due to COVID-19 pandemic restrictions and faculty turnover. Initiatives were officially launched in the 2022–2023 academic year, and data regarding their impact and effectiveness is pending. Thus far, human trafficking education ambassadors have given well-received lectures to JHS-affiliated clinics and to JMH's family medicine, pediatrics, and internal medicine residencies.

Overall Program Feedback and Support

Since the program is under the direct guidance of a current residency associate program director, there is continual communication between Social EM directors and EM residency leadership. Residency leadership actively engages with and provides insights into pillar initiatives, leading to timely changes to the program when deemed necessary. For example, previous feedback led to the development of the elective and longitudinal tracks. Residents in the core Social EM leadership team also obtain regular qualitative feedback from their peers and share this feedback with the Social EM directors. This program is also reviewed during the annual residency program evaluation committee meeting. This program has full EM departmental support.

Additional Recognition/Awards

Overall, this robust, multimodal, resident-led Social EM program has rapidly grown over the last three years, despite the COVID-19 pandemic. In 2023, six of the 14 PGY-3 residents graduated with a Social EM distinction. The program's interdisciplinary nature ensured its success, as multiple initiatives were launched without significant funding or administrative restrictions. The program is receiving increasing recognition. In addition to the previously mentioned community service award for the Narcan Initiative, the MDC chapter of the Stop the Bleed Campaign received a 2021 award from the mayor for its education initiatives in local high schools. In 2023, we were also honored to receive the 2023 ACEP Social EM Section Distinguished Program Award.

LIMITATIONS

Residents' availability often limits consistent participation in Social EM. Residents have multiple clinical and academic responsibilities, and as they progress through training, their time is further limited by searching for jobs and applying for fellowships. In response to this limitation, the elective and longitudinal track were developed to allow for flexible but regular participation, as many requirements can be

completed during lighter rotations. The didactic curriculum also ensures that all residents will graduate with the same baseline knowledge of Social EM tenets. Additionally, the Social EM leadership will transition every two years, allowing junior residents with leadership roles to pass on their duties to incoming residents as they become senior residents.

Certain aspects of this program were designed to address some of the social issues that are particularly prevalent in MDC and may not be generalizable to other EM residency programs in the United States. Other residency programs seeking to develop their own Social EM initiatives should consider the unique needs of their patient populations when doing so.

The program's first three years were dedicated to overall development, garnering participants, finding community partners, and launching initiatives in each pillar. Therefore, data collection to formally assess the program's impact and effectiveness is still in process and is currently limited to initial data (unblinded pre- and post-tests completed by resident participants) from the launch of the 18-month didactic curriculum. This data may also be subject to selection bias, as most residents, faculty, students, and staff are excited about the Social EM program and want it to succeed.

CONCLUSION

The University of Miami-Jackson Health System Social EM program was launched in 2020 to address the SDoH of patients in Miami-Dade County—an area of significant medical and social need. It targets critical social issues through four pillars: Curriculum Integration; Community Outreach; Access to Care; and Social Justice. This multimodal, resident-run program achieved rapid success in three years by developing sustainable initiatives in partnership with local organizations and other UM-JHS departments. Rather than focusing solely on service opportunities, this program enhances residents' knowledge of SDoH, fosters the development of quality improvement initiatives, and provides opportunities to create meaningful change in the ED and the community. This program also provides residents with leadership and scholarly opportunities. We hope that this article will inspire other residencies to develop similar programs.

ACKNOWLEDGMENTS

The University of Miami-Jackson Health System Social Emergency Medicine program would not have been possible without the ingenuity, dedication, and passion of the following individuals and organizations. Thank you for everything that you have done and continue to do to create

positive change: Rachel Armstrong, MD; Tatiana Barriga, MD; Kelley Benck; Daniel Bergholtz, MD; Emily Brauer, MD; Sandra Cabrera, MD; Danielle Cohen, MD; Emily Dawra, MD; Luis De La Rosa, RN; Shaheen Emami; Joshua Goldstein, MD; Jamie Harris; Kristopher Hendershot, MD; Armen Henderson, MD, MBA; Brooke Hensley, MD; Sarah Jabre, MD; Jackson Health System Clinical Pharmacy Residency; Johnnatan Marin, MD; Julia Martinez, MD; Elizabeth May-Smith, MD; Kelly Medwid, MD; Miami Street Medicine; Victoria McKee; Samantha Mosle; Lien Morcate; Kaleb Morris; Judith Oriental-Pierre, MD; Roxcy Bolton Rape Treatment Center; Stop the Bleed; University of Miami IDEA Needle Exchange Clinic; University of Miami, Michael Wolfson Department of Community Service; and Melissa Velasquez, MD.

Address for Correspondence: Naomi P. Newton, MD, Emory University Hospital, 531 Asbury Circle, Annex Suite N340, Atlanta, GA 30322. Email: naomi.newton@emory.edu

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. No author has professional or financial relationships with any companies that are relevant to this study. There are no conflicts of interest or sources of funding to declare.

Copyright: © 2024 Newton et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. U.S. Department of Health and Human Services. Healthy people 2030-social determinants of health. Available at: <https://health.gov/healthypeople/priority-areas/social-determinants-health>. Accessed May 2, 2023.
2. Zibulewsky J. The Emergency Medical Treatment and Active Labor Act (EMTALA): what it is and what it means for physicians. *Proc (Bayl Univ Med Cent)*. 2001;14(4):339–46.
3. American College of Emergency Physicians. Understanding EMTALA. Available at: <https://www.acep.org/life-as-a-physician/ethics-legal/emtala/emtala-fact-sheet/>. Accessed May 30, 2023.
4. U-Health-University of Miami Health System. Community health needs assessment 2019–2021. Available at: <https://umiamihealth.org/-/media/uhealth/chna/umhc-2019-community-health-needs-assessment.ashx>. Accessed May 15, 2023.
5. Florida Department of Health in Miami-Dade County. 2022 Community health assessment. 2022. Available at: https://www.floridahealth.gov/_media/miami-dade/community-reports/miamidade-cha.pdf. Accessed May 15, 2023.
6. Thomas PA, Kern DE, Hughes MT, et al. (Eds.). (2016). Goals and objectives. In: *Curriculum Development for Medical Education* (53–56;130–133). 3rd ed. Baltimore, MD: Johns Hopkins University Press, 2016.
7. Emergency Treatment for Suspected Opioid Overdose. *Chapter 381 Section 887–2021 Florida Statutes*, State of Florida 2021 Legislative Statutes, 1 Jan. 2021.
8. American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders*, 5th ed. (541–542). Washington, DC: American Psychiatric Association.
9. Benk KN, Goldstein J, Newton N, et al. Social emergency medicine: naloxone out of the ED & into the community. 2022. Available at: <https://fcep.org/social-emergency-medicine-naloxone-out-of-the-ed-and-into-the-community/>. Accessed February 3, 2022.
10. Lin VW, Lin J, Zhang X. US social worker workforce report card: forecasting nationwide shortages. *Soc Work*. 2016;61(1):7–15.
11. Miami-Dade State Attorney's Office. Florida facts regarding human trafficking. Available at: <https://humantrafficking.miamisao.com/human-trafficking>. Accessed June 3, 2023.
12. Armstrong RE, Hendershot KA, Newton NP, et al. Addressing emergency department care for patients experiencing incarceration: a narrative review. *West J Emerg Med*. 2023;24(4):654–61.

Effectiveness of a Collaborative, Virtual Outreach Curriculum for 4th-Year EM-bound Students at a Medical School Affiliated with a Historically Black College and University

Cortlyn Brown, MD, MCSO*

Richard Carter, MD[†]

Nicholas Hartman, MD, MPH[‡]

Aaryn Hammond, MD[‡]

Emily MacNeill, MD*

Lynne Holden, MD[§]

Ava Pierce, MD^{||}

Linelle Campbell, MD[§]

Marquita Norman, MD, MBA^{||}

*Atrium Health Carolinas Medical Center, Department of Emergency Medicine, Charlotte, North Carolina

[†]Howard University, College of Medicine, Washington, DC

[‡]Wake Forest University, School of Medicine, Winston-Salem, North Carolina

[§]Albert Einstein College of Medicine, Montefiore Medical Center, Bronx, New York

^{||}UT Southwestern Medical Center, Dallas, Texas

Section Editors: Matthew Tews, MD and Christine Stehman, MD

Submission history: Submitted January 24, 2024; Revision received October 9, 2024; Accepted October 17, 2024

Electronically published December 16, 2024

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.18748

Background: Diversity within the physician workforce is associated with improved clinical outcomes and patient satisfaction. Despite this, the US physician workforce, particularly in emergency medicine (EM), remains relatively homogeneous. Of all Black medical school students in the US, 14% attend the four Historically Black Colleges and Universities (HBCU) that have a medical school. Unfortunately, none of these schools are affiliated with an academic EM program. Because of this, there is less professional mentorship focused on obtaining a career in EM and potentially less formal curricula for senior medical students doing their home sub-internship in EM.

Objectives: Our objective was to fill the gap left by the absence of an academic EM department at Howard University College of Medicine (HUCOM) by creating a collaborative educational experience for fourth-year medical students during their home EM sub-internship. The curricular objectives were to teach core principles of EM, build relationships with students, and prepare them for pursuing EM residency training.

Curricular Design: Four EM academic departments collaborated to create and implement a virtual curriculum using the six-step approach to curricular development.

Impact/Effectiveness: After completion of the course, five students (100%) reported strongly agreeing with the following statements. These sessions 1) helped me learn the approach to core EM topics more than I would have been able to do on my own; 2) helped me learn key skills for excelling in an EM rotation more than I would have been able to do on my own; and 3) allowed me to connect with faculty and resident mentors to learn more about the field of EM. Of these five students, 80% and 20% reported strongly agreeing and agreeing, respectively, that these sessions helped them learn about the process of applying to and selecting an EM residency program. [West J Emerg Med. 2025;26(1.2)63–68.]

INTRODUCTION

Need for Innovation

Medical students interested in emergency medicine (EM) who attend a historically Black college or university (HBCU) do not have the teaching and mentorship that occurs when a medical school is affiliated with an academic EM program. We formed a collaborative program among four academic EM departments to help fill this need for EM-bound students at Howard University College of Medicine (HUCOM). To our knowledge, this is the first such program to be reported in the literature.

Background

A diverse physician workforce is associated with increased access to and utilization of the healthcare system, improved health outcomes and patient experience, and improved fiscal margins for hospitals.^{1-4,6} Despite this, the medical field as a whole has made minimal advances in increasing physician diversity. In 2008 the percentage of Black or Hispanic US physicians from all specialties was 6.3% and 5.5%, respectively. By 2018, however, those percentages were only 5.0% and 5.8%, respectively. Even more concerning given the diverse patient population that the emergency department (ED) serves, EM remains among the medical specialties with the lowest number of physicians from backgrounds under-represented in medicine (URiM). Between 2008–2018, the percentage of emergency physicians who identified as Black decreased from 5.0% to 4.5%, and stayed constant at 5.3% for Hispanic/Latinos.⁷

When surveyed, 35% of EM program directors reported that the small number of URiM residency applicants was the greatest barrier to obtaining a diverse residency class.⁸ Of all Black medical school students in the US, 14% attend four HBCUs with a medical school. Because none of these schools are affiliated with an academic EM program, their medical students have decreased exposure to EM in the pre-clinical years, less professional mentorship focused on obtaining a career in EM, and fewer formal curricula for senior medical students doing their home sub-internship (sub-I) in EM. This lack of mentorship has been identified as a critical barrier for URiM students across various specialties, contributing to lower application rates and residency placement. Studies suggest that mentorship increases both career satisfaction and inclusivity and the likelihood of these students entering and succeeding in competitive fields like EM.^{9,10} In addition, a national survey of clerkship directors found that having a structured, standardized sub-I curriculum significantly improved the preparedness of students for residency, especially when these rotations were affiliated with residency programs.¹¹

The Emory University Department of Emergency Medicine created a program with Morehouse School of Medicine to provide guidance to medical students interested in EM. A total of 115 Morehouse students completed an EM

clerkship at Emory, and 62.6% successfully matched into EM.¹² While this program was successful, students typically rely on their home sub-I to prepare for mandatory away rotations. This absence of support from an academic department prior to away rotations may cause the students to find themselves less prepared and at a competitive disadvantage when they begin their away rotations. Furthermore, many EM residencies are not in proximity to a HBCU, requiring students to bear the financial burden of traveling to other cities and states for their away rotations.

At HUCOM, the EM sub-I relied heavily on an older, recorded online lecture series from an external institution, supplemented by bedside teaching from community attendings at one site, Howard Hospital. Students noted that the absence of formal educational components, such as weekly didactics, journal clubs, and simulation, resulted in limited exposure to “cutting-edge” EM practices. Moreover, the lack of interaction with academic attendings who are dedicated to medical student education, along with the absence of residents—who represent the next step in career progression—left students without access to critical mentorship and guidance. This gap hindered students’ ability to visualize their own progression and receive practical advice from individuals at a similar stage in training, further limiting their connection to the broader EM community.

To help overcome that barrier, we created a collaboration between four academic EDs and HUCOM in an attempt to augment curricular offerings for EM-interested students on their HUCOM fourth-year EM home rotation. The collaboration between four academic EDs broadens the exposure students receive to different teaching styles, institutional cultures, and clinical perspectives. This variety provides a more comprehensive educational experience than what can be offered by a single institution alone.

Objective of Innovation

We aimed to address the absence of an academic ED at HUCOM by developing a collaborative educational experience. This program focuses on core principles of EM and residency preparation and was designed specifically for fourth-year medical students during their home EM sub-I at HUCOM. We obtained institutional board review approval from Wake Forest University School of Medicine.

Development Process

We used the six-step approach to curricular development. All final curricular design and content was agreed upon by the faculty representatives at each of the four participating residency sites.^{13,14} 1) *Problem identification and general needs assessment.* Unlike traditional curriculum development where the need assessment is based on a specific health problem, our needs assessment was based on the need to increase the diversity of emergency clinicians by helping prepare under-represented students to succeed in away

rotations and the match. 2) *Determining and prioritizing content*. While individuals at each participating institution were involved with teaching at their own institution, the needs of the HUCOM students were unique. Therefore, educational objectives were developed in conjunction with the faculty advisor to the fourth-year EM rotation at HUCOM who conducted stakeholder interviews with five current medical students and five alumni who had recently graduated and were currently in EM residencies across the country. It was decided that curricular content would include a mix of core EM topics (as determined from stakeholder interviews) and advising sessions.

After all sessions, students were provided with the contact information for the faculty lecturers and were encouraged to reach out. 3) *Goals and objectives*. Broad curricular goals were developed. These were to a) teach the approach to core complaints in EM; b) teach key skills in EM; c) demystify the process of applying to an EM residency program; and d) connect students with residents and faculty in the field of EM. After this, specific measurable lecture goals were developed based on cognitive, affective, and psychomotor objectives for the learner. 4) *Educational strategies*. We created an entirely virtual, four-week didactic program, with content organized into weekly four-hour blocks, each led by a different academic ED, on an interactive platform that allowed for case-based discussions, small-group discussions, and standard lecture format. Since implementation in 2022, the program has been mandatory for all students completing their fourth-year EM sub-I at HUCOM.

Each week, the sessions required the participation of four to five faculty members who volunteered their time, with the majority of lectures delivered by a single faculty member. However, select sessions, such as the “Application and Interviewing Process,” were co-led by a dynamic team consisting of the assistant program director, program director, and chief residents, providing a well-rounded perspective and valuable insights for the participants. Content was mapped and coordinated, and pre-reading was assigned from the Academy for Diversity and Inclusion in Emergency Medicine webinar series “How to Be a Successful

EM Applicant” and the Clerkship Directors in Emergency Medicine/Society of Academic Emergency Medicine M4 curriculum. Each day included a mix of clinical topics and “advising” sessions (Table 1). 5) *Implementation*. Approval from the EM director was obtained, and the curricula was implemented. 6) *Evaluation and feedback*. After each block of content, evaluations for each individual session (including the presenter) were sent to participating students via REDCap (Research Electronic Data Capture, hosted at Howard University School of Medicine).

These evaluations consisted of one question for each session: “Please rate the effectiveness of the following session in accomplishing its learning objectives: *Session, Presenter*.” At the end of the month-long program, an overall evaluation of the program was sent to participating students, also via RedCap. The program evaluation survey tool, including four multiple-choice questions regarding the overall learning objectives, is reflected in Figure 1. The tool also included two free-response questions: 1) “Which parts of the curriculum were of most value to you?”; and 2) “Which parts of the curriculum could be improved?” We refined the curricula each year during an end-of-year debrief.

Implementation Phase

Prior to the first session, students were provided a spreadsheet with pre-session work, curriculum topics, presenting faculty and residents, dates and times, and links to access the weekly virtual sessions. Each EM program provided four hours of interactive didactics to the students according to the scheduled dates and times.

Outcomes

A post-curricular survey found universal agreement from students that the curriculum was effective in meeting the above goals. Of the five students, 100% reported strongly agreeing with the following statements. These sessions 1) helped me learn the approach to core EM topics more than I would have been able to do on my own; 2) helped me learn key skills for excelling in an EM rotation more than I would have been able to do on my own; and 3) allowed me to

Table 1. Curricula from sample block.

	Didactic session one Institution one	Didactic session two Institution two	Didactic session three Institution three	Didactic session four Institution four
Lecture topics	Personal statement	Presentation skills	How to choose the right program for you	Application and interviewing process
	Chest pain	Altered mental status	Toxicology overview	Headache
	Shortness of breath	Abdominal pain	Shock and sepsis	Gynecologic and urologic emergencies
	Radiographs	Electrocardiogram introduction	Vaginal bleeding	Endocrine and electrolytes
	Social emergency medicine	Ultrasound basics	Advanced trauma life support	Advanced cardiac life support, basic life support

Session:
Presenter:

Please rate the effectiveness of the following session in accomplishing its learning objectives on a scale from 1 (not effective) to 5 (very effective)

Questions	These sessions helped me learn the approach to core emergency medicine topics (abdominal pain, chest pain, headache, etc.) more so than I would have been able to do on my own.	These sessions helped me learn key skills for excelling in an emergency medicine rotation including oral presentations, EKG interpretation, x-ray interpretation and ultrasound, more so than I would have been able to do on my own.	These sessions helped me learn about the process of applying to and selecting an EM residency program.	These sessions allowed me to connect with faculty and resident mentors to learn more about the field of emergency medicine.
Response Options	Strongly agree, agree, neutral, disagree, strongly disagree	Strongly agree, agree, neutral, disagree, strongly disagree	Strongly agree, agree, neutral, disagree, strongly disagree	Strongly agree, agree, neutral, disagree, strongly disagree
Response				

Which parts of the curriculum were of most value to you?

Which parts of the curriculum could be improved?

Figure 1. Evaluation form sent to students after each session.

connect with faculty and resident mentors to learn more about the field of EM. Of the five students, 80% and 20% reported strongly agreeing and agreeing, respectively, that these sessions helped them learn about the process of applying to and selecting an EM residency program.

Narrative feedback, such as the quotes below, from students highlighted the value of meeting with faculty and residents from different programs. from going through cases in real time.

Meeting the faculty and program directors at various EM programs really was the highlight of the curriculum. It was great to get an inside look at each program and learn more about their culture, approach, and the people there.

I really enjoyed hearing the residents’ perspective on how to navigate the application process.

Narrative feedback, such as the quotes below, also emphasized the value of the curriculum’s interactive nature and how traditionally in-person topics were effectively adapted for virtual learning.

My favorite part was participating in real-time cases. Being involved as the case unfolded felt like hands-on practice.

It was incredible to have the mechanisms of ultrasound explained in such detail. Breaking it down to the basics really helped me understand ultrasound for the first time.

REFLECTIONS AND LESSONS LEARNED

Engagement of the Home Institution

Successful implementation required active engagement from HUCOM, specifically the clerkship director and administrative staff, who served as lead contacts. Control over rotation scheduling was essential to ensure all students were fully engaged in the sessions. In addition, as participating institutions used various online platforms to communicate and disseminate curricula materials, such as *Tintinalli’s Emergency Medicine*, with their students, it was necessary to have HUCOM manage a central communications- and video-conferencing platform that was accessible to all lecturing institutions and participating students.

Engagement of Collaborating Institutions

Recruiting faculty and residents for each institution's week was challenging, but having representatives with strong connections in medical education made a significant difference. These relationships allowed them to quickly and effectively recruit lecturers, leveraging their networks to secure individuals who were both willing and enthusiastic to participate. This highlights the value of having institutional leads with established ties to their educational infrastructure, streamlining the recruitment process.

Collaborative Power

The success of this project involved a high degree of trust as many of the institutional representatives had not worked together. To develop this trust, we followed the framework of engaging, listening, framing, envisioning, and committing.¹⁵ The power of this program is truly in the collective rather than the individual. While students could learn about atrial fibrillation from one institution, the real learning occurs when they see the collaboration, get a sense of the scope of EM as a professional field, and are able to interact with varied institutions that have different approaches to teaching and the practice of medicine.

Challenges with Small Student Cohorts

Unlike traditional EM rotations that attract students from across the country, our program had a small cohort comprised solely of HUCOM students, as there was no affiliated residency. This small group size meant that if one student missed a session due to interviews, illness, or other reasons, it noticeably impacted the learning environment, limiting group dynamics and peer-to-peer learning.

Program Limitations and Adaptations

Virtual learning posed challenges for teaching interactive skills such as ultrasound. We addressed this by incorporating case-based learning with curated image libraries and real-time feedback. To further enhance the learning experience, future iterations should explore the integration of ultrasound simulation software to better mimic hands-on scenarios.

Scalability and Expansion

Although initially designed for HUCOM students, this model could be expanded to other medical schools without academic EDs, especially those with a high proportion of URiM students. With the opening of additional HBCU medical schools, there is an even greater need for programs that increase access to EM education.

Limitations

Study limitations include the small sample size as well as lack of a comparison group. Future analyses will address

these limitations and include evaluation of match outcomes as well as other learner-centered targets such as performance in Standardized Letters of Evaluation or subsequent rotations and intern year performance.

Address for Correspondence: Cortlyn Brown, MD, MCSO, Atrium Health Carolinas, Department of Emergency Medicine, 1000 Blythe Blvd., Charlotte, NC 28203. Email: Cortlyn.Brown@atriumhealth.org

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. No author has professional or financial relationships with any companies that are relevant to this study. There are no conflicts of interest or sources of funding to declare.

Copyright: © 2025 Brown et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. Komaromy M, Grumbach K, Drake M, et al. The role of Black and Hispanic physicians in providing health care for underserved populations. *N Engl J Med.* 1996;334(20):1305–10.
2. Marrast LM, Zallman L, Woolhandler S, et al. Minority physicians' role in the care of underserved patients: diversifying the physician workforce may be key in addressing health disparities. *JAMA Intern Med.* 2014;174(2):289.
3. Gomez LE and Bernet P. Diversity improves performance and outcomes. *J Natl Med Assoc.* 2019;111(4):383–92.
4. Saha S, Komaromy M, Koepsell TD, et al. Patient-physician racial concordance and the perceived quality and use of health care. *Arch Intern Med.* 1999;159(9):997.
5. Takeshita J, Wang S, Loren AW, et al. Association of racial/ethnic and gender concordance between patients and physicians with patient experience ratings. *JAMA Netw Open.* 2020;3(11):e2024583.
6. Shen MJ, Peterson EB, Costas-Muñiz R, et al. The effects of race and racial concordance on patient-physician communication: a systematic review of the literature. *J Racial Ethn Health Disparities.* 2018;5(1):117–40.
7. Garrick JF, Perez B, Anaebere TC, et al. The diversity snowball effect: the quest to increase diversity in emergency medicine. A case study of Highland's emergency medicine residency program. *Ann Emerg Med.* 2019;73(6):639–47.
8. Boatright D, Tunson J, Caruso E, et al. The impact of the 2008 Council of Emergency Residency Directors (CORD) panel on emergency medicine resident diversity. *J Emerg Med.* 2016;51(5):576–83.
9. Mabeza RM, Christophers B, Ederaine SA, et al. Interventions associated with racial and ethnic diversity in US graduate medical education: a scoping review. *JAMA Netw Open.* 2023;6(1):e2249335.
10. Martinez J, Mieres JH, Roswell RO, URiM (underrepresented in medicine) learner and faculty mentoring. In: Fornari A and Shah DT

- (Eds.), *Mentoring in Health Professions Education (34–43)*. *IAMSE manuals*. Cham, Switzerland: Springer International Publishing, 2021.
11. De La Cruz MSD, Sairenji T, Stumbar SE, et al. Curricular recommendations for a national family medicine subinternship: a qualitative analysis from multiple stakeholders. *Fam Med*. 2021;53(10):835–42.
 12. Goines J, Iledare E, Ander D, et al. A model partnership: mentoring underrepresented students in medicine (URiM) in emergency medicine. *West J Emerg Med*. 2021;22(2):213–7.
 13. Sweet L and Palazzi D. Application of Kern's six-step approach to curriculum development by global health residents. *Educ Health*. 2015;28(2):138.
 14. Thomas PA, Kern DE, Hughes MT, et al. (Eds.). *Curriculum Development for Medical Education: a Six-step Approach*. 3rd Ed. Baltimore, Maryland: Johns Hopkins University Press, 2016.
 15. Maister D, Green C, Gaiford R. *The Trusted Advisor*. New York New York: Simon & Schuster, 20th Anniversary Ed., 2021.



SAEM 25

**PHILADELPHIA MARRIOTT DOWNTOWN • PHILADELPHIA, PA
MAY 13-16, 2025**

**Society For Academic Emergency Medicine | 1111 East Touhy Avenue, Suite 540 | Des Plaines, IL 60018
SAEM.org | Ph: 847-257-7236 | exhibitors@saem.org**

Call for Section Editors


- Clinical Practice
- Education
- EMS
- Infectious Disease
- Toxicology

WestJEM Integrating Emergency Care
with Population Health
westjem.org

Send CV and letter of interest to
Editor@WestJEM.org

WestJEM Integrating Emergency Care
with Population Health

Call for Reviewers!



Please send your CV and letter of interest to
editor@westjem.org

CPC *EM* Clinical
Practice
& Cases

**JOIN
CAL/AAEM!**



CALAAEM

CALIFORNIA CHAPTER DIVISION OF THE
AMERICAN ACADEMY OF EMERGENCY MEDICINE

*Championing
individual physician rights
and workplace fairness*

BENEFITS

- *Western Journal of Emergency Medicine* Subscription
- CAL/AAEM News Service email updates
- Free and discounted registration to CAL/AAEM events
- And more!

CAL/AAEM NEWS SERVICE

- Healthcare industry news
- Public policy
- Government issues
- Legal cases and court decisions

In collaboration with our official journal

WestJEM Integrating Emergency Care
with Population Health



Join the CAL/AAEM Facebook Group to stay up-to-date:
www.facebook.com/groups/calaaem

www.aaem.org/calaaem



CALIFORNIA ACEP

AMERICAN COLLEGE OF EMERGENCY PHYSICIANS

MEDI-CAL RATE INCREASE

**CALIFORNIA ACEP SECURED
\$200M ANNUALLY TO
INCREASE MEDI-CAL
REIMBURSEMENT FOR
EMERGENCY PHYSICIANS!**

**EMERGENCY MEDICINE IS THE ONLY
SPECIALTY WITH ITS OWN BUDGET
ALLOCATION.**

@CALIFORNIAACEP

www.californiaacep.org