

# Acute Care of Patients with Moderate Respiratory Distress: Recommendations from an American College of Emergency Physicians Expert Panel

Christopher W. Baugh, MD, MBA\*

Jim F. Neuenschwander, MD†

Jesslyn Lenox, MHA, RRT-NPS, AE-C‡

Jennifer Hoh, PharmD, BCCCP§

Kara Ward, MD||#

Sara Muramoto, MD¶

John Casey, DO, MA†

Antonio Anzueto, MD\*\*

Hajirah Ishaq, DO, MS††

Jared Mount, BS‡‡

Peter M. DeBlieux, MD§§

\*Brigham and Women's Hospital, Department of Emergency Medicine, Boston, Massachusetts

†OhioHealth Doctors Hospital, Department of Emergency Medicine, Columbus, Ohio

‡South Shore Hospital, Pulmonary Function Lab and Neurodiagnostics, Weymouth, Massachusetts

§University of Louisiana at Monroe, Department of Emergency Medicine, Monroe, Louisiana

||Tammany Health System, Division of Pulmonary & Critical Care, St Covington, Louisiana

#Robotic Critical Care Services

¶Louisiana State University's Spirit of Charity Emergency Medicine Residency Program, New Orleans, Louisiana

\*\*UT Health San Antonio, South Texas Veterans Health Care System, Division of Pulmonary Medicine/ Critical Care, San Antonio, Texas

††Indiana University School of Medicine, Department of Emergency Medicine, Indianapolis, Indiana

‡‡Ohio University Heritage College of Osteopathic Medicine, Department of Medical Education, Athens, Ohio

§§University Medical Center New Orleans, Department of Emergency Medicine, New Orleans, Louisiana

*Section Editor:* Tom Benzoni

Submission history: Submitted February 18, 2025; Revision received May 13, 2025; Accepted May 23, 2025

Electronically published September 27, 2025

Full text available through open access at [http://escholarship.org/uc/uciem\\_westjem](http://escholarship.org/uc/uciem_westjem)

DOI 10.5811/westjem.43539

**Introduction:** Patients with respiratory distress are frequently encountered in the emergency department (ED). Efforts to assess, initiate treatments, and stabilize these patients require a systematic and rapid response. Emergency physicians need a comprehensive and efficient approach for evaluating, treating, and managing patients presenting to the ED with moderate respiratory distress.

**Methods:** The American College of Emergency Physicians convened an expert panel of academic and community emergency physicians, critical care specialists, respiratory therapists, hospitalists, and pharmacists to develop and subsequently disseminate consensus recommendations regarding the diagnosis and treatment of patients with moderate respiratory distress presenting to the ED.

**Results:** A digital tool using a consensus-based framework was developed to aid emergency clinicians in diagnosing and caring for patients with moderate respiratory distress. The tool can be employed at each step in the diagnostic and treatment process.

**Conclusion:** The evidence-based tool is a practical and freely available bedside instrument for emergency clinicians to diagnose and treat patients with moderate respiratory distress. Further studies are needed to examine the effectiveness of this approach. [West J Emerg Med. 2025;26(5)1485–1494.]

## INTRODUCTION

Respiratory distress or shortness of breath represent the third most common reason for visiting the emergency department (ED) in the United States, with 5,918,000 visits in 2021.<sup>1</sup> Shortness of breath is a common manifestation of a wide array of primary pulmonary and respiratory tract illnesses, as well as other potentially life-threatening conditions, such as acute coronary syndrome, pulmonary embolism, cardiac tamponade, severe anemia, toxic metabolic disturbances, and many others.<sup>2</sup> Emergency physicians must manage these patients with limited time and information. They also initiate rapid assessment, diagnosis, and treatment to stabilize and begin reversing life-threatening pathophysiology amid increased stressors such as ED boarding and patients with higher morbidity.<sup>3</sup> Frequent reassessment, adjustments to new data, and observed patient response to therapies are the mainstays of ED care for this patient population.

For emergency physicians faced with the challenge of caring for patients with respiratory distress, many resources, such as specialty society guidelines, are available for disease-specific assessment and treatment.<sup>4</sup> Few resources are readily available for the initial care of patients with undifferentiated respiratory distress, especially those with moderate symptoms and elevated risk.<sup>5-7</sup> Often, immediate aggressive maneuvers, such as rapid sequence intubation, are not indicated. Developing and disseminating digital tools easily accessed via smartphone or desktop computer, and explicitly designed to support real-time clinical care, is needed.<sup>8</sup> Lastly, many authoritative guidelines are rarely updated due to rigorous and time-consuming processes for evidence review and consensus generation. This timing results in a delay between the time when innovative and emerging approaches, such as high-flow nasal cannula, are indicated for use and when they are subsequently incorporated into expert recommendations.

Expert panels can be convened to address challenges in clinical practice that students, physicians, nurses, physician assistants, and nurse practitioners may face and provide consensus recommendations. The initial assessment and optimal diagnostic and therapeutic approach to patients with respiratory distress in the ED can be complex and time sensitive. The American College of Emergency Physicians (ACEP) expert panel is a mechanism to address the diagnosis and management of moderate respiratory distress in the ED by providing a practical, consensus-based guide via an online tool and smartphone app at the point of care.

## METHODS

### Study Design

An expert panel to develop a consensus-based decision tool for emergency clinicians was convened by ACEP. Selected panel members conducted a literature review focusing on terms such as “respiratory distress” to examine recent ( $\leq 5$  years) publications and specialty-society

### *Population Health Research Capsule*

What do we already know about this issue?  
*Patients frequently present to the ED with undifferentiated moderate respiratory distress.*

What was the research question?  
*We convened an expert panel and developed a digital tool to assist clinicians' bedside assessment of respiratory distress in an efficient and evidence-based manner.*

What was the major finding of the study?  
*We emphasize a rapid yet thorough initial history and exam to guide diagnosis, initiate targeted treatments, and select the most appropriate disposition for this target patient population.*

How does this improve population health?  
*As emergency clinicians face challenges to assess and treat complex patients, digital tools like the one we developed and implemented can help improve care for higher risk populations.*

guidelines involving the approach to respiratory distress in the ED. The panel sought more recent literature to reflect the current and emerging science related to the target population. When possible, definitions and recommendations reflect study criteria and treatment processes. Otherwise, the expert panel sought to standardize definitions and processes. The panel participated in four virtual rounds of questioning to make recommendations using a modified Delphi rounds process to reach a consensus.<sup>9</sup> Statements were discussed extensively during the virtual meetings and then serially modified based on discussions. The final expert panel recommendations represented consensus (defined as majority agreement with final adjudication by co-chairs) and majority opinions. This project relied on publicly available sources and was framed as a quality improvement project exempt from institutional review board oversight.

The definition of moderate respiratory distress is intricate and not easily defined. Our recommendations defined moderate respiratory distress as “a patient who requires more than basic respiratory care or is at risk of deterioration.” For example, this definition would include a patient with a respiratory rate  $> 24$  breaths per minute, with or without hypoxia, and with the use of accessory muscles of respiration.

## Setting

The multidisciplinary expert panel, which was diverse in geography and practice settings, was convened virtually between January–July 2024.

## Selection of Participants

The American College of Emergency Physicians selected the expert panel; the chosen co-chairs Jim F. Neuenschwander and Peter M. DeBlieux then selected additional panelists. Criteria for selection included individuals with direct personal clinical experience in treating patients with respiratory distress in the ED, prior contributions to ACEP expert panels and/or academic contributions, and diverse clinical practice settings within the group. Clinical backgrounds included emergency medicine, critical care, respiratory therapy, pharmacy, and hospital medicine. The final panel consisted of three academic emergency physicians, two community emergency physicians, one ED pharmacist, one respiratory therapist, and one hospitalist, with panelists representing a broad geographic distribution across the US (eg, Southwest, South, Midwest, Northeast).

## RESULTS

The ACEP moderate respiratory distress emergency medicine point of care (emPOC) tool was developed to communicate the main steps in the evaluation, treatment, and disposition of patients presenting to the ED with moderate respiratory distress. The tool is available on the [ACEP website](#) and as a smartphone app for [Apple](#) and [Android](#) devices.

## History and Physical Exam

Care begins with the initial assessment, which includes obtaining a history of present illness and performing a physical examination. The history may be obtained from the patient, emergency medical services, family, or other historians to assist with differential diagnosis, initial testing, and treatment strategy. Table 1a outlines the key elements of the history that should be obtained for patients presenting to the ED with moderate respiratory distress. In parallel with obtaining the patient's history, a physical examination should begin immediately as the patient is assessed. We outline the key exam elements in Table 1b.

## Testing

After the initial evaluation, the differential diagnosis generated from the encounter will drive a diagnostic strategy. Testing falls into three broad categories: laboratory studies; imaging studies; and other testing. We outline the key diagnostic considerations in Table 2, which may be modified by local resource availability and the evolution of the differential diagnosis as more information is gathered over the patient's ED course.

**Table 1a.** History elements for patients with moderate respiratory distress.

- 
- Dyspnea at rest and with ambulation
  - Chest pain quality, including pleuritic pain
  - Increased sputum production
  - Fever
  - Heart failure
  - Orthopnea
  - Paroxysmal nocturnal dyspnea
  - Cough
  - Peripheral edema
  - Fatigue
  - Altered mental status
  - Weakness
  - Smoking/vaping use
  - Chronic lung disease (e.g., chronic obstructive pulmonary disease, interstitial lung disease, pulmonary fibrosis)
  - Anemia
  - Asthma
  - Increased sputum/color/production/volume
  - Current oxygen requirement
  - Prior intubation or known difficult airway
  - Recent travel (e.g., elevation, exposure, pulmonary embolism)
  - Known infectious contacts
  - Prior personal or family history of venous thromboembolism
  - Terminal diagnosis/condition
  - Exogenous estrogen use
  - Hemoptysis
  - Hospitalizations
  - Trauma/surgery (e.g., fat emboli, rib fractures, pulmonary contusion, flail chest)
  - Dysphagia/severe reflux
  - Environmental exposure (e.g., carbon monoxide, asbestos)
  - Congenital anomalies (e.g., Marfan, cystic fibrosis, Ehlers-Danlos)
  - Cancer
  - Inflammatory (e.g., lupus, rheumatoid arthritis)
  - Substance use disorder
  - Goals of care
  - Pain
  - Anxiety
  - Immunosuppressed state
- 

## Treatment

Treatments should be targeted at the clinical scene, addressing the underlying pathophysiology driving the patient's presentation and helping support their work of breathing to avoid clinical decompensation. Initial maneuvers to re-position the patient, suction the airway, or other immediate interventions may be helpful to improve the

**Table 1b.** Physical examination elements for the patient with moderate respiratory distress.

- Vitals trending, including prehospital
- Tachypnea
- Tachycardia
- Hypotension
- Upper airway exam
- Voice quality
- Abnormal lung exam (e.g., wheezing, rhonchi, crackles)
- Hypoxia
- Tripoding or accessory muscle use
- Mental status
- Abnormal skin exam (e.g., pallor or discoloration, urticarial rash)
- Swallow
- Cough quality
- Grunting, nasal flaring, or abdominal contractions
- Diaphoresis
- Irregular breathing pattern
- Crepitus
- Extremity edema
- Screening for difficult airway assessment and management: (<https://www.aliem.com/mnemonics-for-difficult-airway/>)

History and physical exam elements for patients presenting to the ED with moderate respiratory distress.

patient's work of breathing. Frequent reassessments are a critical aspect of the care of this patient population. The goal is to choose the best intervention to match the patient's current and anticipated status with frequent evaluation of the need for respiratory support escalation or de-escalation.<sup>10</sup> As a result, the clinician should proceed with continuous information gathering and optimization of a differential diagnosis, work of breathing, serial vital signs, serial mental status exams, repeat imaging (eg, point-of-care ultrasound), lab result analysis (eg, serial blood gasses), disposition trajectory, evolving goals of care, and delivery of more targeted treatments as the diagnosis becomes clearer.

When using a high-flow nasal cannula, the ratio of oxygenation (ROX) index predicts clinical decompensation and the need for intubation.<sup>11-13</sup> The ROX index is the SpO<sub>2</sub> (peripheral oxygen saturation)/FIO<sub>2</sub> (fraction of inspired oxygen) (%) ratio to respiratory rate (breaths/minute). Values  $\geq 4.88$  measured at 2, 6, or 12 hours after high-flow nasal cannula initiation are associated with a lower intubation risk, whereas values  $< 3.85$  indicate a high risk of high-flow nasal cannula failure and intubation consideration. Intermediate ROX values should be reassessed serially, 1-2 hours later. We illustrate an algorithm for treating and reassessing patients with moderate respiratory distress in Figures 1a and 1b for the adult and pediatric populations, respectively.<sup>14,15</sup>

Various respiratory therapies are available in the ED and should match the patient's needs. Table 3 highlights the key similarities and differences between commonly available respiratory therapies, including nasal cannula, facemask, high-flow nasal cannula, non-invasive ventilation, intubation, and surgical airways. We provide additional detail for each treatment in Figure 2. For some patients, intubation and mechanical ventilation avoidance may be an important consideration, as intubation may increase the risk of peri-intubation complications. These patients may require special ventilation strategies and include examples such as status asthmaticus, severe metabolic acidosis, and untreated pneumothorax.

Medications are an essential component of treatment for most patients in the ED with moderate respiratory distress. Given the broad and variable causes of respiratory distress, we outline key pharmacologic treatments for selected common conditions in Supplement Table 1. This resource is not intended to be exhaustive, and once a specific diagnosis is strongly suspected or confirmed, we recommend that clinicians tailor treatments to best treat specific conditions.

### Disposition

Disposition decisions should be made based on the severity of symptoms, concurrent comorbidities, and the social drivers of health that impact a patient's ability to adhere to a care plan. Patients from the ED will be discharged home, admitted to the hospital (including inpatient or observation status), or admitted to the intensive care unit. Patients who want to focus on symptom control and quality of life can be directed to palliative care or hospice.

Patients with mild-to-moderate symptoms who respond well to initial therapy in the ED and do not require new or additional supplemental oxygen may be suitable for discharge home. Patients who are being considered for discharge home should meet the following additional requirements: able to ambulate without a new significant drop in oxygen saturation or return of presenting symptoms; able to obtain prescribed medications promptly; able to express an understanding of their medications and how to properly take them, patient or caregiver can perform discharge instruction "teach back" and demonstrate proper use of new and existing therapies; able to express understanding of return precautions (ie, worrisome signs and symptoms that warrant a return to the ED), and able to follow up with an outpatient clinic within 1-2 weeks of the ED visit. Discharge instructions should include a written discharge plan that contains a list of newly prescribed medications, including their purpose and directions (eg, frequency and duration), a list of all home medications, including any adjustments or changes, signs or symptoms that would warrant returning to the ED, and clear directions to follow up with a primary care physician or specialist in a specified time frame. For patients who smoke or live with

**Table 2.** Diagnostic approach to the patient with moderate respiratory distress.

## General Labs

Complete blood count (CBC)

anemia, pneumonia/infections

Comprehensive metabolic panel (CMP)

ketoacidosis (diabetic ketoacidosis [DKA]/starvation ketoacidosis [SKA]/alcoholic ketoacidosis [AKA]), pulmonary edema (cardiorenal effects), electrolytes (cardiac arrhythmia), metabolic acidosis

Point of care blood gas

## Targeted Labs

Blood gas: use pulse ox and bicarb on basic metabolic panel (BMP)

Co-oximetry

carbon monoxide, methemoglobinemia

Venous blood gas (VBG)

chronic obstructive pulmonary disease (COPD), asthma, respiratory failure

Brain natriuretic peptide (BNP)

heart failure, cardiomyopathy, acute coronary syndrome (ACS)

International normalized ratio (INR) with a history of anticoagulation or coagulopathies

Anti-factor Xa test if actively bleeding

Troponin

acute coronary syndrome (ACS)/pulmonary embolism (PE)/venous thromboembolism (VTE)

D-dimer

PE rule out

Lactate

seizure, sepsis

Procalcitonin

pneumonia or other infection

Blood cultures

Nasal swab for viral panel

C-reactive protein (CRP)

chronic lung disease

Toxicology

salicylate

alcohol and urine tox

## Targeted imaging/studies

Chest x-ray (CXR)

pneumonia, pleural effusion, enlarged cardiac silhouette, widened mediastinum, pneumomediastinum, rib fractures, pneumothorax, subcutaneous air, acute respiratory distress syndrome (ARDS), mass, atelectasis, small inspiratory lung volumes (neuromuscular disease)

Expiratory films for focal air trapping/unilateral wheezing

foreign body, mucous plug

Computed tomography chest

with contrast

concern for mediastinal mass or pleural metastases

without contrast


pneumonia, ground glass opacities, eval pulmonary nodule/mass seen on x-ray, pulmonary contusions, ARDS, pulmonary edema, rib fractures

Diagnostic testing considerations for patients presenting to the emergency department with moderate respiratory distress.

**Table 2.** Continued.

CT angiogram (CTA) Chest
PE aortic dissection, trauma
Ventilation/Perfusion (V/Q) scan - hard to interpret with existing pulmonary disease/abnormal CXR
PE
Point of care ultrasound
bedside: pneumonia, pneumothorax, pleural or pericardial effusion, pulmonary edema, cardiac tamponade, right heart strain (PE, right heart failure), left ventricular (LV) function, ability to tolerate fluids, inferior vena cava (IVC), lung windows
Consider Ancillary Testing:
End-tidal CO <sub>2</sub> (ETCO <sub>2</sub> )
waveform analysis for overdose, asthma, and COPD
Electrocardiogram (EKG)
arrhythmia, ACS
Negative inspiratory force (NIF)
neuromuscular disease
Physical exam maneuvers
fluid status
passive leg raise-fluid responsiveness

Diagnostic testing considerations for patients presenting to the emergency department with moderate respiratory distress. ACS, acute coronary syndrome; CO<sub>2</sub>, carbon dioxide; COPD, chronic obstructive pulmonary disease.



**Figure 1. A.** Adult moderate respiratory distress algorithm. **B.** Pediatric moderate respiratory distress algorithm. <sup>†</sup>Criteria for immediate intubation include inability to protect the airway, anticipated clinical course, and failure of oxygenation or ventilation despite other treatments (e.g., supplemental oxygen, non-invasive ventilation, or not a candidate for those therapies). <sup>‡</sup>The extent and reliability of ROX index validation in children remain limited and variable. Patient care algorithm for adult and pediatric patients presenting to the emergency department with moderate respiratory distress.

Lpm, liters per minute; RR, respiratory rate; NIV, noninvasive ventilation; WOB, work of breathing; HFNC, high-flow nasal cannula; SpO<sub>2</sub>, peripheral oxygen saturation; ROX, ratio of oxygenation; PEEP, positive-end expiratory pressure; FiO<sub>2</sub>, fraction of inspired oxygen. **\*\***Figure 1B adapted from: Mosier JM, Tidswell M, Wang HE. Noninvasive respiratory support in the emergency department: Controversies and state-of-the-art recommendations. *J Am Coll Emerg Physicians Open.* 2024;5(2):e13118. doi:10.1002/emp2.13118 and Kwon JW. High-flow nasal cannula oxygen therapy in children: a clinical review. *Clin Exp Pediatr.* 2020;63(1):3-7. doi:10.3345/kjp.2019.00626.

**Table 3.** Comparison of respiratory therapies.

NC, mask, NRB	HFNC	NIV	ETT/Surgical neck airway
FiO <sub>2</sub> estimated		FiO <sub>2</sub> controlled	
Well-tolerated	Tolerated	Uncomfortable	Medicated
Inexpensive, simple	Intermediate cost, Increased resource intensity		Expensive, resource intensive
No PEEP, No Pressure Support	PEEP (Low level)		PEEP +/- Pressure Support
	Cooperation Not Needed	Cooperation Needed†	Not Needed
	Patient clears secretions	Assisted secretion clearance	

†Patient participation is required.

Respiratory therapy options for patients presenting to the emergency department with moderate respiratory distress.

NC, nasal cannula; Mask, facemask; NRB, nonrebreather; HFNC, high-flow nasal cannula; NIV, non-invasive ventilation; ETT, endotracheal tube; FiO<sub>2</sub>, fraction of inspired oxygen; PEEP, positive end-expiratory pressure.

\*\*Adapted from Baker K, Greaves T, Fraser JF. How to use humidified high-flow nasal cannula in breathless adults in the emergency department. *Emerg Med Australas*. 2019 Oct;31(5):863-868. doi: 10.1111/1742-6723.13372. Epub 2019 Aug 6. PMID: 31389171.

someone who smokes, smoking cessation information should be given with the discharge paperwork and discussed at the bedside before discharge.

Patients should be hospitalized if they have a new supplemental oxygen requirement. This includes those who have an increased need for supplemental oxygen above their baseline, advanced lung disease, and/or multiple comorbidities, and those who are otherwise unlikely to do well if discharged home. Additionally, those with a new concurrent condition, such as pneumonia, arrhythmias, heart failure, and sepsis, or who have persistent symptoms despite initial ED treatment and require constant monitoring or titration of ventilatory support, should be hospitalized. Finally, those not meeting the criteria for a safe discharge home, including those with social drivers of health likely to disrupt the discharge plan, should remain until a safe plan is identified. Many hospitals have disease-specific protocols that help emergency clinicians determine patient eligibility for observation vs inpatient care, including using ED observation units and home hospital services. We recommend that clinicians reference their local protocols to help direct patients toward the most appropriate site of care in their institution. Example ED observation protocols can be found on [ACEP's Observation Services Toolkit](#) page.

Based on institution-specific capabilities and policies, consider intensive care unit admission for patients with moderate-to-severe symptoms who require escalating respiratory support, are hemodynamically unstable, are intubated, have severe acid-base disturbances, have a new altered mental status, or other symptoms that necessitate close monitoring. Complex patients who may exceed local capabilities or are better served at a higher resource facility require an interfacility transfer. Finally, patients who want to focus on symptom control and quality of life primarily can be directed to palliative care or hospice.









## DISCUSSION

In this paper, we describe our process and consensus

recommendations generated by an ACEP-convened expert panel focused on ED care of patients presenting with moderate respiratory distress. Our recommendations emphasize a rapid yet thorough initial history and exam to create and narrow a differential diagnosis, initiate treatments, and select the most appropriate disposition for this target patient population. The framework is presented as a digital tool, available via smartphone app and desktop computer and designed for real-time accessibility to assist clinicians at the bedside for this challenging patient population.

Prior investigations and tools are overwhelmingly targeted at a specific diagnostic tool or therapy for a disease process. For example, evidence abounds for the roles of point-of-care ultrasound and high-flow nasal cannula for the initial evaluation and treatment of patients with undifferentiated respiratory distress.<sup>16-19</sup> Disease-specific guidance for respiratory illnesses such as chronic obstructive pulmonary disease, heart failure, pulmonary embolism, and pneumonia is also available.<sup>4,20-22</sup> Alternatively, other resources with a broad approach have not been designed as a tool for real-time clinical support.<sup>2,23</sup> As a result, our novel approach of combining broad patient assessment and treatment guidance with a format optimized for bedside use is novel and designed for maximal clinical impact. This combination best reflects the real-world scenario of the initial ED encounter, where the diagnosis is not yet confirmed, and the support of a digital tool may be most helpful.

The role of electronic clinical decision-support tools is rapidly evolving. Following the framework of the five “rights” of clinical decision-support, our moderate respiratory distress tool aims to provide the right information to the right people in the right channel, format, and time in the workflow.<sup>24</sup> Evidence is increasing for the use of interactive digital tools at the bedside for high-acuity scenarios, such as pediatric cardiac arrest.<sup>25</sup> Enabling prompts for clinicians to gather key information and act quickly with evidence-based interventions has the promise of improving care for vulnerable patients,

<p><b>Nasal Cannula (NC)</b></p> <p>Pros: Comfortable, enables communication, eating &amp; drinking. Well tolerated. Easily travels</p> <p>Cons: Can be drying without humidification. Cannot provide a lot of oxygen</p>	
<p><b>Indication:</b> Low flow, Low <math>FI_{O_2}</math> %</p> <p><b>Flow (Lpm):</b> 0.1 → 6</p> <p><b>Oxygen %:</b> 24-40 (v)</p> <p><i>Fixed (f) or Variable (v)</i></p> <p><b>Delivery Interface:</b> Nasal Cannula</p> <p><b>Humidification:</b> Possible</p>	
<p><b>Simple Face Mask</b></p> <p>Pros: Enables communication, easily removed for eating &amp; drinking. Easily travels</p> <p>Cons: Can be drying. Cannot provide a lot of oxygen. Dangerous at flows &lt;6 Lpm</p>	
<p><b>Indication:</b> Med flow, Low <math>FI_{O_2}</math> %</p> <p><b>Flow (Lpm):</b> 6 → 10</p> <p><b>Oxygen %:</b> 35-60 (v)</p> <p><i>Fixed (f) or Variable (v)</i></p> <p><b>Delivery Interface:</b> Face Mask</p> <p><b>Humidification:</b> No</p>	
<p><b>Aerosol Mask / Face Tent / Trach Mask</b></p> <p>Pros: Generally well tolerated. Enables humidification &amp; medication delivery</p> <p>Cons: Cannot provide a lot of oxygen. Dangerous at flows &lt;6 Lpm</p>	
<p><b>Indication:</b> Med Flow, Med Delivery</p> <p><b>Flow (Lpm):</b> 6 → 10</p> <p><b>Oxygen %:</b> 35-60 (v)</p> <p><i>Fixed (f) or Variable (v)</i></p> <p><b>Delivery Interface:</b> Face Mask</p> <p><b>Humidification:</b> Yes</p>	
<p><b>Non-invasive Ventilation</b></p> <p>Pros: Fixed <math>FI_{O_2}</math>%. Enables single or bi-levels of pressure delivery.</p> <p>Cons: Requires a seal to deliver pressure, can cause skin damage. Inhibits communication. Can be very drying.</p>	
<p><b>Indication:</b> OSA, Augmentation of Oxygenation Support (CPAP), Ventilation Support (BIPAP), RR possible</p> <p><b>Flow (Lpm):</b> _____</p> <p><b>Oxygen %:</b> 21-100 (f)</p> <p><i>Fixed (f) or Variable (v)</i></p> <p><b>Delivery Interface:</b> Face or Nasal Mask, Nasal Pillows</p> <p><b>Humidification:</b> Possible</p>	
<p><b>High Flow Nasal Cannula</b></p> <p>Pros: Optimal humidification control, enables communication, easily removed for eating &amp; drinking. Travels</p> <p>Cons: Unable to deliver high levels of pressure.</p>	
<p><b>Indication:</b> Higher Flow &amp; <math>FI_{O_2}</math> %, Augmentation of Oxygenation Support (some +pressure), Ventilation Support (Dead space washout)</p> <p><b>Flow (Lpm):</b> 2 → 70</p> <p><b>Oxygen %:</b> 21-100 (f)</p> <p><i>Fixed (f) or Variable (v)</i></p> <p><b>Delivery Interface:</b> Large Bore Nasal Cannula</p> <p><b>Humidification:</b> Yes</p>	
<p><b>Venturi Mask</b></p> <p>Pros: Generally well tolerated. Uniquely allows for precise oxygen titration.</p> <p>Cons: Can be drying. Cannot provide high amounts of oxygen. Dangerous at flows &lt;6 Lpm</p>	
<p><b>Indication:</b> Precise <math>O_2</math> titration</p> <p><b>Flow (Lpm):</b> 6 → 10</p> <p><b>Oxygen %:</b> 24-60</p> <p><i>Fixed (f) or Variable (v)</i></p> <p><b>Delivery Interface:</b> Aerosol Mask</p> <p><b>Humidification:</b> No</p>	
<p><b>"Higher Flow" NC</b></p> <p>Pros: Comfortable, enables communication, eating &amp; drinking. Well tolerated. Easily travels</p> <p>Cons: Can be very drying. Can not deliver higher than 15 Lpm flow, may be inadequate to meet pt's flow or <math>O_2</math> needs.</p>	
<p><b>Indication:</b> Med Flow &amp; Med <math>FI_{O_2}</math>%</p> <p><b>Flow (Lpm):</b> 2 → 15</p> <p><b>Oxygen %:</b> 24-60 (v)</p> <p><i>Fixed (f) or Variable (v)</i></p> <p><b>Delivery Interface:</b> Nasal Cannula</p> <p><b>Humidification:</b> No</p>	
<p><b>Non-Rebreather</b></p> <p>Pros: Comfortable, enables communication, eating &amp; drinking. Well tolerated. Easily travels</p> <p>Cons: Can be very drying. Can not deliver higher than 15 Lpm flow, may be inadequate to meet pt's flow or <math>O_2</math> needs.</p>	
<p><b>Indication:</b> Higher Flow &amp; <math>FI_{O_2}</math> %</p> <p><b>Flow (Lpm):</b> 15</p> <p><b>Oxygen %:</b> 100 (f)</p> <p><i>Fixed (f) or Variable (v)</i></p> <p><b>Delivery Interface:</b> Face Mask</p> <p><b>Humidification:</b> NO</p>	

**Figure 2.** Summary of respiratory therapy options.

Additional details for respiratory therapy options for patients presenting to the emergency department with moderate respiratory distress. *BIPAP*, bilevel positive airway pressure; *CPAP*, continuous positive airway pressure; *FI<sub>O<sub>2</sub></sub>*, fraction of inspired oxygen; *Lpm*, liters per minute *O<sub>2</sub>*, oxygen; *RR*, respiratory rate, *pt*, patient.

**TABLE 4:** Suggested outcome measures for patients with moderate respiratory distress.

- Emergency department door-to-disposition decision times
- Intensive care unit admission volume and rate
- Hospital length of stay
- Therapy switches (i.e., high flow nasal cannula to NIPPV) and escalations to intubation
- 30-day emergency department revisits and hospital readmissions

NIPPV, non-invasive positive pressure ventilation.

including those with moderate respiratory distress. Future efforts should focus on iterating the content to keep it up to date and improve accessibility, perhaps allowing for electronic health record integration. Such integration may allow for incorporating clinical data feeds (eg, vital signs, laboratory results) and facilitating bedside orders, test result interpretation, and clinical documentation.<sup>26</sup> While these suggestions have been thoughtful in their creation, we recommend outcome measures in Table 4 that can be used as measures of success.

## LIMITATIONS

Convening an expert panel to develop recommendations and create a bedside tool for emergency physicians has several limitations. First, when evidence was lacking, some aspects of the best practices in the ED were derived only from expert panel discussions. A rigorous systematic literature review and grading consistent with Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) standards or Grading of Recommendations Assessment, Development, and Evaluation (GRADE) methodology were not performed, and specialty society treatment guidelines formulated via a structured or systematic review of the evidence are typically restricted to interventions with the highest quality of evidence.<sup>27,28</sup> This expert panel's recommendations are distinct from ACEP clinical policies, governed by the Clinical Policies Committee under a separate process. In addition, outside the relative cost information briefly provided in Table 3, recommendations from this expert panel do not include more detailed economic analyses of the direct and net financial impact of various treatment and disposition (ie, post-ED care) options.

## CONCLUSION

The evidence-based emPOC tool was developed by a multidisciplinary panel of experts convened by ACEP to be a resource for emergency clinicians caring for patients with moderate respiratory distress. Studies examining the functionality of this tool in real-world practice are warranted, as data related to its daily use in the ED setting will allow for improvements and modifications to be rapidly incorporated over time, given the tool's flexibility to be easily updated.

Future studies examining the impact of specific interventions, such as various respiratory therapies initiated during the ED visit vs deferring to the inpatient team, are important to demonstrate the real-world effect of digital tools and further support the benefits of specific interventions on patient outcomes.

*Address for Correspondence:* Christopher W. Baugh, MD, MBA, Brigham and Women's Hospital, 75 Francis Street, Department of Emergency Medicine, Neville House 2nd Floor, 10 Vining Street, Boston, MA 02115. Email: cbaugh@bwh.harvard.edu.

*Conflicts of Interest:* By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. Christopher W. Baugh is a paid speaker for CE Symmetry, Octapharma, and Roche Diagnostics, an investigator for Abbott Laboratories, Roche and Sanofi, an advisory board participant for Roche Diagnostics, Salix Pharmaceuticals, Pfizer Inc., and AstraZeneca, a consultant for Abbott Laboratories, Pfizer Inc., and Roche Diagnostics, and an advisor to Vera Health, Quai, and Lucia Health Guidelines. Jim F. Neuenschwander has received research funding from BridgeSource Medical, CSL Behring, and Siemens Healthineers USA and reports being a speaker and consultant for Abbott Laboratories, AstraZeneca, Janssen (now J&J Innovative Medicine), Thermo Fisher Scientific, Fisher & Paykel Healthcare, and Pfizer Inc.; and is an owner in AsceptiScope, Inc. EO has nothing to disclose. Antonio Anzueto is a consultant for GSK, AstraZeneca, Sanofi/Regeneron, Viatrix, and Pfizer. No other author has professional or financial relationships with any companies that are relevant to this study. There are no other conflicts of interest or sources of funding to declare.

*Copyright:* © 2025 Baugh et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

## REFERENCES

1. Cairns C, Kang, K. National Hospital Ambulatory Medical Care Survey: 2021 emergency department summary tables. Available at: [https://ftp.cdc.gov/pub/Health\\_Statistics/NCHS/Dataset\\_Documentation/NHAMCS/doc21-ed-508.pdf](https://ftp.cdc.gov/pub/Health_Statistics/NCHS/Dataset_Documentation/NHAMCS/doc21-ed-508.pdf). Accessed December 10, 2024.
2. Schwartzstein R. Approach to the patient with dyspnea. *UpToDate*, Connor RF (Ed), Wolters Kluwer. Available at: Approach to the patient with dyspnea - UpToDate. Accessed: March 29, 2025.
3. Oskvarek JJ, Zocchi MS, Black BS, et al. Emergency department volume, severity, and crowding since the onset of the coronavirus disease 2019 pandemic. *Ann Emerg Med*. 2023;82(6):650-60.
4. Metlay JP, Waterer GW, Long AC, et al. Diagnosis and treatment of adults with community-acquired pneumonia. An official clinical practice guideline of the American Thoracic Society and Infectious Diseases Society of America. *Am J Respir Crit Care Med*. 2019;200(7):e45-67.
5. Rochweg B, Brochard L, Elliott MW, et al. Official ERS/ATS clinical

- practice guidelines: noninvasive ventilation for acute respiratory failure. *Eur Respir J*. 2017;50(2):1602426.
6. Rochweg B, Einav S, Chaudhuri D, et al. The role for high flow nasal cannula as a respiratory support strategy in adults: a clinical practice guideline. *Intensive Care Med*. 2020;46(12):2226-37.
  7. Ware LB. Go with the flow: expanding the definition of acute respiratory distress syndrome to include high-flow nasal oxygen. *Am J Respir Crit Care Med*. 2022;205(4):380-2.
  8. Dubin J. How smartphones and apps are changing emergency department care. June 26, 2017. Available at: <https://www.medstarhealth.org/blog/smartphone-medical-apps-emergency-department>. Accessed: February 17, 2025.
  9. Diamond IR, Grant RC, Feldman BM, et al. Defining consensus: a systematic review recommends methodologic criteria for reporting of Delphi studies. *J Clin Epidemiol*. 2014;67(4):401-9.
  10. Baker K, Greaves T, Fraser JF. How to use humidified high-flow nasal cannula in breathless adults in the emergency department. *Emerg Med Australas*. 2019;31(5):863-8.
  11. Roca O, Caralt B, Messika J, et al. An index combining respiratory rate and oxygenation to predict outcome of nasal high-flow therapy. *Am J Respir Crit Care Med*. 2019;199(11):1368-76.
  12. Prakash J, Bhattacharya PK, Yadav AK, et al. ROX index as a good predictor of high flow nasal cannula failure in COVID-19 patients with acute hypoxemic respiratory failure: a systematic review and meta-analysis. *J Crit Care*. 2021;66:102-8.
  13. Melo-Diaz LL, Kieling GA. The ROX index: "propelled" by high-flow nasal cannula therapy during the COVID-19 pandemic into greater applicability in respiratory support. *Can J Respir Ther*. 2022;58:182-4.
  14. Mosier JM, Tidswell M, Wang HE. Noninvasive respiratory support in the emergency department: controversies and state-of-the-art recommendations. *J Am Coll Emerg Physicians Open*. 2024;5(2):e13118.
  15. Kwon JW. High-flow nasal cannula oxygen therapy in children: a clinical review. *Clin Exp Pediatr*. 2020;63(1):3-7.
  16. Wimalasena Y, Kocierz L, Strong D, et al. Lung ultrasound: a useful tool in the assessment of the dyspnoeic patient in the emergency department. Fact or fiction? *Emerg Med J*. 2018;35(4):258-66.
  17. Giorno EPC, Foronda FK, De Paulis M, et al. Point-of-care lung ultrasound score for predicting escalated care in children with respiratory distress. *Am J Emerg Med*. 2023;68:112-8.
  18. Schibler A, Franklin D. Respiratory support for children in the emergency department. *J Paediatr Child Health*. 2016;52(2):192-6.
  19. Long B, Liang SY, Lentz S. High flow nasal cannula for adult acute hypoxemic respiratory failure in the ED setting. *Am J Emerg Med*. 2021;49:352-9.
  20. Agustí A, Celli BR, Criner GJ, et al. Global Initiative for Chronic Obstructive Lung Disease 2023 Report: GOLD Executive Summary. *Eur Respir J*. 2023;61(4):2300239.
  21. Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: Executive Summary: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2022;79(17):1757-80.
  22. Konstantinides SV, Meyer G, Becattini C, et al. 2019 ESC Guidelines for the diagnosis and management of acute pulmonary embolism developed in collaboration with the European Respiratory Society (ERS). *Eur Heart J*. 2020;41(4):543-603.
  23. DeVos E, Jacobson L. Approach to adult patients with acute dyspnea. *Emerg Med Clin North Am*. 2016;34(1):129-49.
  24. Osheroff J, Teich J, Levick D, et al. (2012). *Improving Outcomes with Clinical Decision Support: An Implementer's Guide, 2nd Ed*. New York: HIMSS Publishing.
  25. Corazza F, Arpone M, Tardini G, et al. Effectiveness of a novel tablet application in reducing guideline deviations during pediatric cardiac arrest: a randomized clinical trial. *JAMA Netw Open*. 2023;6(8):e2327272.
  26. Sutton RT, Pincock D, Baumgart DC, et al. An overview of clinical decision support systems: benefits, risks, and strategies for success. *NPJ Digit Med*. 2020;3:17.
  27. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Medicine*. 2009;6(7):e1000097.
  28. Malmivaara A. Methodological considerations of the GRADE method. *Ann Med*. 2015;47(1):1-5.