

Does Prolonged Length of Stay in the Emergency Department Affect Outcome for Stroke Patients?

Minal Jain, MBBS, MPH*

Dushyant Damania, MBBS*

Anunaya R. Jain, MBBS, MCEM*

Abhijit R. Kanthala, MD*

Latha Ganti, MD, MS, MBA†

Babak S. Jahromi, MD, PhD, FRCSC*

* Department of Neurosurgery, University of Rochester Medical Center, Rochester, New York

† Departments of Emergency Medicine and Neurological Surgery, University of Florida College of Medicine, Gainesville, Florida

Supervising Section Editor: John Sarko, MD

Submission history: Submitted February 18, 2013; Revision received July 7, 2013; Accepted August 13, 2013

Electronically published April 15, 2014

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.2013.8.16186

Introduction: Conflicting data exist regarding the association between the length of stay (LOS) of critically ill patients in the emergency department (ED) and their subsequent outcome. However, such patients are an overall heterogeneous group, and we therefore sought to study the association between EDLOS and outcomes in a specific subgroup of critically ill patients, namely those with acute ischemic stroke/transient ischemic attack (AIS/TIA).

Methods: This was a retrospective review of adult patients with a discharge diagnosis of AIS/TIA presenting to an ED between July 2009 and February 2010. We collected demographics, EDLOS, arrival stroke severity (National Institutes of Health Stroke Scale - NIHSS), intravenous tissue plasminogen activator (IV tPA) use, functional outcome at discharge, discharge destination and hospital-LOS. We analyzed relationship between EDLOS, outcomes and discharge destination after controlling for confounders.

Results: 190 patients were included in the cohort. Median EDLOS was 332 minutes (Inter-Quartile Range -IQR: 250.3–557.8). There was a significant inverse linear association between EDLOS and hospital-LOS ($p=0.049$). Patients who received IV tPA had a shorter median EDLOS (238 minutes, IQR: 194–299) than patients who did not (median: 387 minutes, IQR: 285–588 minutes; $p<0.0001$). There was no significant association between EDLOS and poor outcome ($p=0.40$), discharge destination ($p=0.20$), or death ($p=0.44$). This remained true even after controlling for IV tPA use, NIHSS and hospital-LOS; and did not change even when analysis was restricted to AIS patients alone.

Conclusion: There was no significant association between prolonged EDLOS and outcome for AIS/TIA patients at our institution. We therefore suggest that EDLOS alone is an insufficient indicator of stroke care in the ED, and that the ED can provide appropriate acute care for AIS/TIA patients. [West J Emerg Med. 2014;15(3):267–275.]

INTRODUCTION

Acute ischemic stroke (AIS) is a clinical emergency that requires immediate and aggressive treatment, from prompt identification of symptoms in the emergency department (ED) through rapid investigations and treatment to hospital

discharge. Bench-marked timelines have been mandated for every step in the process,¹ leading to the creation of Primary Stroke Centers to standardize care delivery. In particular, as per the recommendations of the National Institute of Neurological Disorders and Stroke (NINDS) guidelines, transferring a stroke

patient to an inpatient setting should be achieved within 3 hours of arrival.² However, the limited availability of inpatient beds often results in access block, leading to an increase in the ED length of stay (EDLOS) for patients with stroke and transient ischemic attack (TIA).³⁻⁵ Studies have reported the median EDLOS for stroke patients to be approximately 5 hours.^{6,7} This can place an additional stress on the already overburdened ED, requiring personnel to not only provide emergent care but also devote time and resources for ongoing supportive care to stroke patients before transitioning to other areas of care within the hospital.

Numerous studies have evaluated the effects of prolonged LOS on the outcome of critically ill patients, with conflicting results.^{6,8-11} Rincon et al⁶ reported that critically ill stroke subjects waiting to be transitioned to intensive care unit (ICU) had a worse outcome when the EDLOS was > 5 hrs. Contrary to this, Saukkonen and his colleagues in a similar study with a diverse population of 1,675 critically ill patients concluded that mortality rate for patients with EDLOS beyond 24 hours was not significantly different than those with a shorter EDLOS.¹⁰ Elmer et al⁷ found no association between EDLOS and functional outcome in patients with hemorrhagic stroke. However, the effect of EDLOS on outcome of patients with ischemic stroke or TIA (representing more than 85% of stroke patients)¹² remains largely unexamined.

Rapid patient transfer from ED is believed to prevent crowding, provide more streamlined care to the patients, and promote efficient hospital operation and improved patient satisfaction.¹³ However, early discharge/transfer from the ED on the assumption that a prolonged LOS worsens outcome/mortality could promote inappropriately early transfers before the patient is stabilized and/or the recipient patient care unit is ready. Time criteria (such as that proposed by the NINDS) may be difficult to meet in busy tertiary care hospitals. Such perceived time pressures among providers can also lead to increased patient and family anxiety during this process.

Our aim in this study was to evaluate the effect of EDLOS on outcomes for patients presenting with AIS/TIA at our tertiary care academic center, after adjusting for common confounders such as initial stroke severity (National Institutes of Health Stroke Scale - NIHSS) and delivery of time-sensitive intravenous tissue plasminogen activator (IV tPA).

METHODS

Study Design and Data Sources

This was an institutional review board-approved retrospective record review of consecutive adult stroke/TIA patients presenting to our tertiary care center from July 2009 to February 2010. Our institution is a large academic primary stroke care center in upstate New York, with an annual turnover of nearly 600 acute ischemic stroke/TIA patients. Of these, approximately, half of them per year present within 12 hours of symptom onset. A flow chart is presented in Figure 1. Our protocol for management of AIS/TIA patients includes

investigations (blood glucose, serum electrolytes, ECG, cardiac ischemia markers, PT, INR, aPTT, oxygen saturation) along with multi-modal CT imaging (plain CT, CT angiography & perfusion CT imaging). We reviewed medical records to identify all adult patients (>18 years age) with a final discharge diagnosis of AIS/TIA who presented within 12 hours since last seen normal and had documented completion of per-protocol multimodal CT, initial stroke severity, time of ED registration, and time of departure from the ED. We collected data on patient demographics, time since last seen normal, EDLOS, initial stroke severity (NIHSS), past medical history, hospital LOS, functional outcome as measured by modified Rankin scale (mRS) at discharge, discharge destination [home, home with services, skilled nursing facility (SNF), or acute rehabilitation unit (ARU)] and mortality at discharge. The EDLOS was documented from time of registration (time of arrival at ED) to time of departure from the ED. Functional outcome at discharge was dichotomized and mRS ≥ 3 was considered as poor outcome. To improve uniformity of the cohort under investigation, we excluded from this study pediatric patients, patients with hemorrhagic strokes, and patients who did not receive all protocol mandated investigations or imaging studies.

Data Collection

To improve accuracy and reduce errors/inconsistencies in abstracted data, we followed the guidelines stated by Lowenstein et al¹⁴ during data collection process. All records were reviewed by a trained ED physician and a research fellow. Variables were explicitly defined and data was extracted from a standardized electronic medical record system. We resolved any discrepancy in the data by a common consensus between the abstractors. We calculated modified Rankin score at discharge based on occupational therapy and physical therapy notes. Inter-rater agreement was assessed for this parameter by having a sample of charts independently reviewed by the 2 data abstractors. Data was collected in MS Excel, then re-reviewed and cleaned before transfer into JMP statistical software by SAS Institute, Inc, for further analysis.

Statistical Analysis

We reported median and inter-quartile range (IQR) of continuous variables. We performed a bivariate Spearman's Rank correlation between EDLOS and age, time since last seen normal, initial stroke severity and hospital LOS. Spearman's ρ , 95% CI (Fisher's Z transformed) and p values were calculated for each correlation. We performed Wilcoxon/Kruskal Wallis tests to compare the association between EDLOS and administration of IV tPA and past history of comorbidities. A logistic fit was done to test the association between increasing EDLOS and poor outcome (mRS ≥ 3), death at discharge, discharge to SNF/death/ARU, and the association between hospital LOS and outcome. Finally, we performed a logistic regression to analyze the association between EDLOS on outcome, and discharge destination, after controlling for




Figure 1. Flow chart indicating the final cohort of patients. This flow chart indicates that our hospital managed about 380 acute ischemic stroke (AIS) or transient ischemic stroke (TIA) patients from July 2009 to February 2010, of which 55% presented within 12 hours of symptom onset. Of these 209 patients, 19 (9%) patients were ineligible as they were either in hospital admissions/ did not have emergency department length of stay documented, leaving a final cohort of 190 patients).

administration of IV tPA, initial stroke severity and hospital LOS. This was done because high initial stroke severity, non-thrombolysis and prolonged hospital LOS are usually associated with poor outcome^{15–18}. We repeated the above analysis for a sub cohort with only AIS patients, excluding patients with a final discharge diagnosis of TIA.

We performed all contingency analysis and multivariate analysis using JMP 9.0[®] and SAS 9.2[®] statistical software. The level of significant association was predetermined at $p < 0.05$ for all analyses.

RESULTS

Patient Demographics

A total of 209 patients were eligible for this study. Complete data were available for 190 (91%) of these patients (Figure 1). Of these, 138 (72.6%) patients had a discharge diagnosis of AIS and the remaining 52 (27.4%) had a discharge diagnosis of TIA. The characteristics of patients are described in Table 1.

EDLOS and Outcome in Patients with AIS or TIA

There were no statistically significant associations between EDLOS and age, time since last seen normal, or various stroke risk factors, such as past history of stroke/TIA ($p=0.25$), CAD ($p=0.18$), dyslipidemia ($p=0.63$), hypertension ($p=0.85$), diabetes mellitus ($p=0.22$), smoking ($p=0.09$) or atrial fibrillation (AF) ($p=0.19$) (Table 2). Lower EDLOS was associated with higher initial stroke severity ($p=0.0005$) and increased hospital LOS ($p=0.0497$). There was no significant association between EDLOS and functional outcome/mRS ≥ 3

(Figure 2) ($p=0.40$), discharge destination (Figure 3) (SNF/death/ARU, $p=0.20$) or death at discharge ($p=0.44$).

Patients with poor outcome had a significantly longer hospital LOS (median 6 days, IQR 4–9) as compared to those with a good outcome (median hospital LOS 2 days, IQR 1–4 days; $p < 0.0001$). Patients who received IV tPA had a significantly lower median EDLOS (238 minutes, IQR 194 – 299) as compared to patients who did not receive IV tPA (median EDLOS 387 minutes, IQR 285 – 588; $p < 0.0001$).

We performed a logistic regression to assess the effect of EDLOS on outcomes (mRS at discharge ≥ 3 and discharge destination other than home) after adjusting for the above confounders - IV tPA, initial stroke severity and hospital LOS. This odds ratio and 95% CI are shown in Table 3. After controlling for the above variables, the association between EDLOS and poor outcome ($p=0.695$), and discharge destination other than home ($p=0.5019$) remained statistically insignificant.

EDLOS and Outcome in Patients with AIS Alone

Restricting the analysis to only AIS patients revealed similar results, with no statistically significant associations between EDLOS and age, and stroke risk factors [past history of stroke/TIA ($p=0.36$), CAD ($p=0.44$), dyslipidemia ($p=0.95$), hypertension ($p=0.46$), diabetes mellitus ($p=0.16$), smoking ($p=0.23$) and AF ($p=0.28$)] (Table 4). There was a significant association between lower EDLOS and higher initial stroke severity ($p=0.001$). Although the trend between EDLOS and hospital LOS was similar as for all AIS/TIA patients, the association did not remain significant for the AIS-only cohort

Table 1. Characteristics of patients presenting with acute ischemic stroke or transient ischemic attack.

Variable	AIS/TIA (n=190) (n, %)	AIS (n=138) (n, %)
Age (Median, IQR)	71 (60–80)	69 (60–82)
Female gender	88 (46.3)	46 (42.6)
Time since last seen normal (Median, IQR)	138.5 (69–367.5)	155 (68 – 528)
NIHSS (Median, IQR)	4 (1–11)	6 (3–13)
Past history of comorbidities		
Stroke/TIA	61 (32.1)	41 (29.7)
Coronary artery disease	45 (23.7)	37 (26.8)
Dyslipidemia	85 (44.7)	58 (42.0)
Hypertension	142 (74.7)	106 (76.8)
Diabetes mellitus	45 (23.7)	39 (28.3)
Smoking	63 (33.2)	52 (37.7)
Atrial fibrillation	37 (19.5)	26 (18.8)
EDLOS (Median, IQR)	332 (250.3–557.8)	315 (232.5–561.8)
Thrombolysis	35 (18.4)	33 (23.9)
Poor outcome on discharge (mRS \geq 3)	56 (29.5)	55 (39.9)
Hospital LOS	3 (2–6)	4 (2–7.3)
Discharge destination		
Home	92 (48.4)	53 (38.4)
Home with services	24 (12.6)	18 (13.0)
SNF	41 (21.6)	36 (26.1)
ARU	19 (10.0)	17 (12.3)
Mortality at discharge	14 (7.4)	14 (10.1)

AIS, acute ischemic stroke; TIA, transient ischemic attack; IQR, interquartile range; NIHSS, National Institutes of Health stroke scale; ED, emergency department; LOS, length of stay; mRS, modified Rankin score; SNF, skilled nursing facility; ARU, acute rehabilitation unit

($p=0.13$). There was no association between EDLOS and poor outcome or discharge destination in this more narrowly defined cohort (Figures 4–5). Specifically, we found no statistically significant association between EDLOS and poor outcome

(mRS \geq 3) ($p=0.41$), discharge to SNF/ARU/death ($p=0.20$) or death at discharge ($p=0.44$) in AIS patients.

Patients with a poor outcome had a significantly longer hospital LOS (median 6 days, IQR 4–9) as compared to patients

Table 2. Emergency department length of stay and outcome in patients with acute ischemic stroke or transient ischemic attack.

Variable	Association with EDLOS	95% CI	p value
Age	$\rho = -0.12$	-0.26 to 0.02	0.10
Time since last seen normal	$\rho = 0.14$	-0.002 to 0.276	0.06
NIHSS	$\rho = -0.25$	-0.38 to -0.11	0.0005*
Hospital LOS	$\rho = -0.14$	-0.28 to 0.002	0.0497*
Poor outcome (mRS \geq 3)	OR = 0.99	0.99 to 1.00	0.40
Discharge destination			
Home/Home with services	OR = 1.00	0.99 to 1.00	0.20
SNF/death	OR = 0.99	0.99 to 1.00	0.54
ARU	OR = 0.99	0.99 to 1.00	0.19
Mortality at discharge	OR = 0.99	0.99 to 1.00	0.44

mRS, modified Rankin score, SNF, skilled nursing facility, ARU, acute rehabilitation unit, OR, odds ratio, CI, confidence intervals; ρ , Spearman's correlation coefficient

* A statistically significant inverse correlation was found between EDLOS and only initial stroke severity (NIHSS) and hospital LOS for patients presenting with AIS or TIA.




Figure 2. Emergency department length of stay and outcome in patients with acute ischemic stroke or transient ischemic attack. There was no significant difference in median emergency department length of stay of patients with poor outcome (modified Rankin score [mRS] ≥ 3) (307.5 min, interquartile range [IQR] 228.3–492.3) versus patients with good outcome (337 min, IQR 257.8–577.3; $p=0.15$).

with good outcome (median hospital LOS 3 days, IQR 2–5 days; $p<0.0001$). In the AIS group, patients who received IV tPA ($n=33$) had a significantly shorter median EDLOS (238 minutes, IQR 195.5 – 292) as compared to patients who did not receive IV tPA (median EDLOS 399 minutes, IQR 280.5 – 623; $p<0.0001$).

After adjusting for potential confounders (differences in IV tPA administration rates, initial stroke severity and hospital LOS), we did not find any statistically significant association between EDLOS and poor outcome ($p=0.98$), or discharge destination other than home ($p=0.70$) in patients with AIS alone. This is shown in Table 5.

DISCUSSION

We chose to focus our study on patients with AIS/TIA because of strict guidelines and compliance mandated by the Joint Commission and the State Department of Health for Primary Stroke Centers with respect to this patient population. Our data suggest that there is no significant impact of increased EDLOS on outcome of AIS/TIA patients, even after controlling for presenting stroke severity, IV tPA delivery and total hospital LOS at our tertiary care academic center. In contrast, when examining all stroke subtypes, Rincon et al⁶ found that of 519 stroke patients, EDLOS adversely affected outcome in 75 patients who were admitted to a neurointensive care unit. Despite adjusting for presenting NIHSS, we did not find this association in our subpopulation of AIS/TIA patients. Our data parallel the findings by Elmer et al,⁷ which showed no

association between EDLOS and functional outcome in patients with hemorrhagic stroke. In a broader population, Varon et al¹⁹ studied 50 consecutive patients admitted to ICU from the ED and found similar survival rates in patients who received critical care procedures in the ED versus patients who received similar interventions in an ICU setting.

We found no difference in mortality or discharge destination with respect to EDLOS in our patients. Similarly, Saukkonen et al¹⁰ looked at a broader critically ill patient population, including patients with neurological diseases, and reported that EDLOS (>24 hours) was not associated with hospital mortality or quality of life at 6 months post discharge. Using a similar >24 hour cut-off point for EDLOS, a study of 443 critically ill patients in 2001–2002 found that mortality rate for stroke patients was unaffected by EDLOS.¹¹ However, at our institute (and many others in the U.S. today) patient stays in the ED >24 hours are considered as “sentinel events” (only 5.8% subjects in our study population had EDLOS >24 hours), with strict protocols designed to prevent such an occurrence. The results of prior studies may therefore be less applicable in the environment that exists today. Data from our study suggest that even with current protocols mandating early transfer to in-patient units or ICU, increased EDLOS does not affect functional outcomes or mortality in AIS/TIA patients.

With changing Medicare and Medicaid reimbursements, there has been an increasing focus on reduction of overall length of stay in the hospital. Studies have shown that




Figure 3. Emergency department length of stay and discharge destination in patients with acute ischemic stroke (AIS) or transient ischemic attack (TIA). There was no significant association between EDLOS and discharge destination, with median EDLOS (IQR) of 337 (254–574), 312 (253.5–483.3), and 301 (228.3–519.3) minutes for discharge to home/home with services, acute rehabilitation unit (ARU) or skilled nursing facility (SNF)/death, respectively.

Table 3. Logistic regression output for association between emergency department length of stay and outcome in acute ischemic stroke and transient ischemic attack patients.

Variable	OR	95% CI	P
Poor outcome (mRS ≥ 3)			
EDLOS	1	0.999 – 1.000	0.8179
Initial stroke severity (NIHSS)	1.20	1.13 – 1.29	<0.0001
Thrombolysis	0.61	0.21 – 1.63	0.3392
Hospital LOS	1.02	0.98 – 1.10	0.4178
Discharge destination other than home			
EDLOS	1	0.999 – 1.000	0.5019
Initial stroke severity (NIHSS)	1.19	1.12 – 1.28	<0.0001
Thrombolysis	0.68	0.24 – 1.82	0.4546
Hospital LOS	1.17	1.05 – 1.32	0.006

mRS, modified Rankin score, NIHSS: National Institutes of Health stroke scale; OR, odds ratio; CI, confidence intervals

prolonged EDLOS contributes to longer hospital LOS, which is considered a surrogate marker for poor outcome or inefficient care.²⁰ In contrast, we found that a shorter EDLOS at our institution was paradoxically associated with increased hospital LOS, although this may be due to the association between shorter EDLOS and higher initial stroke severity in our cohort. Overall, we found no significant association between EDLOS and poor outcome, suggesting that delivery of care in the ED can be efficient and congruent with the care delivered in subsequent hospital settings for this patient subgroup.

Some authors have explained the association of long EDLOS to poor outcome by linking it to triage preferences, suggesting that patients with good prognosis might be treated, triaged and/or transferred early from the ED at the expense of patients with worse prognosis and/or comfort care order.⁹ By only analyzing AIS/TIA patients who had uniformly completed all investigations and imaging studies, as well as adjusting for

time dependent IV tPA delivery and initial stroke severity, we attempted to avoid such limitations in our analysis.

Our data shows that EDLOS does not affect outcome, but does not indicate why. It is possible that outcome after AIS/TIA is mostly dependent upon factors such as initial stroke severity, thrombolysis and time from onset, none of which are affected by EDLOS. (Thrombolysis usually occurs within the ED setting). Yet even after adjusting for the above factors, there was no significant association between EDLOS and outcome. Differences in EDLOS may reflect practical availability of beds elsewhere in the hospital, and overall minor variations in EDLOS (measured in hours rather than days) may be unlikely to affect the overall outcome of AIS/TIA. This of course assumes that the ED is capable of providing the appropriate acute/critical care needed within the first few hours of AIS/TIA, which we believe holds at most institutions, including our own. The above reasons, while speculative, do call into question the

Table 4. Emergency department length of stay and outcome in patients with acute ischemic stroke alone.

Variable	Association with EDLOS	95% CI	p value
Age	$\rho = -0.11$	-0.27 to 0.06	0.18
Time since last seen normal	$\rho = 0.19$	0.02 to 0.35	0.03*
NIHSS	$\rho = -0.27$	-0.42 to -0.11	0.001*
Hospital LOS	$\rho = -0.13$	-0.29 to 0.04	0.13
Poor outcome (mRS ≥ 3)	OR = 0.999	0.998 to 1.00	0.41
Discharge destination			
Home/Home with services	OR = 1.00	0.99 to 1.00	0.20
SNF/death	OR = 0.99	0.99 to 1.00	0.59
ARU	OR = 0.99	0.99 to 1.00	0.19
Mortality at discharge	OR = 0.999	0.996 to 1.00	0.44

NIHSS, National Institutes of Health stroke scale; mRS, modified Rankin score; SNF, skilled nursing facility; ARU, acute rehabilitation unit; OR, odds ratio; CI, confidence intervals; ρ , Spearman's correlation coefficient

* Considering the subgroup of patients presenting with AIS alone, a statistically significant inverse correlation was found between EDLOS and only initial stroke severity (NIHSS).




Figure 4. Emergency department length of stay and outcome in patients with acute ischemic stroke (AIS) alone. There was no significant difference in median EDLOS of AIS patients with poor outcome (modified Rankin score [mRS] ≥ 3) (309 min, interquartile range [IQR] 229–499) versus patients with good outcome (316 min, IQR 233–592; $p=0.38$).

focus on EDLOS as a measure of quality of care, if no direct correlation can be found between EDLOS and eventual patient outcomes.

We emphasize that our study does not provide data regarding the benefits of transferring AIS/TIA patients from ED to higher levels of care, and believe that transferring appropriate patients early helps operational efficiency in the ED by decreasing crowding and resource utilization. We only suggest that focusing on EDLOS as the sole or key indicator of quality of stroke care in the ED may be inaccurate, as increased EDLOS alone does not seem to bear upon outcomes of patients with AIS/TIA.

LIMITATIONS

There were several limitations identified during our study. The primary outcome was mRS at discharge. Some have argued that mRS and death may not be the best outcome measures to study the effect of EDLOS, and have proposed looking at other surrogate markers, such as quality adherence and time to antibiotics, as they are less patient specific.^{22,23} However, the inability of surrogate markers to produce consistent results is well documented, and we felt it more reasonable to use mRS, which is a routinely used functional outcome score for stroke patients.^{24,25} In addition to this, we reported outcomes at discharge. While prospective large-scale stroke trials have typically assessed outcome at 3 months, we were unable to do so due to the retrospective nature of our study and the

availability of data within the existing records. We hope to address this limitation in a future prospective study.

We were also not able to do a formal power calculation for this study, since this was not a prospective comparison of 2 groups with a pre-specified EDLOS cutoff. Rather, our study retrospectively analyzed patient outcome in relation to EDLOS as a continuous variable, and used multivariate regression to adjust for confounders in this analysis.

Our study was conducted at a single center and the conclusions should be cautiously generalized to other hospitals due to variability in the geographic location, level of care offered, patient management and outcome metrics of other institutes. The study was conducted at a tertiary academic teaching center, which is a certified Primary Stroke Center, with a 24x7 availability of in-house neurologists/ neurosurgeons providing immediate care to stroke patients. Thus, there is simultaneous involvement of multiple specialties – ED, Neurology, and Neurosurgery within the ED itself to coordinate and deliver care. This could also limit the generalizability of our results to other institutions without a similar structural and functional setup.

CONCLUSION

Our study concluded that there was no significant association between ED length of stay and mRS, mortality, or




Figure 5. Emergency department length of stay (EDLOS) and discharge destination in patients with acute ischemic stroke (AIS) alone. There was no significant association between EDLOS and discharge destination, with median EDLOS (IQR) of 317.5 minutes (229–600.3), 312 minutes (253.5–483.3), and 301 (228.8–535.5) minutes for discharge to home/home with services, acute rehabilitation unit (ARU), or skilled nursing facility (SNF)/death, respectively.

Table 5. Logistic regression output for association between emergency department length of stay and outcome in acute ischemic stroke patients only.

Variable	OR	95% CI	P
Poor outcome (mRS ≥ 3)			
EDLOS	1	0.99 – 1.003	0.9839
Initial stroke severity (NIHSS)	1.18	1.11 – 1.27	<0.0001*
Thrombolysis	0.63	0.23 – 1.65	0.3581
Hospital LOS	1.01	0.97 – 1.1	0.6176
Discharge destination other than home			
EDLOS	1	0.999 – 1.004	0.7001
Initial stroke severity (NIHSS)	1.18	1.10 – 1.28	<0.0001*
Thrombolysis	0.71	0.25 – 1.92	0.5076
Hospital LOS	1.10	1.00 – 1.25	0.094

mRS, modified Rankin score, NIHSS, National Institutes of Health stroke scale; SNF, skilled nursing facility, ARU, acute rehabilitation unit, OR, odds ratio; CI, confidence intervals

discharge destination for acute ischemic stroke or TIA patients presenting to a tertiary care academic ED.

Address for Correspondence: Babak S. Jahromi, MD, PhD, FRCSC. University of Rochester Medical Center, School of Medicine, 601 Elmwood Ave. Box 670, Rochester, NY 14642. Email: babak_jahromi@urmc.rochester.edu.

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.

REFERENCES

- Adams HP, Jr., del Zoppo G, Alberts MJ, et al. Guidelines for the early management of adults with ischemic stroke: a guideline from the American Heart Association/American Stroke Association Stroke Council, Clinical Cardiology Council, Cardiovascular Radiology and Intervention Council, and the Atherosclerotic Peripheral Vascular Disease and Quality of Care Outcomes in Research Interdisciplinary Working Groups: the American Academy of Neurology affirms the value of this guideline as an educational tool for neurologists. *Stroke* 2007;38:1655–1711
- National Institute of Neurological Disorders and Stroke, National Institute of Health. NINDS: Stroke Proceedings: Executive Summary. *Proceedings of a National Symposium on Rapid Identification and Treatment of Acute Stroke* Bethesda, MD; 2011
- Fromm RE, Jr., Gibbs LR, McCallum WG, et al. Critical care in the emergency department: a time-based study. *Crit Care Med* 1993;21:970–976
- Nelson M, Waldrop RD, Jones J, et al. Critical care provided in an urban emergency department. *Am J of Emerg Med* 1998;16:56–59
- United States General Accounting Office. Hospital emergency departments. Crowded conditions vary among hospitals and communities.; 2003
- Rincon F, Mayer SA, Rivolta J, et al. Impact of delayed transfer of critically ill stroke patients from the Emergency Department to the Neuro-ICU. *Neurocrit Care* 2010;13:75–81
- Elmer J, Pallin DJ, Liu S, et al. Prolonged Emergency Department length of stay is not associated with worse outcomes in patients with intracerebral hemorrhage. *Neurocrit Care* 2011
- Rapoport J, Teres D, Lemeshow S, et al. Timing of intensive care unit admission in relation to ICU outcome. *Crit Care Med* 1990;18:1231–1235
- Chalfin DB, Trzeciak S, Likourezos A, et al. Impact of delayed transfer of critically ill patients from the emergency department to the intensive care unit. *Crit Care Med* 2007;35:1477–1483
- Saukkonen KA, Varpula M, Rasanen P, et al. The effect of emergency department delay on outcome in critically ill medical patients: evaluation using hospital mortality and quality of life at 6 months. *Jl of Intern Med* 2006;260:586–591
- Tilluckdharry L, Tickoo S, Amoateng-Adjepong Y, et al. Outcomes of critically ill patients. *Am J of Emerg Med* 2005;23:336–339
- Roger VL, Go AS, Lloyd-Jones DM, et al. Heart disease and stroke statistics–2012 update: a report from the American Heart Association. *Circulation* 2012;125:e2–e220
- Stead LG, Jain A, Decker WW. Emergency department over-crowding: a global perspective. *Int J of Emerg Med* 2009;2:133–134
- Lowenstein SR. Medical record reviews in emergency medicine: the blessing and the curse. *Ann Emerg Med* 2005;45:452–455
- Tissue plasminogen activator for acute ischemic stroke. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. *The New England Journal of Medicine* 1995;333:1581–1587
- Adams HP, Jr., Davis PH, Leira EC, et al. Baseline NIH Stroke Scale score strongly predicts outcome after stroke: A report of the Trial of Org 10172 in Acute Stroke Treatment (TOAST). *Neurology* 1999;53:126–131

17. Wong DT, Gomez M, McGuire GP, et al. Utilization of intensive care unit days in a Canadian medical-surgical intensive care unit. *Crit Care Med* 1999;27:1319–1324
18. Fakhry SM, Kercher KW, Rutledge R. Survival, quality of life, and charges in critically ill surgical patients requiring prolonged ICU stays. *J Trauma* 1996;41:999–1007
19. Varon J, Fromm RE, Jr., Levine RL. Emergency department procedures and length of stay for critically ill medical patients. *Ann Emerg Med* 1994; 23:546–549
20. Krochmal P, Riley TA. Increased health care costs associated with ED overcrowding. *Am J of Emerg Med* 1994;12:265–266
21. Heidenreich PA, Trogon JG, Khavjou OA, et al. Forecasting the future of cardiovascular disease in the United States: a policy statement from the American Heart Association. *Circulation* 2011;123:933–944
22. Pines JM. Measuring antibiotic timing for pneumonia in the emergency department: another nail in the coffin. *Ann Emerg Med* 2007;49:561–563
23. Diercks DB, Roe MT, Chen AY, et al. Prolonged emergency department stays of non-ST-segment-elevation myocardial infarction patients are associated with worse adherence to the American College of Cardiology/American Heart Association guidelines for management and increased adverse events. *Ann Emerg Med* 2007;50:489–496
24. Wachter RM, Flanders SA, Fee C, et al. Public reporting of antibiotic timing in patients with pneumonia: lessons from a flawed performance measure. *Ann Intern Med* 2008;149:29–32
25. Fonarow GC, Abraham WT, Albert NM, et al. Association between performance measures and clinical outcomes for patients hospitalized with heart failure. *JAMA* 2007;297:61–70