

Adolescent with Foot Pain

Nicole Mansfield, MD
Sarab Sodhi, MD
Richard Pescatore, DO
Andrew Nyce, MD

Cooper University Hospital, Department of Emergency Medicine, Camden,
New Jersey

Section Editor: Rick A. McPheeters, DO

Submission history: Submitted February 20, 2016; Revision received May 21, 2016; Accepted June 7, 2016

Electronically published July 19, 2016

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.2016.6.30142

[West J Emerg Med. 2016;17(5)617-618.]

A previously healthy 14-year-old female presented to the emergency department because of a left foot injury. She reported “twisting” her foot while walking. She complained of left foot pain, but was able to ambulate with a limp. Physical exam was significant for soft tissue edema and isolated tenderness to the dorsum of the left foot in the region of the cuboid. She had full range of motion at the ankle joint without pain or tenderness and was neurovascularly intact. Radiographs were obtained and revealed cuboid subluxation (Figure).

DIAGNOSIS

Cuboid subluxation. Cuboid subluxation is generally



Figure. Anterior-posterior (left) and oblique (right) radiograph of the left foot with lateral displacement of the cuboid relative to the base of the fourth metatarsal (arrows).

diagnosed clinically, based on history and physical exam, and definitive radiographic evidence is rare. Injury presentation is varied, but case reports often describe pain isolated over the cuboid following a forceful inversion of the foot while bearing weight. Physical exam typically reveals tenderness over the cuboid. Because of normal variations between the cuboid and its surrounding structures, traditional radiographs, computerized tomography, and magnetic resonance imaging are generally non-diagnostic for this injury.¹⁻⁸ Additionally, midtarsal joint abnormalities that cause pain while weight bearing may be missed on non-weight bearing films.² Imaging, however, is often ordered even when suspicion for cuboid subluxation is high to rule out other causes of foot pain.

In the case image above (Figure), there is lateral displacement of the cuboid relative to the base of the fourth metatarsal (arrows). A good rule-of-thumb and highly suggestive radiographic finding of cuboid subluxation is a disturbed relationship between the medial-cuboid and its articulation with the fourth metatarsal.

Treatment may require manual reduction and responds well to conservative physiotherapy management. Our patient was fitted with a controlled ankle motion (CAM) walking boot and was given crutches. She was referred to the orthopedic surgery clinic for further management.

Address for Correspondence: Nicole Mansfield, MD, Cooper University Hospital, Department of Emergency Medicine, One Cooper Plaza, Kelemen 152 Camden, New Jersey 08103. Email: nickyjmansfield2@gmail.com.

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.

Copyright: © 2016 Mansfield et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. Adams E and Madden C. Cuboid subluxation: a case study and review of the literature. *Curr Sports Med Rep.* 2009;8(6):300-7.
2. Durall CJ. Examination and treatment of cuboid syndrome: a literature review. *Sports Health.* 2011;3(6):514-9.
3. Jennings J and Davies GJ. Treatment of cuboid syndrome secondary to lateral ankle sprains: a case series. *J Orthop Sports Phys Ther.* 2005;35(7):409-15.
4. Marshall P and Hamilton WG. Cuboid subluxation in ballet dancers. *Am J Sports Med.* 1992;20(2):169-75.
5. Mooney M and Maffey-Ward L. Cuboid plantar and dorsal subluxations: assessment and treatment. *J Orthop Sports Phys Ther.* 1994;20(4):220-6.
6. Newell SG and Woodle A. Cuboid Syndrome. *Physician and Sports Medicine*1981:71-6.
7. Patterson SM. Cuboid syndrome: a review of the literature. *J Sports Sci Med.* 2006;5(4):597-606.
8. Zirm R. Cuboid Syndrome. *Podiatry Institute.* 1997.